

Noncommutative general relativity

Article (Published Version)

Calmet, Xavier and Kobakhidze, Archil (2005) Noncommutative general relativity. *Physical Review D*, 72 (4). ISSN 1550-7998

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/31376/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Noncommutative general relativityXavier Calmet^{1,*} and Archil Kobakhidze^{2,3,†}¹*Service de Physique Théorique, Université Libre de Bruxelles, CP225, Boulevard du Triomphe (Campus plaine), B-1050 Brussels, Belgium*²*Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA*³*Andronikashvili Institute of Physics, 0177 Tbilisi, Georgia*

(Received 28 June 2005; published 11 August 2005)

We define a theory of noncommutative general relativity for canonical noncommutative spaces. We find a subclass of general coordinate transformations acting on canonical noncommutative spacetimes to be volume-preserving transformations. Local Lorentz invariance is treated as a gauge theory with the spin connection field taken in the $so(3,1)$ enveloping algebra. The resulting theory appears to be a noncommutative extension of the unimodular theory of gravitation. We compute the leading order noncommutative correction to the action and derive the noncommutative correction to the equations of motion of the weak gravitation field.

DOI: [10.1103/PhysRevD.72.045010](https://doi.org/10.1103/PhysRevD.72.045010)

PACS numbers: 11.10.Nx, 04.20.-q

I. INTRODUCTION

General Relativity [1] is a very successful theory when it comes to describe macroscopic effects of gravitation. However, it is widely believed that a unification of Quantum Mechanics and General Relativity requires a short distance modification of spacetime. It can be shown that classical General Relativity considered together with Quantum Mechanics implies the existence of a fundamental length [2]. A class of models that incorporate the notion of a fundamental length in gauge theories are gauge theories formulated on noncommutative spaces.

It is a challenge to formulate General Relativity on noncommutative spaces and there are thus different approaches in the literature. In [3] for example a deformation of Einstein's gravity was studied using a construction based on gauging the noncommutative $SO(4,1)$ de Sitter group and the Seiberg-Witten map with subsequent contraction to $ISO(3,1)$. Most recently another construction of a noncommutative gravitational theory [4] was proposed based on a twisted Poincaré algebra [5,6]. These approaches, although mathematically consistent, are not minimal formulations of Einstein's General Relativity on noncommutative spaces. The main problem in formulating a theory of gravity on noncommutative manifolds is that it is difficult to implement symmetries such as general coordinate covariance and local Lorentz invariance and to define derivatives which are torsion free and satisfy the metricity condition.

Similar obstacles appear in constructing models of particle physics on flat spacetime with canonical noncommutativity defined by the algebra $[\hat{x}^a, \hat{x}^b] = i\theta^{ab}$ (θ^{ab} is constant and antisymmetric). Indeed, it turns out to be rather difficult to implement most symmetries particle physicists are so familiar with. In particular, Lorentz in-

variance is explicitly violated by the noncommutative algebra. However, it has been shown in [7] that there is another exact symmetry, the noncommutative Lorentz invariance, based on the usual Lorentz algebra $so(3,1)$ which is undeformed. Another example is the implementation of noncommutative local gauge theories. A formulation of noncommutative gauge theories within the enveloping algebra approach has been proposed in [8]. The fields taken in the enveloping algebra are expanded in term of a series in θ . Each of the terms of this series is however a function of the classical variables [9,10]. The number of degrees of freedom is thus finite and the same as in the corresponding commutative gauge theory.

In this work, based partially on the above achievements in implementing symmetries on flat noncommutative spacetimes, we would like to propose a theory of General Relativity on curved spacetimes with canonical noncommutativity. We shall use the tetrad approach to General Relativity. This formalism applied to noncommutative General Relativity allows one to follow closely the usual construction of noncommutative gauge theories. This requires one to implement two gauge symmetries: local Lorentz transformations which can be seen as a local gauge theory based on the algebra $so(3,1)$ for the spin connection field and general coordinate transformations which are inhomogeneous translations with the tetrad as a gauge field.

The gauging of noncommutative $so(3,1)$ algebra is only possible if the corresponding gauge field, the spin connection, is assumed to be in the enveloping algebra. Hence, to implement local Lorentz invariance we follow the approach developed in [8]. The invariance under the general coordinate transformations, however, is explicitly violated by the canonical noncommutative algebra. Nevertheless, we find a restricted class of coordinate transformations which preserves the canonical structure. It turns out that these transformations correspond to volume-preserving diffeomorphisms. Volume-preserving transformations

*Electronic address: xcalmet@ulb.ac.be†Electronic address: kobakhid@physics.unc.edu

have been previously discussed in [11,12] in different physical frameworks. Thus, the basic new ideas for constructing a theory of noncommutative General Relativity are to formulate local Lorentz invariance by gauging $so(3,1)$ within the enveloping algebra approach and to introduce volume-preserving coordinate transformations in place of general coordinate transformations, which is indeed an exact symmetry of the canonical noncommutative spacetime.

II. NONCOMMUTATIVE GENERAL RELATIVITY

Let us start from a noncommutative spacetime and assume that the coordinates fulfill canonical commutation relations:

$$[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu}. \quad (1)$$

Obviously, the commutator (1) explicitly violates general coordinate covariance since $\theta^{\mu\nu}$ is constant in all reference frames. However, we can identify a subclass of general coordinate transformations,

$$\hat{x}^{\mu'} = \hat{x}^\mu + \hat{\xi}^\mu(\hat{x}), \quad (2)$$

which are compatible with the algebra given by (1). The hat on the function $\hat{\xi}^\mu(\hat{x})$ indicates that it is in the enveloping algebra. Under the change of coordinates (2), the commutator (1) transforms as

$$\begin{aligned} [\hat{x}^{\mu'}, \hat{x}^{\nu'}] &= \hat{x}^{\mu'}\hat{x}^{\nu'} - \hat{x}^{\nu'}\hat{x}^{\mu'} \\ &= i\theta^{\mu\nu} + [\hat{x}^\mu, \hat{\xi}^\nu] + [\hat{\xi}^\mu, \hat{x}^\nu] + \mathcal{O}(\hat{\xi}^2). \end{aligned} \quad (3)$$

Requiring that θ remains constant yields the following partial differential equations:

$$\theta^{\mu\alpha}\hat{\partial}_\alpha\hat{\xi}^\nu(\hat{x}) = \theta^{\nu\beta}\hat{\partial}_\beta\hat{\xi}^\mu(\hat{x}). \quad (4)$$

A nontrivial solution to this condition can be easily found:

$$\hat{\xi}^\mu(\hat{x}) = \theta^{\mu\nu}\hat{\partial}_\nu\hat{f}(\hat{x}), \quad (5)$$

where $\hat{f}(\hat{x})$ is an arbitrary field. This noncommutative general coordinate transformation corresponds to the following classical transformation: $\hat{\xi}^\mu(x) = \theta^{\mu\nu}\partial_\nu\hat{f}(x)$. The Jacobian of this restricted coordinate transformation is equal to 1, meaning that the volume element is invariant: $d^4x' = d^4x$. The version of General Relativity based on volume-preserving diffeomorphism is known as the unimodular theory of gravitation [13]. Thus we come to the conclusion that symmetries of canonical noncommutative spacetime naturally lead to the noncommutative version of unimodular gravity.

Now we need to implement two gauge symmetries mentioned above. A noncommutative gauge transformation $\hat{\Lambda}(\hat{x})$ valued in the $iso(3,1)$ Lie algebra can be decomposed using the generators of the inhomogeneous translations $p_\mu = -i\partial_\mu$, which are anti-Hermitian, and the generators of the local Lorentz algebra $so(3,1)$ Σ_{ab} , which are Hermitian. One finds

$$\hat{\Lambda}(\hat{x}) = \hat{\xi}(\hat{x}) + \hat{\Lambda}(\hat{x}) = \hat{\xi}^\mu(\hat{x})p_\mu + \frac{1}{2}\hat{\lambda}^{ab}(\hat{x})\Sigma_{ab}, \quad (6)$$

where $\hat{\xi}^\mu$ is subject to the constraint (5). Note that p_μ acts on the coordinates and functions, including $\hat{\lambda}^{ab}$, and a, b, \dots run over the tangent space indices. As in [7], the algebra of generators is undeformed. It is easy to verify that the commutator of two noncommutative gauge transformations $[\hat{\Lambda}_1(\hat{x}), \hat{\Lambda}_2(\hat{x})]$ is in general not a noncommutative gauge transformation if the transformations are Lie algebra valued. As in the Yang-Mills case, the solution is to assume that the noncommutative gauge transformations are in the enveloping algebra. Let us introduce a noncommutative vector potential which corresponds to the noncommutative gauge transformation (6)

$$\hat{A}_a(\hat{x}) = (\hat{D}_a) = i\hat{E}_a^\mu(\hat{x})p_\mu + \frac{i}{2}\hat{\omega}(\hat{x})_a^{bc}\Sigma_{bc}, \quad (7)$$

where $\hat{E}_a^\mu(\hat{x})$ are the components of the noncommutative tetrad $\hat{E}_a(\hat{x})$, i.e. the gauge fields corresponding to general coordinate transformations, and $\hat{\omega}(\hat{x})_a^{bc}$ are the spin connections fields associated with local Lorentz invariance. Note that $\hat{A}_a(\hat{x})$ plays a dual role. It can be viewed as a covariant derivative as well. It is also worth noticing that $\hat{E}_a = \hat{E}_a^\mu\hat{\partial}_\mu = \hat{\partial}_a$, which implies that the noncommutative tetrad is mapped trivially on the commutative one: $\hat{E}_a = e_a$ to all orders in θ .

Let us now assume that the gauge transformations and the spin connection field are in the enveloping algebra:

$$\hat{\Lambda} = \Lambda(x) + \Lambda^{(1)}(x, \omega_a) + \mathcal{O}(\theta^2), \quad (8)$$

and

$$\hat{\omega}_a = \omega_a(x) + \omega_a^{(1)}(x, \omega_a) + \mathcal{O}(\theta^2), \quad (9)$$

respectively, with $\Lambda(x) = \xi^\mu(x)p_\mu + \frac{1}{2}\lambda^{ab}(x)\Sigma_{ab}$ and $\omega_a(x) = \frac{1}{2}\omega_a^{bc}\Sigma_{bc}$. We require that the commutator of two noncommutative gauge transformations with $\hat{\Lambda}_1$ and $\hat{\Lambda}_2$ be a gauge transformation $\hat{\Lambda} \underset{\Lambda_1 \times \Lambda_2}{\frown} :$

$$(\hat{\delta}_{\hat{\Lambda}_1}\hat{\delta}_{\hat{\Lambda}_2} - \hat{\delta}_{\hat{\Lambda}_2}\hat{\delta}_{\hat{\Lambda}_1}) \star \hat{\phi}(x) = (i\hat{\delta}_{\hat{\Lambda}_1}\hat{\Lambda}_2[\omega_a] - i\hat{\delta}_{\hat{\Lambda}_2}\hat{\Lambda}_1[\omega_a] + [\hat{\Lambda}_1[\omega_a]; \hat{\Lambda}_2[\omega_a]) \star \hat{\phi}(x) = \hat{\Lambda} \underset{\Lambda_1 \times \Lambda_2}{\frown} \star \hat{\phi}(x). \quad (10)$$

One finds as usual

$$[\Lambda_1, \Lambda_2] = i\Lambda_{\Lambda_1 \times \Lambda_2} \quad (11)$$

in the zeroth order in θ and

$$i\delta_{\Lambda_1}\Lambda_2^{(1)} - i\delta_{\Lambda_2}\Lambda_1^{(1)} + i\theta^{ab}\{\partial_a\Lambda_1, \partial_b\Lambda_2\} + [\Lambda_1, \Lambda_2^{(1)}] - [\Lambda_2, \Lambda_1^{(1)}] = \Lambda_{\Lambda_1 \times \Lambda_2}^{(1)} \quad (12)$$

in the leading order in θ . A solution to this consistency equation is

$$\Lambda_1^{(1)} = \frac{1}{4}\theta^{ab}\{\partial_a\Lambda_1, \omega_b\} \quad (13)$$

and analogously for $\Lambda_2^{(1)}$ and $\Lambda_{\Lambda_1 \times \Lambda_2}^{(1)}$ and where we have used $\theta^{ab} = \theta^{\mu\nu}e_\mu^a e_\nu^b$ and $\partial_a = e_a^\mu \partial_\mu$. Note that the consistency condition is derived in the leading order in θ and $\xi(x)$. However, since $\xi(x)$ is itself proportional to θ , the relevant part of the volume-preserving diffeomorphism transformation is trivial. In other words, the terms proportional to $\xi(x)$ can be dropped in Eq. (13) and it actually determines $\lambda^{(1)}$ which is the first nontrivial term in the Seiberg-Witten map for the $\text{so}(3,1)$ gauge transformation.

The consistency condition for the spin connection is given by

$$\begin{aligned} \delta_\Lambda \omega_a^{(1)} &= \partial_a \Lambda^{(1)} - \frac{1}{2}\theta^{bc}(\partial_b \Lambda \partial_c \omega_a - \partial_b \omega_a \partial_c \Lambda) \\ &+ i[\Lambda, \omega_a^{(1)}] + i[\Lambda^{(1)}, \omega_a]. \end{aligned} \quad (14)$$

A solution is

$$\omega_a^{(1)} = -\frac{1}{4}\theta^{bc}\{\omega_b, \partial_c \omega_a + F_{ca}\}. \quad (15)$$

One thus has $\hat{E}_a^\mu = e_a^\mu$ to all orders in θ and $\hat{\omega}_a = \omega_a + \omega_a^{(1)} + \mathcal{O}(\theta^2)$.

The Seiberg-Witten map for the field strength is given by $\hat{F}_{ab} = F_{ab} + F_{ab}^{(1)} + \mathcal{O}(\theta^2)$, where

$$F_{ab}^{(1)} = \frac{1}{2}\theta^{cd}\{F_{ac}, F_{bd}\} - \frac{1}{4}\theta^{cd}\{\omega_c, (\partial_d + D_d)F_{ab}\}, \quad (16)$$

where $D_a = A_a = ie_a^\mu p_\mu + \frac{i}{2}\omega_a^{bc}\Sigma_{bc}$ is the commutative covariant derivative.

The commutative field strength F_{ab} contains the Riemann tensor $R_{ab}{}^{cd}$ as well as a torsion $T_{ab}{}^c$:

$$F_{ab} = i[D_a, D_b] = \frac{1}{2}R_{ab}{}^{cd}\Sigma_{cd} + T_{ab}{}^c D_c \quad (17)$$

with $R_{ab} = \frac{1}{2}R_{ab}{}^{cd}\Sigma_{cd}$ and $T_{ab}{}^c = (D_a e_b^\nu - D_b e_a^\nu)e_\nu^c$. The commutative covariant derivative D_a is torsion free ($T_{ab}{}^c = 0$) and compatible with a metric: $e_\mu^a D_a e_\nu^b = 0$. We now have all the required tools to consider actions that are invariant under general coordinate transformations.

III. ACTION FOR NONCOMMUTATIVE GENERAL RELATIVITY

The Seiberg-Witten map for the Riemann tensor R_{ab} which is the field strength tensor corresponding to a local noncommutative Lorentz transformation can be read from Eqs. (16) and (17) where the classical torsion is being set to

zero:

$$\hat{R}_{ab} = R_{ab} + R_{ab}^{(1)} + \mathcal{O}(\theta^2), \quad (18)$$

with

$$R_{ab}^{(1)} = \frac{1}{2}\theta^{cd}\{R_{ac}, R_{bd}\} - \frac{1}{4}\theta^{cd}\{\omega_c, (\partial_d + D_d)R_{ab}\}. \quad (19)$$

The noncommutative Riemann tensor is then given by

$$\hat{R}_{ab}(\hat{x}) = \frac{1}{2}\hat{R}_{ab}{}^{cd}(\hat{x})\Sigma_{cd}, \quad (20)$$

from which we can determine the corresponding noncommutative Ricci tensor, $\hat{R}_{ab}{}^{bd}$, and a Ricci scalar $\hat{R} = \hat{R}_{ab}{}^{ab}$ in terms of the classical fields using the above Seiberg-Witten map.

The noncommutative action is then given by

$$\begin{aligned} S &= \int d^4x \frac{1}{2\kappa^2} \hat{R}(\hat{x}) \\ &= \int d^4x \frac{1}{2\kappa^2} (R(x) + R^{(1)}(x)) + \mathcal{O}(\theta^2). \end{aligned} \quad (21)$$

In the second line we have made use of the Weyl quantization procedure which allows to replace the noncommutative variables by commuting ones by expanding the noncommutative fields using the Seiberg-Witten maps. The only correction to leading order in θ comes from the Seiberg-Witten map for the $\text{so}(3,1)$ gauge field. It is easy to verify that this action is Hermitian and invariant under unimodular coordinate transformations and local Lorentz transformations. This noncommutative general relativity theory is, by construction, torsion free. In the leading order in the expansion in θ we can use the classical relations:

$$\omega_\mu^{ab}(x) = \frac{1}{2}e_\mu^c(x)(\Omega^{ab}{}_c(x) - \Omega^b{}_c{}^a(x) - \Omega_c{}^{ab}(x)) \quad (22)$$

with

$$\Omega^{ab}{}_c(x) = e_\mu^a(x)e_\nu^b(x)(\partial^\mu e_c^\nu(x) - \partial^\nu e_c^\mu(x)). \quad (23)$$

Equation (21) represents an action for the noncommutative version of the unimodular theory of gravitation. The unimodular theory is known [13] to be classically equivalent to Einstein's General Relativity with a cosmological constant. Indeed, we can rewrite the action (21) in the form of an Einstein-Hilbert action by introducing a Lagrange multiplier Λ which appears to be an arbitrary integration constant:

$$\begin{aligned} S &= \int d^4x \left(\frac{1}{2\kappa^2} \det(e_\mu^a(x))(R(x) + R^{(1)}(x)) \right. \\ &\quad \left. + \Lambda(\det(e_\mu^a(x)) - 1) + \mathcal{O}(\theta^2) \right). \end{aligned} \quad (24)$$

In deriving (24) we have used

$$\det \star(e_\mu^a(x)) \stackrel{\text{def}}{=} \frac{1}{4!} \epsilon^{\mu\nu\rho\sigma} \epsilon_{abcd} e_\mu^a(x) \star e_\nu^b(x) \star e_\rho^c(x) \star e_\sigma^d(x) = \det(e_\mu^a(x)) + \mathcal{O}(\theta^2), \quad (25)$$

where \star is the star product.

Let us now consider the weak field approximation of the noncommutative action (24). Although the theory defined by the action (24) does not admit flat spacetime as a background solution, we can still locally (for regions with volume $V \ll 1/\Lambda$) expand the tetrad around flat spacetime:

$$e_a^\mu(x) = \eta_a^\mu - \frac{1}{2} h_a^\mu(x). \quad (26)$$

Here $h_a^\mu(x)$ is a weak gravitational field subject to the traceless condition [$\det(e_\mu^a(x)) = 1$ follows from (24)]. The noncommutative correction to Einstein's action in the weak field limit reads

$$\frac{1}{8} \theta^{ln} \tilde{R}_{kl}{}^{ab} \tilde{R}_{mn}{}^{cd} d_{abcd}^{km} - \frac{1}{16} \theta^{ln} \tilde{R}_{ln}{}^{ab} \tilde{R}_{km}{}^{cd} d_{abcd}^{km} \quad (27)$$

where d_{abcd}^{km} are the structure constants defined by $d_{abcd}^{km} \Sigma_{km} = 2\{\Sigma_{ab}, \Sigma_{cd}\}$. Notice that

$$\tilde{R}_{km}{}^{ab} = -\frac{1}{2} \partial_k \left(\partial^a h_m^b + \frac{1}{2} \partial^b h_m^a \right) - \partial_m (\partial^a h_k^b - \partial^b h_k^a) \quad (28)$$

is the leading order of the weak field approximation of $R_{ab}{}^{cd}$. This modification of the linearized noncommutative action implies a noncommutative correction of the equations of motion for the weak gravitational field:

$$R^a{}_b - \frac{1}{2} \eta^a{}_b R = \frac{1}{8} \theta^{an} (\partial^r \partial_k \tilde{R}_{mn}{}^{cd} + \partial^r \partial_n \tilde{R}_{km}{}^{cd}) d_{rbcd}^{km} - \frac{1}{8} \theta^{ln} \partial^r \partial_l \tilde{R}_{mn}{}^{cd} d_{rbcd}^{am} \quad (29)$$

where $R^a{}_b$ is the usual Ricci tensor and R is the corresponding Ricci scalar in the linearized approximation (we have omitted the contribution coming from the cosmological constant). This modification might have some interesting physical implications that will be studied elsewhere.

Finally we briefly discuss the relation between the tetrad formalism considered here and a second-order formalism which involves the metric tensor. The noncommutative metric tensor defined naively as $\hat{g}_{\mu\nu}(\hat{x}) = E_\mu^a(\hat{x}) E_\nu^b(\hat{x}) \eta_{ab}$

is neither real nor symmetric. This raises a more generic question of the geometrical interpretation of the noncommutative deformation of Einstein's General Relativity considered above. The simple prescription to define the metric in our case is to solve the deformed equations of motion for the classical tetrads at each given order of θ expansion and then to determine the metric in the standard way.

IV. CONCLUSIONS

We have constructed a theory of noncommutative General Relativity on a canonical noncommutative spacetime. General coordinate transformations are shown to be restricted to the volume-preserving transformations. Thus the General Relativity on canonical noncommutative spacetimes is the noncommutative version of the unimodular theory of gravitation. The local Lorentz invariance is described as a noncommutative gauge theory by taking the spin connection field in the enveloping algebra.

The action for noncommutative General Relativity was constructed and the expansion in first order of the noncommutative parameter θ has been calculated. We derived the noncommutative correction to the equations of motion of the weak gravitational field. An interesting question is whether the effects of spacetime noncommutativity coming from the noncommutative modifications of gravity are stronger than the ones coming from the modifications of the interactions of the standard model [14].

It will also be interesting to consider classical solutions of the noncommutative action defined in this work. The minimal length introduced in our version of General Relativity could have interesting consequences for the horizon and the singularity of black holes. It will also be worth studying cosmological implications and the quantization of the noncommutative action.

ACKNOWLEDGMENTS

The work of X. C. was supported in part by the IISN and of the Belgian science policy office (IAP V/27). The work of A. K. was partially supported by the GRDF Grant No. 3305.

- [1] A. Einstein, Ann. Phys. (Leipzig) **49**, 769 (1916).
- [2] X. Calmet, M. Graesser, and S. D. H. Hsu, Phys. Rev. Lett. **93**, 211101 (2004).
- [3] A. H. Chamseddine, Phys. Lett. B **504**, 33 (2001).

- [4] P. Aschieri, C. Blohmann, M. Dimitrijevic, F. Meyer, P. Schupp, and J. Wess, hep-th/0504183.
- [5] J. Wess, in Lectures given at the BW2003 Workshop on Mathematical, Theoretical and Phenomenological

- Challenges Beyond the Standard Model: Perspectives of Balkans Collaboration, Vrnjacka Banja, Serbia, 2003 (hep-th/0408080).
- [6] M. Chaichian, P. P. Kulish, K. Nishijima, and A. Tureanu, Phys. Lett. B **604**, 98 (2004).
- [7] X. Calmet, Phys. Rev. D **71**, 085012 (2005).
- [8] B. Jurco, S. Schraml, P. Schupp, and J. Wess, Eur. Phys. J. C **17**, 521 (2000).
- [9] J. Madore, S. Schraml, P. Schupp, and J. Wess, Eur. Phys. J. C **16**, 161 (2000).
- [10] N. Seiberg and E. Witten, J. High Energy Phys. **09** (1999) 032.
- [11] R. Jackiw, S. Y. Pi, and A. P. Polychronakos, Ann. Phys. (N.Y.) **301**, 157 (2002).
- [12] J. Lukierski, P. C. Stichel, and W. J. Zakrzewski, Ann. Phys. (N.Y.) **306**, 78 (2003).
- [13] The equations of motion corresponding to this theory had first been written down by Albert Einstein in Sitz. Preuss. Akad. Wiss. Berlin (Math. Phys.) (1919); A. Einstein *et al.*, *Do Gravitational Fields Play an Essential Role in the Structure of Elementary Particles of Matter?* in *The Principle of Relativity: A Collection of Original Memoirs on the Special and General Theory of Relativity* (Dover, New York, 1923), pp. 191–198, English translation. The theory has been rediscovered in J. J. van der Bij, H. van Dam, and Y. J. Ng, Physica **116A**, 307 (1982), and further developed by a number of authors. See e.g., F. Wilczek, Phys. Rep. **104**, 143 (1984); W. Buchmuller and N. Dragon, Phys. Lett. B **207**, 292 (1988); M. Henneaux and C. Teitelboim, Phys. Lett. B **222**, 195 (1989); W. G. Unruh, Phys. Rev. D **40**, 1048 (1989).
- [14] X. Calmet, Eur. Phys. J. C **41**, 269 (2005).