

## Second order noncommutative corrections to gravity

Article (Published Version)

Calmet, Xavier and Kobakhidze, Archil (2006) Second order noncommutative corrections to gravity. *Physical Review D*, 74 (4). ISSN 1550-2368

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/41210/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

## Second order noncommutative corrections to gravity

Xavier Calmet<sup>1,\*</sup> and Archil Kobakhidze<sup>2,†</sup>

<sup>1</sup>*Université Libre de Bruxelles, Service de Physique Théorique, CP225 Boulevard du Triomphe (Campus plaine), B-1050 Brussels, Belgium*

<sup>2</sup>*Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA*  
(Received 30 May 2006; published 29 August 2006)

In this work, we calculate the leading order corrections to general relativity formulated on a canonical noncommutative spacetime. These corrections appear in the second order of the expansion in theta. First order corrections can only appear in the gravity-matter interactions. Some implications are briefly discussed.

DOI: [10.1103/PhysRevD.74.047702](https://doi.org/10.1103/PhysRevD.74.047702)

PACS numbers: 11.10.Nx, 04.20.-q

It is difficult to formulate general relativity on noncommutative spaces, and there are thus different approaches in the literature. In [1] for example a deformation of Einstein's gravity was studied using a construction based on gauging the noncommutative SO(4,1) de Sitter group and the Seiberg-Witten map with subsequent contraction to ISO(3,1). Most recently constructions of a noncommutative gravitational theory [2,3] were proposed based on a twisted Poincaré algebra [4,5]. The main problem in formulating a theory of gravity on noncommutative manifolds is that it is difficult to implement symmetries such as general coordinate covariance and local Lorentz invariance and to define derivatives which are torsion-free and satisfy the metricity condition.

Another approach has been proposed based on true physical symmetries [6,7] (see also [8]). In that approach one restricts the noncommutative action to symmetries of the noncommutative algebra

$$[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu}, \quad (1)$$

(see also [9] where this idea was applied to Lorentz symmetry). Obviously, the commutator (1) explicitly violates general coordinate covariance since  $\theta^{\mu\nu}$  is constant in all reference frames. However, we can identify a subclass of general coordinate transformations,

$$\hat{x}^{\mu'} = \hat{x}^\mu + \hat{\xi}^\mu(\hat{x}), \quad (2)$$

which are compatible with the algebra given by (1). The hat on the function  $\hat{\xi}(\hat{x})$  indicates that it is in the enveloping algebra. Under the change of coordinates (2) the commutator (1) transforms as

$$\begin{aligned} [\hat{x}^{\mu'}, \hat{x}^{\nu'}] &= \hat{x}^{\mu'} \hat{x}^{\nu'} - \hat{x}^{\nu'} \hat{x}^{\mu'} \\ &= i\theta^{\mu\nu} + [\hat{x}^\mu, \hat{\xi}^\nu] + [\hat{\xi}^\mu, \hat{x}^\nu] + \mathcal{O}(\hat{\xi}^2). \end{aligned} \quad (3)$$

Requiring that  $\theta^{\mu\nu}$  remains constant yields the following partial differential equations

$$\theta^{\mu\alpha} \hat{\partial}_\alpha \hat{\xi}^\nu(\hat{x}) = \theta^{\nu\beta} \hat{\partial}_\beta \hat{\xi}^\mu(\hat{x}). \quad (4)$$

A nontrivial solution to this condition can be easily found

$$\hat{\xi}^\mu(\hat{x}) = \theta^{\mu\nu} \hat{\partial}_\nu \hat{f}(\hat{x}), \quad (5)$$

where  $\hat{f}(\hat{x})$  is an arbitrary field. This noncommutative general coordinate transformation corresponds to the following transformation:  $\hat{\xi}^\mu(x) = \theta^{\mu\nu} \partial_\nu \hat{f}(x)$ . The Jacobian of this restricted coordinate transformations is equal to 1, meaning that the volume element is invariant:  $d^4x' = d^4x$ . The version of general relativity based on volume-preserving diffeomorphism is known as the unimodular theory of gravitation [10]. Thus we came to the conclusion that symmetries of canonical noncommutative spacetime naturally lead to the noncommutative version of unimodular gravity. We obtain the noncommutative field-strength of the SO(3,1) gauge symmetry

$$\hat{R}_{ab} = R_{ab} + R_{ab}^{(1)} + R_{ab}^{(2)} + \mathcal{O}(\theta^3), \quad (6)$$

with

$$R_{ab}^{(1)} = \frac{1}{2}\theta^{cd}\{R_{ac}, R_{bd}\} - \frac{1}{4}\theta^{cd}\{\omega_c, (\partial_d + D_d)R_{ab}\}. \quad (7)$$

The noncommutative Riemann tensor is then given by

$$\hat{R}_{ab}(\hat{x}) = \frac{1}{2}\hat{R}_{ab}{}^{cd}(\hat{x})\hat{\Sigma}_{cd}, \quad (8)$$

and the leading order correction in  $\theta^{ab}$  is found explicitly to be

$$\begin{aligned} \hat{R}_{ab}^{(1)pq}(x) &= \frac{1}{2}\theta^{cd}R_{ac}{}^{ij}R_{bd}{}^{kl}d_{ijkl}^{pq} \\ &\quad - \frac{1}{4}\theta^{cd}\omega_c^{ij}(\partial_d + D_d)R_{ab}{}^{kl}d_{ijkl}^{(1)pq} \end{aligned} \quad (9)$$

the coefficient  $d_{ijkl}^{(1)pq}$  is defined by

$$d_{ijkl}^{(1)pq} = \text{Tr}(\{\Sigma_{ij}, \Sigma_{kl}\}\Sigma^{pq}), \quad (10)$$

where the trace goes over the matrix indices of the SO(3,1) generators  $\Sigma_{ij}$ . The group-theoretic coefficients of Eq. (10) are all vanishing by virtue of antisymmetry property of the

\*Electronic address: [xcalmet@ulb.ac.be](mailto:xcalmet@ulb.ac.be)

†Electronic address: [kobakhid@physics.unc.edu](mailto:kobakhid@physics.unc.edu)

SO(3,1) generators,  $\Sigma_{ij}^T = -\Sigma_{ij}$  and cyclic properties of the trace [11]. This can be explicitly demonstrated for an arbitrary representation for the generators, e.g.  $\Sigma_{ab} = \frac{i}{4} \times [\gamma_a, \gamma_b]$

The new result of this work is the second order correction in  $\theta^{ab}$  which is given by

$$\begin{aligned}
R_{rs}^{(2)mn} = & \frac{1}{32} \theta^{ij} \theta^{kl} (2\omega_i^{ab} \partial_j \omega_k^{cd} \partial_l R_{rs}^{ef} d_{abcdef}^{(2)mn} \\
& - 2R_{ik}^{ab} \omega_j^{cd} \partial_l R_{rs}^{ef} d_{abcdef}^{(2)mn} \\
& - 2\partial_l \partial_i R_{rs}^{ab} \omega_j^{cd} \omega_k^{ef} d_{abcdef}^{(2)mn} \\
& + 2i\omega_k^{ab} \partial_l \omega_i^{cd} R_{rs}^{ef} \omega_j^{gh} d_{abcdefgh}^{(3)mn} \\
& - 4\omega_k^{ef} \partial_l (R_{ri}^{ab} R_{sj}^{cd}) d_{abcdef}^{(2)mn} \\
& - 4i\partial_i \omega_k^{ab} \partial_j \partial_l R_{rs}^{cd} d_{abcd}^{(4)mn} \\
& + i\omega_i^{ab} \partial_j \omega_k^{cd} R_{rs}^{ef} \omega_l^{gh} d_{abcdefgh}^{(5)mn} \\
& - iR_{ik}^{ab} \omega_j^{cd} R_{rs}^{ef} \omega_l^{gh} d_{abcdefgh}^{(5)mn} \\
& - 2i\omega_k^{ab} R_{ri}^{cd} R_{sj}^{ef} \omega_l^{gh} d_{abcdefgh}^{(5)mn} \\
& + i\omega_k^{ab} R_{rs}^{cd} \omega_i^{ef} \partial_j \omega_l^{gh} d_{abcdefgh}^{(6)mn} \\
& - i\omega_k^{ab} R_{rs}^{cd} R_{il}^{ef} \omega_j^{gh} d_{abcdefgh}^{(6)mn} \\
& + 2\partial_i \omega_k^{ab} \partial_j R_{rs}^{cd} \omega_l^{ef} d_{abcdef}^{(2)mn} \\
& + 2\partial_i (\omega_k^{ab} R_{rs}^{cd}) \partial_j \omega_l^{ef} d_{abcdef}^{(7)mn} \\
& + 2\partial_i R_{rk}^{ab} \omega_j^{cd} R_{sl}^{ef} d_{abcdef}^{(2)mn} \\
& - 2i\omega_i^{ab} R_{rk}^{cd} \omega_j^{ef} R_{sl}^{gh} d_{ghabcdef}^{(3)mn} \\
& - 2R_{rk}^{ab} R_i^{cd} \partial_j R_{sl}^{ef} d_{efabcd}^{(2)mn} \\
& + R_{rk}^{ab} R_{is}^{cd} R_{jl}^{ef} d_{efabcd}^{(2)mn} \\
& + 4i\partial_i R_{rk}^{ab} \partial_j R_{sl}^{cd} d_{abcd}^{(4)mn} \\
& + 2\omega_k^{ab} \partial_l (\omega_i^{cd} \partial_j R_{rs}^{ef} d_{abcdef}^{(2)mn} \\
& + i\omega_k^{ab} \omega_i^{cd} \partial_j R_{rs}^{ef} \omega_l^{gh} d_{abcdefgh}^{(5)mn} \\
& - i\omega_k^{ab} \partial_i R_{rs}^{cd} \omega_j^{ef} \omega_l^{gh} d_{abcdefgh}^{(5)mn} \\
& - \omega_k^{ab} \omega_i^{cd} R_{rs}^{ef} \omega_j^{gh} \omega_l^{pq} d_{abcdefghpq}^{(8)mn}) \quad (11)
\end{aligned}$$

using the result obtained for a generic noncommutative gauge theory in [12] and where the coefficients  $d^{(i)}$  are defined by:

$$d_{abcdef}^{(2)mn} = \text{Tr}(\{\{\Sigma_{ab}, \Sigma_{cd}\}, \Sigma_{ef}\} \Sigma^{mn}), \quad (12)$$

$$d_{abcdefgh}^{(3)mn} = \text{Tr}(\{\Sigma_{ab}, \{\{\Sigma_{cd}, \Sigma_{ef}\}, \Sigma_{gh}\}\} \Sigma^{mn}), \quad (13)$$

$$d_{abcd}^{(4)mn} = \text{Tr}([\Sigma_{ab}, \Sigma_{cd}] \Sigma^{mn}), \quad (14)$$

$$d_{abcdefgh}^{(5)mn} = \text{Tr}(\{\{\{\Sigma_{ab}, \Sigma_{cd}\}, \Sigma_{ef}\}, \Sigma_{gh}\} \Sigma^{mn}), \quad (15)$$

$$d_{abcdefgh}^{(6)mn} = \text{Tr}(\{\{\Sigma_{ab}, \Sigma_{cd}\}, \{\Sigma_{ef}, \Sigma_{gh}\}\} \Sigma^{mn}), \quad (16)$$

$$d_{abcdef}^{(7)mn} = \text{Tr}([\Sigma_{ab}, \Sigma_{cd}], \Sigma_{ef}) \Sigma^{mn}, \quad (17)$$

$$d_{abcdefghpq}^{(8)mn} = \text{Tr}(\{\{\Sigma_{ab}, \{\{\Sigma_{cd}, \Sigma_{ef}\}, \Sigma_{gh}\}\}, \Sigma_{pq}\} \Sigma^{mn}). \quad (18)$$

These coefficients are easily calculable using a specific representation, e.g. spinorial representation, for the matrices  $\Sigma_{ab}$  and a computer algebra program such as Mathematica with the routine TRACER [13]. The noncommutative action is then given by

$$\begin{aligned}
S &= \int d^4x \frac{1}{2\kappa^2} \hat{R}(\hat{x}) \\
&= \int d^4x \frac{1}{2\kappa^2} (R(x) + R^{(2)}(x)) + \mathcal{O}(\theta^3). \quad (19)
\end{aligned}$$

This equation is an action for the noncommutative version of the unimodular theory of gravitation. The unimodular theory is known [10] to be classically equivalent to Einstein's general relativity with a cosmological constant and it can be put in the form

$$S_{NC} = \frac{-1}{16\pi G} \int d^4x \sqrt{-g} R(g^{\mu\nu}) + \mathcal{O}(\theta), \quad (20)$$

where  $R(g^{\mu\nu})$  is the usual Ricci scalar and  $g$  is the determinant of the metric. If we restrict ourselves to the transformations (5), the determinant of the metric is always equal to minus one, the term  $\sqrt{-g}$  in the action is thus trivial. However, as mentioned previously, we recover full general coordinate invariance in the limit  $\theta^{\mu\nu}$  to zero and it is thus important to write this term explicitly to study the symmetries of the action. In order to obtain the equations of motion corresponding to this action, we need to consider variations of (20) that preserve  $g = \det g^{\mu\nu} = -1$ , i.e. not all the components of  $g_{\mu\nu}$  are independent. One thus introduces a new variable  $\tilde{g}_{\mu\nu} = g^{1/4} g_{\mu\nu}$ , which has explicitly a determinant equal to one. The field equations are then

$$R^{\mu\nu} - \frac{1}{4} g^{\mu\nu} R + \mathcal{O}(\theta) = 0. \quad (21)$$

As done in e.g. [10] we can use the Bianchi identities for  $R$  and find:

$$R_{;\mu} = 0, \quad (22)$$

which can be integrated easily and gives  $R = \Lambda$ , where  $\Lambda$  is an integration constant. It can then be shown that the differential equations (21) imply

$$R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R - \Lambda g^{\mu\nu} + \mathcal{O}(\theta) = 0, \quad (23)$$

i.e. Einstein's equations of general relativity with a cos-

mological constant  $\Lambda$  that appears as an integration constant. Because any solution of Einstein's equations with a cosmological constant can, at least over any topologically  $R^4$  open subset of spacetime, be written in a coordinate system with  $g = -1$ , the physical content of unimodular gravity is identical at the classical level to that of Einstein's gravity with some cosmological constant [10].

The form of the  $\mathcal{O}(\theta^2)$  corrections in Eq. (11) suggests that in the linearized approximation, gravity is not affected by spacetime noncommutativity. Note also that in the full

gravity-matter action the dominant  $\mathcal{O}(\theta)$  will generally be present in the matter Lagrangian, that in turn could affect the solutions for the metric in this order. It would be very interesting to study cosmological perturbations in the above setting.

The work of X. C. was supported in part by the IISN and the Belgian science policy office (IAP V/27). We are grateful to C. P. Martin for a useful discussion.

- 
- [1] A. H. Chamseddine, Phys. Lett. B **504**, 33 (2001).
  - [2] P. Aschieri, C. Blohmann, M. Dimitrijevic, F. Meyer, P. Schupp, and J. Wess, Classical Quantum Gravity **22**, 3511 (2005); P. Aschieri, M. Dimitrijevic, F. Meyer, and J. Wess, Classical Quantum Gravity **23**, 1883 (2006).
  - [3] A. Kobakhidze, hep-th/0603132.
  - [4] M. Chaichian, P. P. Kulish, K. Nishijima, and A. Tureanu, Phys. Lett. B **604**, 98 (2004).
  - [5] J. Wess, report “*Workshop on Mathematical, Theoretical and Phenomenological Challenges Beyond the Standard Model: Perspectives of Balkans Collaboration, Vrnjacka Banja, Serbia, 2003*,” (unpublished).
  - [6] X. Calmet and A. Kobakhidze, Phys. Rev. D **72**, 045010 (2005).
  - [7] X. Calmet, hep-th/0510165.
  - [8] E. Harikumar and V. O. Rivelles, hep-th/0607115.
  - [9] X. Calmet, Phys. Rev. D **71**, 085012 (2005); Proceedings of 41st Rencontres de Moriond on Electroweak Interactions and Unified Theories, La Thuile, Aosta Valley, Italy, 2006.
  - [10] The equations of motion corresponding to this theory have first been written down by Albert Einstein in: A. Einstein (Siz. Preuss. Acad. Scis., 1919); A. Einstein *et al.* *Do Gravitational Fields Play an Essential Role in the Structure of Elementary Particle of Matter*, in The Principle of Relativity: A Collection of Original Memoirs on the Special and General Theory of Relativity (Dover, New York, 1923), p. 191, English translation. The theory has been rediscovered in J. J. van der Bij, H. van Dam, and Y. J. Ng, Physica A (Amsterdam) **116**, 307 (1982), and further developed by a number of authors, see e.g., F. Wilczek, Phys. Rep. **104**, 143 (1984); W. Buchmuller and N. Dragon, Phys. Lett. B **207**, 292 (1988); M. Henneaux and C. Teitelboim, Phys. Lett. B **222**, 195 (1989); W. G. Unruh, Phys. Rev. D **40**, 1048 (1989).
  - [11] P. Mukherjee (private communication).
  - [12] L. Moller, J. High Energy Phys. **10** (2004) 063.
  - [13] M. Jamin and M. E. Lautenbacher, Comput. Phys. Commun. **74**, 265 (1993).