

The role of mutation rate variation and genetic diversity in the architecture of human disease

Article (Published Version)

Chen Eyre-Walker, Ying and Eyre-Walker, Adam (2014) The role of mutation rate variation and genetic diversity in the architecture of human disease. PLoS ONE, 9. e90166. ISSN 1932-6203

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/47655/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

The Role of Mutation Rate Variation and Genetic Diversity in the Architecture of Human Disease

Ying Chen Eyre-Walker, Adam Eyre-Walker*

School of Life Sciences, University of Sussex, Brighton, United Kingdom

Abstract

Background: We have investigated the role that the mutation rate and the structure of genetic variation at a locus play in determining whether a gene is involved in disease. We predict that the mutation rate and its genetic diversity should be higher in genes associated with disease, unless all genes that could cause disease have already been identified.

Results: Consistent with our predictions we find that genes associated with Mendelian and complex disease are substantially longer than non-disease genes. However, we find that both Mendelian and complex disease genes are found in regions of the genome with relatively low mutation rates, as inferred from intron divergence between humans and chimpanzees, and they are predicted to have similar rates of non-synonymous mutation as other genes. Finally, we find that disease genes are in regions of significantly elevated genetic diversity, even when variation in the rate of mutation is controlled for. The effect is small nevertheless.

Conclusions: Our results suggest that gene length contributes to whether a gene is associated with disease. However, the mutation rate and the genetic architecture of the locus appear to play only a minor role in determining whether a gene is associated with disease.

Citation: Eyre-Walker YC, Eyre-Walker A (2014) The Role of Mutation Rate Variation and Genetic Diversity in the Architecture of Human Disease. PLoS ONE 9(2): e90166. doi:10.1371/journal.pone.0090166

Editor: I. King Jordan, Georgia Institute of Technology, United States of America

Received: August 23, 2013; **Accepted:** January 28, 2014; **Published:** February 27, 2014

Copyright: © 2014 Eyre-Walker, Eyre-Walker. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: These authors have no support or funding to report.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: a.c.eyre-walker@sussex.ac.uk

Introduction

Why do humans suffer from the diseases that we do? In part this is clearly due to our anatomy and physiology, and that of the organisms that infect us - we cannot have a disease of an organ that we do not possess. But why do we suffer from cystic fibrosis rather than some other disease of the lungs? One simple reason might be variation in the mutation rate. Those genes and genomic regions that have high mutation rates are more likely to generate disease mutations, and hence be associated with a disease. The rate of mutation of a locus will depend upon two factors: the rate of mutation per site and the number of sites at which a mutation can generate a disease phenotype. The per site mutation rate is known to vary across the human genome at a number of different scales such that some genes have mutation rates that are several fold higher than other genes (reviewed in Hodgkinson *et al.* [1]). Genes also vary considerably in their length, with some of the largest, such as the dystrophin gene, being associated with disease.

A more subtle factor affecting the likelihood of a gene being associated with a disease is the genealogy. At each site in the genome there is an underlying genealogy whereby every chromosome in the population is related via a bifurcating tree to every other chromosome at that site. If there is no recombination between sites then sites share the same genealogy. The shape and depth of the genealogy depends on several factors. The first is chance; for example, the average total length of a genealogy for a neutral locus in a population of stationary size is expected to be

proportional to $4N$ generations in a diploid species, where N is the population size, but this is expected to have a variance of at least $(4N)^2$ generations [2]. Second, the genealogy depends on the effective population size of the locus (N_e). N_e is thought to vary across the human genome as a consequence of natural selection [3,4]. Selection can reduce the N_e of a genomic region through either a selective sweep caused by the passage of an advantageous mutation through the population [5], or via background selection caused by the removal of deleterious mutations [6]. Those regions of the genome with low rates of recombination or a high density of selected sites are expected to have low N_e , and this is expected to reduce the genetic diversity of neutral and weakly selected variants in these regions (reviewed in [7]). Analyses suggest that N_e varies across the human genome by a few-fold [4]. The effective population size is not expected to affect the frequency of deleterious mutations in which the product of N_e and the strength of selection is greater than one. However, stochastic factors affecting the genealogy are expected to be important irrespective of the selection acting upon a mutation.

Previous analyses have shown that Mendelian disease genes are 30% longer than non-disease genes [8,9]. Comparative analyses have also shown that genes associated with Mendelian diseases have significantly, but only slightly higher rates of mutation per site, as inferred from levels of synonymous divergence between species [8,10]. The rather modest differences between disease and non-disease genes in the inferred mutation rate might be due to time frame over which the mutation rate was inferred: Smith and




Figure 1. CDS length. (A) Mean total CDS length, and (B) Mean average CDS length. Total CDS length is the sum of all constitutive and alternately spliced exons; average CDS length is the average CDS length of each transcript. Error bars represent the 95% confidence intervals. doi:10.1371/journal.pone.0090166.g001

Eyre-Walker [8] considered the divergence between human and mouse, and Huang et al. [10] considered the divergence between mouse and rat. This will give a poor estimate of the current mutation rate at a locus in humans because the relative mutation rate of a locus appears to have evolved through time [1,11]. The mutation rate has also recently been predicted, based on a model fitted to the locations of *de novo* mutations in humans, to be slightly higher in disease associated genes [12], but the accuracy of this model is unproven, and they consider the total mutation rate of the exon, rather than the rate at non-synonymous sites. Here we

consider the divergence between humans and their most closely related extant relative, chimpanzee, as our measure of the mutation rate. We also consider whether the density of single nucleotide polymorphism (SNP) is greater in disease than non-disease genes.

Materials and Methods

To estimate mutation rate for each gene, we estimated their intron divergence between the human and chimpanzee genomes as follows. Alignments using the NCBI build 36 version of the




Figure 2. Mutation rates. The mutation rate per site, as inferred from intron divergence between human and chimpanzee. A) Intron divergence per site between human and chimpanzee; B) the predicted non-synonymous mutation rate per CDS site. Error bars represent the 95% confidence intervals.

doi:10.1371/journal.pone.0090166.g002

human genome (hg18) and PanTro2 version of the chimp genome were downloaded from the UCSC website (<http://genome.ucsc.edu/>). Alignments were parsed into individual genic sequences and realigned with MAFFT version 6 (<http://mafft.cbrc.jp/alignment/software/>). Exon sequences were masked according

to exon annotation of the NCBI build 36 version of the human genome from the ensemble database (<http://www.ensembl.org/>). We did not correct for multiple hits; this is not necessary since the average intron divergence between human and chimpanzee sequences is 1.05% [13]. We calculated the rates of intron

divergence for CpG and nonCpG sites separately since the former have much higher rates of mutation. We used these intron divergences to infer the rate of non-synonymous mutation in human exons, by calculating the number of CpG and non-CpG sites in each exon which when mutated would give a non-synonymous change; in this calculation we assumed that all mutations at CpGs are transitions, which is a good approximation [14], and that 60% of mutations at other sites were transitions. If a gene had multiple transcripts we made these calculations for each transcript and averaged the result.

DNA sequence diversity data were taken from the 1000 genome project [15].

Genes were designated as being associated with Mendelian disease based upon the compilation made by Blekman *et al.* [16]. Genes associated with genome-wide association studies (GWAS) were obtained from the GWAS catalog (<http://www.genome.gov/gwastudies/>); a gene in which the strongest GWAS signal was found within the boundaries of a gene were designated as being a GWAS gene.

To investigate what factors might influence patterns of genic mutations, estimated by intron divergence, we considered a number of variables. Intron GC content, nucleosome occupancy, replication timing and male and female recombination rates were downloaded from the UCSC website (<http://genome.ucsc.edu/>). We used A365 values to study the influence of nucleosome occupancy on the distribution of genic mutations rate across the genome. Recombination rates per MB were from Kong *et al.* [17]. Replication time data were from Chen *et al.* [18] and Hansen *et al.* [19]. Qualitatively similar results were obtained using each of four replication time datasets, so we only present the analysis using data from an embryonic stem cell line BG02 [19]. Germ-line expression data were from a study by McVicker and Green [20].

The dataset for this analysis is available as Table S1.

Results

We predict that unless all possible diseases with a genetic basis, and all the genes that can cause them, have already been discovered, then genes associated with diseases should have higher genic mutation rates than non-disease genes, where the genic mutation rate is determined by the product of gene length and the mutation rate per site. We also predict that disease genes should be in relatively diverse regions of the genome. To investigate these predictions we compiled data from 17577 nuclear genes with introns, of which 854 genes are known to cause a Mendelian

disease. We also analysed 1732 genes in which the strongest signal in a genomic region in a genome wide association study (GWAS) lay within the boundaries of the gene (i.e. all exons and introns between the start and stop codon). The presence of an association signal within the boundaries of the gene does not necessarily mean that the causative mutation is within the protein coding sequence or even within the boundaries of the gene, and many of these associations may be in regulatory sequences [21]. We subsequently excluded genes on the sex chromosomes since the Y-chromosome is known to have a higher mutation rate and the X-chromosome a lower mutation rate than the autosomes [22]. This yielded a dataset of 17062 autosomal genes including 820 associated with a Mendelian disease and 1726 with a GWAS signal. Details of the dataset are given in Table S1.

Gene Length

Consistent with the hypothesis that disease genes should have higher overall rates of mutation we find, as others have in the past for genes causing Mendelian disease [8,9], that genes associated with disease are significantly longer, in terms of their total coding sequence (CDS) length (i.e. the sum of all constitutive and alternatively spliced exons), than non-disease genes; Mendelian disease genes are ~28% and GWAS genes ~44% longer than non-disease genes (One-way ANOVA $p < 0.001$) (Figure 1a). A similar pattern is evident for average CDS length; both Mendelian and GWAS disease genes are 50% longer than non-disease genes (Figure 1b). The difference in average CDS length is greater than in previous studies [8,9], but this is likely to be due to the improvement in genome annotation; the average length of genes is slightly shorter than in previous analyses.

Strikingly, the difference in length is as great or greater for the GWAS than the Mendelian disease genes despite the fact that many of the GWAS signals are likely to be outside the protein coding sequence [21]. GWAS genes might have longer CDSs for three reasons. First, genes with longer CDSs have a greater chance of generating a disease mutation. Second, longer genes are more likely to have a non-causative marker SNP in the CDS that is associated with the disease. And finally since intron and total CDS lengths are correlated ($r = 0.36$, $p < 0.001$), genes with long CDSs have longer introns and hence an increase chance of having causative or non-causative SNPs in their introns. However, if we control for the correlation between intron and CDS length by regressing CDS length against intron length and taking the residuals, we find that GWAS genes have longer CDSs, than non-disease genes, even given their longer introns (t-test $p < 0.001$;

Table 1. Standardised regression coefficients from multiple regressions.

Factor	Intron Divergence	Predicted non-synonymous mutation rate	Intron SNP density	Average genealogy length
GC content	0.525***	0.325***	0.192***	-0.212***
Nucleosome occupancy	-0.396***	-0.167***	-0.412***	-0.035
Female recombination rate	-0.020*	0.018*	0.058***	0.042***
Male recombination rate	0.202***	0.143***	0.129***	-0.048***
Germ-line expression	-0.062***	-0.116***	-0.020*	0.032***
Replication time	-0.132***	-0.157***	-0.071***	0.038***
Distance to telomere	-0.158***	-0.097***	-0.117***	0.060***
Distance to centromere	-0.018*	-0.016	0.020*	0.014

Note that the replication time data is such that a negative slope indicates an increase in the variable through the cell cycle * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. doi:10.1371/journal.pone.0090166.t001




Figure 3. Diversity and genealogy estimates. The diversity in disease and non-disease genes measured as the A) average intron SNP density, B) the average intron SNP density divided by intron divergence, C) and the mean minor allele frequency (MAF). Error bars represent the 95% confidence intervals.
doi:10.1371/journal.pone.0090166.g003

similar results are obtained if we regress log CDS length against

log intron length ($p < 0.001$). This suggests that GWAS genes are not simply longer because they have longer introns; it therefore seems that either GWAS genes are more likely to be associated with disease because some causative mutations are within their exons, or because there is a greater number of marker SNPs in exons.

Mutation Rates

However, contrary to our expectations, we find that disease genes are found in regions of the genome with significantly lower per site mutation rates, as measured by intron divergence between human and chimpanzee. The difference is highly significant (one-way ANOVA $p < 0.001$), but the difference is small with disease genes having approximately 5% lower intron divergence than non-disease genes (Figure 2a). The pattern differs between CpG and non-CpG sites, with disease genes having lower divergence at CpG sites and either similar or higher divergence at non-CpG sites (results not shown). If we calculate the expected non-synonymous mutation rate in the CDS by multiplying the proportion of non-synonymous sites that are CpG and non-CpG in the CDS by the respective levels of intron divergence, we still find that both Mendelian and complex disease genes have slightly lower mutation rates per site than non-disease genes ($p = 0.004$) (Figure 2b). As expected, both Mendelian and complex disease genes have significantly higher overall predicted rates of non-synonymous mutation ($p < 0.001$), driven by the fact that disease genes have longer CDSs.

The fact that disease genes have lower predicted rates of non-synonymous mutation per site is inconsistent with our hypothesis, but this might be due to the fact that they have features which predispose them to lower mutation rates - for example they might be transcribed at lower levels and hence have lower rates of mutation [23]. Divergence at intronic and intergenic sites is known to be significantly correlated to a number of other variables including GC-content [3,18,24,25], recombination rate [3,24,26,27,28], replication time [18,29,30], distance to the telomere and centromere [3,13,18,24], gene density [3,24], nucleosome occupancy [11] and expression level [23]. We confirm previous results and show that intron divergence is positive correlated to GC content and male recombination rate within a multiple regression; and that intron divergence is negatively correlated to replication time (later genes have higher divergence), distance to the telomere, distance to the centromere, female recombination rate, nucleosome occupancy and germ-line expression (Table 1). Similar patterns are evident for the predicted non-synonymous mutation rate (Table 1). If we take the residuals from a multiple regression of intron divergence against all the genomic variables above we find that intron divergence and the predicted rate of non-synonymous mutation do not differ significantly between disease and non-disease genes.

Genetic Diversity

Although, disease genes are found in regions of the genome with relatively low rates of intron mutation we find that disease genes have a significantly greater density of polymorphisms segregating in their introns than non-disease genes; the difference is 11% and 17% for the Mendelian and GWAS genes respectively (Figure 3a). If we divide the density of SNPs by the divergence of introns to estimate a quantity that is proportional to the average length of the genealogies at the locus, we find that Mendelian and GWAS genes have significantly longer average genealogy lengths that are 9% and 12% greater than non-disease genes (ANOVA $p < 0.001$; t-test of Mendelian versus non-disease $p < 0.001$; t-test of GWAS versus non-disease $p < 0.001$). It is odd that the difference between disease

and non-disease genes is less pronounced for average genealogy length than diversity given that disease genes have lower intron divergence than non-disease genes. This is probably due to non-linearities associated with ratios.

Although, we find that disease genes have higher diversities and average genealogy lengths than non-disease genes, we find no evidence that the predicted non-synonymous population mutation rate in the CDS (calculated as the proportion of non-synonymous sites that are CpG multiplied by the SNP density at CpG sites in introns plus the proportion of non-synonymous sites that are non-CpG multiplied by the SNP density at non-CpG in introns) differs between disease and non-disease genes. However, the calculation of the predicted non-synonymous population mutation rate is subject to considerable error because we have relatively few intron CpG sites and SNP density is very low in humans.

It is possible that disease genes have higher diversities and average genealogy lengths because disease genes have features that predispose them to higher values, not because by having higher values they are more likely to be associated with disease. We find that intron SNP density is positively correlated to GC content, female and male rates of recombination and distance to the centromere and negatively correlated to the time of replication (late genes have higher diversity), nucleosome occupancy, germline expression and distance to the telomere (Table 1). If control for these factors by taking the residuals from the multiple regression we find that SNP density is still significantly greater in both Mendelian and GWAS genes, than in non-disease genes (ANOVA $p < 0.001$; individual t-tests $p < 0.001$). Likewise we find the average genealogy length is positively correlated to all variables except GC content, nucleosome occupancy and male recombination rate (Table 1), and that after controlling for these associations, disease genes still have significantly greater average genealogy lengths than non-disease genes (ANOVA $p = 0.019$; individual t-tests Mendelian versus non-disease $p = 0.21$, GWAS versus non-disease $p = 0.001$).

Although disease genes have a greater number of SNPs per bp than non-disease genes the distribution of the genetic variation varies in an inconsistent manner between categories of genes; the average minor allele frequency is $\sim 10\%$ greater in Mendelian, and $\sim 10\%$ lower in GWAS genes, than in non-disease genes (ANOVA $p < 0.01$) (Figure 3b).

Discussion

We have found that genes associated with disease are longer and reside in regions of the genome with greater intron diversities and average genealogy lengths than non-disease genes. This is consistent with a role for mutation and genetic variation in determining whether a gene becomes associated with disease.

References

- Hodgkinson A, Eyre-Walker A (2011) Variation in the mutation rate across mammalian genomes. *Nature Reviews Genetics* 12: 756–766.
- Charlesworth B, Charlesworth D (2010) *Elements of evolutionary genetics*. Greenwood Village: Ben Roberts.
- Hellmann I, Prüfer K, Ji H, Zody MC, Paabo S, et al. (2005) Why do human diversity levels vary at a megabase scale? *Genome Res* 15: 1222–1231.
- Gossmann TI, Woolfit M, Eyre-Walker A (2011) Quantifying the variation in the effective population size within a genome. *Genetics* 189: 1389–1402.
- Maynard Smith J, Haigh J (1974) The hitch-hiking effect of a favourable gene. *Genet Res* 23: 23–35.
- Charlesworth B, Morgan MT, Charlesworth D (1993) The effect of deleterious mutations on neutral molecular variation. *Genetics* 134: 1289–1303.
- Charlesworth B (2009) Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. *Nat Rev Genet* 10: 195–205.
- Smith NG, Eyre-Walker A (2003) Human disease genes: patterns and predictions. *Gene* 318: 169–175.
- Kondrashov FA, Ogurtsov AY, Kondrashov AS (2004) Bioinformatical assay of human gene morbidity. *Nucleic acids research* 32: 1731–1737.
- Huang H, Winter EE, Wang H, Weinstock KG, Xing H, et al. (2004) Evolutionary conservation and selection of human disease gene orthologs in the rat and mouse genomes. *Genome biology* 5: R47.
- Hodgkinson A, Chen Y, Eyre-Walker A (2012) The large scale distribution of somatic mutations in cancer genomes. *Human Mutation* 33: 136–143.
- Michaelson JJ, Shi Y, Gujral M, Zheng H, Malhotra D, et al. (2012) Whole-genome sequencing in autism identifies hot spots for de novo germline mutation. *Cell* 151: 1431–1442.
- Chimpanzee-Sequencing-and-Analysis-Consortium (2005) Initial sequence of the chimpanzee genome and comparison with the human genome. *Nature* 437: 69–87.
- Nachman MW, Crowell SL (2000) Estimate of the mutation rate per nucleotide in humans. *Genetics* 156: 297–304.
- _Genomes_Project_Consortium (2012) An integrated map of genetic variation from 1,092 human genomes. *Nature* 491: 56–65.

However, we find no evidence that the mutation rate per site is greater in disease than non-disease genes. Nevertheless, what is ultimately important is the mutation rate of the gene, and we find that the overall mutation rate of disease genes is greater than non-disease genes because disease genes are longer ($p < 0.001$). The effect of gene length may be more conspicuous than for the other variables, because there is substantially more variation in CDS length per gene (coefficient of variation (CV) = 0.78) than in intron divergence (CV = 0.56), intron SNP density (CV = 0.42) and average genealogy length (CV = 0.47); in reality the differences in CV are even larger because intron divergence, and in particular SNP density and average genealogy length, are likely to be subject to large sampling error variances that CDS length is not.

We have interpreted the fact that disease genes are longer than non-disease genes as evidence that genes with higher mutation rates are more likely to generate disease mutations, however, it is possible that disease genes are longer simply because genes involved in particular processes that could cause disease to be longer. It is difficult to test this hypothesis without knowing all the genes that might cause disease. We have also interpreted the greater diversity in disease genes as being what causes them to be associated with disease. However, in the case of the complex disease genes this might simply reflect a bias towards a better ability to detect GWAS signals in regions of higher diversity.

Although, we have found that disease genes are longer than non-disease genes, and that they have greater diversity and average genealogy lengths, the differences are fairly small. It is therefore evident that either most disease associated genes have been discovered, which seems unlikely, or that the function of the gene is far more important in determining whether a gene causes disease than its effective mutation rate.

Supporting Information

Table S1 The data matrix used in the analysis. A description of column headings is provided as a separate worksheet within the Excel spreadsheet. (XLSX)

Acknowledgments

The authors are grateful to the comments of three anonymous referees.

Author Contributions

Conceived and designed the experiments: YEW AEW. Performed the experiments: YEW AEW. Analyzed the data: YEW AEW. Contributed reagents/materials/analysis tools: YEW AEW. Wrote the paper: YEW AEW.

16. Blekhnman R, Man O, Herrmann L, Boyko AR, Indap A, et al. (2008) Natural selection on genes that underlie human disease susceptibility. *Curr Biol* 18: 883–889.
17. Kong A, Gudbjartsson DF, Sainz J, Jonsdottir GM, Gudjonsson SA, et al. (2002) A high-resolution recombination map of the human genome. *Nat Genet* 31: 241–247.
18. Chen CL, Rappailles A, Duquenne L, Huvet M, Guilbaud G, et al. (2010) Impact of replication timing on non-CpG and CpG substitution rates in mammalian genomes. *Genome Res* 20: 447–457.
19. Hansen RS, Thomas S, Sandstrom R, Canfield TK, Thurman RE, et al. (2010) Sequencing newly replicated DNA reveals widespread plasticity in human replication timing. *Proceedings of the National Academy of Sciences of the United States of America* 107: 139–144.
20. McVicker G, Green P (2010) Genomic signatures of germline gene expression. *Genome research* 20: 1503–1511.
21. Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, et al. (2012) Systematic localization of common disease-associated variation in regulatory DNA. *Science* 337: 1190–1195.
22. Ellegren H (2007) Characteristics, causes and evolutionary consequences of male-biased mutation. *Proc Biol Sci* 274: 1–10.
23. Park C, Qian W, Zhang J (2012) Genomic evidence for elevated mutation rates in highly expressed genes. *EMBO reports* 13: 1123–1129.
24. Tyekuceva S, Makova KD, Karro JE, Hardison RC, Miller W, et al. (2008) Human-macaque comparisons illuminate variation in neutral substitution rates. *Genome Biol* 9: R76.
25. Wolfe KH, Sharp PM, Li W-H (1989) Mutation rates differ among regions of the mammalian genome. *Nature* 337: 283–285.
26. Duret L, Arndt PF (2008) The impact of recombination on nucleotide substitutions in the human genome. *PLoS Genet* 4: e1000071.
27. Hellmann I, Ebersberger I, Ptak SE, Paabo S, Przeworski M (2003) A neutral explanation for the correlation of diversity with recombination rates in humans. *Am J Hum Genet* 72: 1527–1535.
28. Lercher MJ, Hurst LD (2002) Human SNP variability and mutation rate are higher in regions of high recombination. *Trends Genet* 18: 337–340.
29. Pink CJ, Hurst LD (2010) Timing of replication is a determinant of neutral substitution rates but does not explain slow Y chromosome evolution in rodents. *Mol Biol Evol* 27: 1077–1086.
30. Stamatoyannopoulos JA, Adzhubei I, Thurman RE, Kryukov GV, Mirkin SM, et al. (2009) Human mutation rate associated with DNA replication timing. *Nat Genet* 41: 393–395.