

Natural control of the mosquito population via *Odonata* and *Toxorhynchites*.

F. O. Faithpraise^{1*}, J. Idung¹, B. Usibe¹, C. R. Chatwin², R. Young², P. Birch²

Ph.D Student, Engineering & Design, (Biomedical Engineering) School of Engineering and Informatics, University of Sussex, Brighton –UK^{1*}.

Lecturer II, Zoology and Environmental Biology; University of Calabar, Nigeria¹.

Lecturer II, Electronic & Computer Technology Unit, Physics Department; University of Calabar, Nigeria¹.

Professor, Engineering & Design, (Biomedical Engineering) School of Engineering and Informatics, University of Sussex, Brighton –UK².

Reader, Engineering & Design, (Biomedical Engineering) School of Engineering and Informatics, University of Sussex, Brighton –UK².

Senior Lecturer, Engineering & Design, (Biomedical Engineering) School of Engineering and Informatics, University of Sussex, Brighton –UK².

Abstract: The main impact of mosquito pests is the transmission of many dangerous diseases and death. Hence, the reduction of their population by the use of a natural control method is a primary objective of this research. This mosquito reduction method utilises different species of predators (*Odonata*) and (*Toxorhynchites*) to substantially improve the environment. The frequency of capturing the pest mosquitoes by the predators is determined using a Pascal distribution, whilst insect mortality is modelled using a Weibull distribution. The results from the model show that by using insect predators, a significant reduction of the mosquito population is possible in less than eighty days.

Keywords: Natural control; Mosquito control; *Odonata*; *Toxorhynchites*; Disease Vector Control

I. INTRODUCTION

Several species of mosquitoes are a public health nuisance for the world and Africa as a continent as they drive extremely dangerous disease vectors. In order to manage public health, it is important to understand the lifecycle of the dangerous mosquito species and the agents that can control them.

Mosquito Lifecycle

All mosquito pest go through complete metamorphoses, from: egg to larva to pupa to adult, see figure 1. Their cycles are identical with slight differences and preferences relating to water clarity. The cycle begins when a female mosquito obtains a blood meal from either a human-being or other mammal to supply the required nutrients to produce as many as two hundred and fifty eggs at a time. She then seeks an aquatic location usually on the surface of stagnant water, or in a water filled depression, or on the edge of a container, where rainwater may have collected to lay eggs. After two days, the eggs hatch into larvae. The larvae live and feed on microorganisms in the water for seven to fourteen days and develop into pupae, which no longer feed. The mosquito then emerges from the pupa shell as a fully developed adult after four days. The moment the body of the mosquito finishes moulting, it is hyper-sensitive to carbon dioxide exhaled from mammals, it has poor eyesight and is very sensitive to mammal sweat scent

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

from a half mile distance [1], [2]; this enables it to locate a mammal target to seek out a blood meal by a bite on the animal or human skin. As the mosquito punctures the skin it injects its saliva into the flesh. If the mosquito is infected with any kind of disease, it is transferred into the prey through the saliva. There are about 2500 species of mosquitoes on the planet, of which 300 are well known disease carriers. Different species carry different diseases that are local to the area where they live; mosquitoes can attack and breed at different times of the day. Worldwide approximately seventy million people per annum catch diseases from mosquito bites [3]. Several attempts to manage and control the population of the harmful mosquito species have failed. Most of these dangerous species can be classified into: Permanent Water Mosquitoes, Floodwater Mosquitoes and Tree Hole/artificial container Mosquitoes. Permanent Flood Water Mosquitoes are found in swamps, ponds, lagoons, and ditches that permanently have water all year around, they include: *Aedes* illustrated by [4],[5], [6], [7]; *Culex pipiens*, [8]; *Culicinae*, [9]; *Culiseta*, [10] and *Anopheles gambiae* [11]. Floodwater Mosquitoes are found in areas that are flooded temporarily or seasonally and have the fastest breeding cycle of all mosquitoes up to 7 days. Tree hole/artificial container Mosquitoes are found and breed in: empty buckets, old tires, and anything that can hold water. Research has confirmed that these species of mosquitoes are epidemiologically a major vector for the spread of infectious disease, [12]; yellow fever virus, [13]; dengus fever, [14], [15]; Chikungunya fever, [16], [17]; Japanese encephalitis [18], [19]; Meningitis [20], [21]; Urticaria [22], [23]; West Nile virus [24], [25] and *Dirofilaria immitis* [26].

II. RELATED WORK

Most notable attempts made to control and manage the mosquito population and the deleterious effect of some classes of mosquitoes includes the design of an adhesive film trap to catch the egg depositing mosquitoes by Facchinelli, et al [27], and Gama, et al [28]. The comparison of the catching effect of gold standard traps with other traps was done by [29], and observed that for the trapping system to be successful users must comply with the deployment instructions, otherwise the trapping method is rendered ineffective with a corresponding increase in mosquito population density. [30], [31] implemented successful control of some species of mosquitoes using *Bacillus sphaericus* but *Bacillus sphaericus* usage could not continue due to its long term side effects on humans and some arthropods [32]. [33] evaluated the control of mosquitoes with *Romanomermis yunanensis* (Nematoda: Mermithidae). [34] demonstrated the substantial controls of the *Culex* mosquitoes larvae by the used of fish muddy loaches (*Misgurnus mizolepis*), but the adult mosquitoes were not controlled. [35] proposed the combination of some chemical component like *imidacloprid* and *permethrin*. [36] proposed the use of entomopathogenic fungus for the control of adult mosquito. [37] proposed the use of predator (Odonata) and its life cycle stages. Because of the long term harmful effect of most of the chemical pesticides and the increased resistance of mosquitoes [38], more favourable and better methods of mosquito control are still under research. [39],[40] suggested and demonstrated programmes based on environmental modification of pond and pool filling and several preventive measures based on mosquito breeding habitats as follows:

- Empty water from containers such as flower pots, birdbaths, pet water dishes, cans, gutters, tires and buckets regularly to disrupt the mosquito breeding cycle.
- Keep windows and door screens in good working order to prevent mosquitoes from entering domiciles.
- Put on long-sleeved cloths while outdoors, consider staying indoors early in the morning and evening when mosquitoes are most active.
- Use mosquito netting both in and outdoors.
- Consider using an insect repellent.
- Keep gutters clean and unclogged.
- Walk properties after rainfall and be sure downspouts are drained properly, without leaving puddles in the drainage area.
- Keep swimming pools cleaned and chlorinated, even when not in use to prevent mosquito breeding.
- Ornamental ponds should be aerated to keep water moving and discourage mosquitoes from laying eggs. Alternatively, stock the pond with mosquito-eating fish.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

This kind of programme is important but requires considerable time and financial investment, it does rely on people being very active, which is often difficult to sustain.

Temporary mosquito control methods

Some temporary mosquito control methods available on the market today include: Mosquito candles, [41], [42], [43]. Citronella candles are made from the essential oils of citronella grass with natural mosquito repellents. While the fragrance may deter mosquitoes, it will still not prevent the bite once the plume of the fragrance the candle releases expire. Conceal candles was made available as an improvement which delivers an inhibitor molecule found in the essential oils of certain plants to render the mosquitoes incapable of smelling and thereby incapable of finding someone to bite. However this product still has short comings, as it must be used in conjunction with more effective mosquito control methods to repel mosquitoes.

Mosquito repellent [44], Bug Zappers an electrical discharge insect control system [45], [46]. The bug zapper actually kills harmless and beneficial bugs. This makes this device an ineffective form of mosquito control [47]. Scientific studies prove that mosquitoes are attracted to carbon dioxide, heat and moisture, not to sound or light [48]. Mosquito nets [49], [50], [51] an effective barrier when the entrance and exit areas are not breached by human traffic. Moreover, the coverage is limited to the area that the net covers and Mosquito Magnet® trap [52] a Mosquito Magnet trap significantly decreases the mosquito population by killing enough mosquitoes to interrupt their breeding cycle. This mosquito trap is made to mimic a warm-blooded creature by giving off carbon dioxide and a sweat scent as well as heat. These devices are still in the research and development phase.

As illustrated from the epistle of temporary control measures, success can be as low as 2%, the mosquito control methods may only be effective at repelling or preventing bites, not actually killing the mosquitoes; furthermore, the protected area is small and there is a limit to the amount of protection time.

Therefore, there is great urgency for the development and implementation of mosquito vector control interventions until a satisfactory solution is attained. [53] reports the importance of mathematical models as a first step to assess control strategies and the efficacy of proposed methods prior to implementation. [54] proposed a system of ordinary differential equations to model the interactions among the larval and adult stages of mosquitoes and water mites and [55] have developed mathematical models to show that it is a crucial element in developing optimised control techniques, especially to understand the pest population dynamics, which controls the strength of the disease vector.

Natural control methods

Due to the ineffectiveness of the control measures that we have reviewed, it is imperative to develop a Natural Control Model (NCM), which is based on the deployment of mosquito predators. *Odonata* and *Toxorhynchites* are both naturally beneficial predators of all classes of mosquitoes [56]; hence, we investigate their effectiveness in the following model.

The NCM concept provides a general opportunity for the control of all species of mosquitoes as it is based on the interaction between the population of all species of mosquito adults and its life cycle stages (egg, larvae and pupae) and the naturally beneficial predator *Odonata* adult and its life cycle stages (egg and nymph) and *Toxorhynchites* adult and its life cycle stages. Fig. 1 illustrates the predatory action of *Odonata* on the mosquitoes and *Toxorhynchites* and the predatory action of *Toxorhynchites* on the disease vector mosquito larva. The green arrows indicate the predation of *Odonata* nymphs on pest mosquito and *Toxorhynchites*: eggs, larvae and pupae. The orange arrows indicate the predation of *Odonata* adults on disease vector mosquitoes and *Toxorhynchites* adults. The red arrow indicates predation of *Toxorhynchites* on disease vector mosquito larvae.




Fig.1The predatory action of *Odonata* and *Toxorhynchites* on Mosquitoes

III. MATERIALS AND METHODS

To understand the mosquito challenge, time was spent assessing the problems in Atimbo a residential area, located in Calabar; the capital city of Cross River State, Southern Nigeria. A more than 100 km² residential area is filled with ideal breedingplaces for mosquitoes, open grassy areas are found to feature unmanaged gardens and tree waste, discarded food waste, recyclable containers, flower pots, birdbaths, water dishes, cans, gutters, tires, buckets, marsh areas and puddles in the drainage areas, as illustrated in Fig. 2



Fig. 2 Mosquito breeding zones – Atimbo, Calabar

Because the population of insects and mosquitoes observed in the Atimbo area were so numerous, we collected some species samples and presented them to the pest detection system, developed by [57]. The results obtained from the detection system confirmed the presence of more than one species of disease vector mosquito, see Fig. 3.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014




Fig. 3 Different species of mosquito identified by the detection system

Model of the interaction between the mosquitoes and predators

The *Odonata* adult and nymph are recommended because the *Odonata* nymph is very effective in reducing the mosquito population by eating the mosquito: eggs, larva and pupa, see Fig. 1. *Odonata* adults are very important predators and a valuable ally for humanity as they feed on adult mosquitoes, especially when their populations are in abundance. [58] experimentally confirmed the feeding ability of the *Odonata* nymph for the control of several species of mosquitoes. [59] discussed the agility of the dragonfly compared to mosquitoes and houseflies as thus: “the dragonfly is many times the size of a mosquito or a housefly and only needs to flap its wings a mere 30 times a minute when compared to a mosquito’s 600 times a minute and the housefly’s 1000 flaps a minute to maintain flight with peak manoeuvrability. Such is its power that the dragonfly is equipped with a low-energy, high speed capability; very few insects can escape its basket shaped grabbing limbs that it uses to clutch on to its prey before crushing them into a gooey mass, with its powerful mandibles and swallowing it”.

Toxorhynchites are unique in that none feed on blood and, unlike many other mosquitoes, they are harmless to mankind. The larvae of all *Toxorhynchites* are predaceous on other mosquito larvae or small aquatic arthropods, they are therefore beneficial to humankind, see Fig. 1. They have the ability to consume 10 to 20 mosquito larva in a day and up to 5000 during their whole larva stage [60], [61], [62].

Our goal is population reduction as recommended by [63], [64]. Therefore the following equations provide a dynamic model of the evolving disease vector mosquito life stages, *Odonata* and its life cycle stages and *Toxorhynchites* adult and its life cycle stages per square kilometre. The model includes reproduction, mortality and predation using Weibull probability distribution function and Pascal (negative binomial) distributions.

The Pascal (negative binomial) distribution evaluates the distribution of the number of trials needed to get a fixed number of successes (r) [65], [66]. The negative binomial distribution has been found to be useful in several areas of study, including the fitting of several frequency distributions of different biological data [67]. The enquiry into the frequency distribution of multiple events with particular reference to the occurrence of multiple attacks of disease or repeated accidents [68]. The consideration of the problem of mortality curve fitting over the entire life range [69], [70] analysed organism frequency count data using the negative binomial distribution.

$$\frac{dN_m^e}{dt} = \beta_m N_m^h - \varepsilon_m N_m^e - a N_m^e N_d^n - m_m^e N_m^e \tag{Eqn. 1}$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

$$\frac{dN_m^l}{dt} = \varepsilon_m N_m^e - \lambda_m N_m^l - b N_m^l N_d^n - \xi N_m^l N_e^l - m_m^l N_m^l \tag{Eqn. 2}$$

$$\frac{dN_m^p}{dt} = \lambda_m N_m^l - \rho_m N_m^p - c N_m^p N_d^n - m_m^p N_m^p \tag{Eqn. 3}$$

$$\frac{dN_m^h}{dt} = \{ \rho_m N_m^p - \zeta N_m^h N_d^h - m_m^h N_m^h \} \left[N_m^h \left(\frac{K_m^h - N_m^h}{K_m^h} \right) \right] \tag{Eqn. 4}$$

$$\frac{dN_e^e}{dt} = \beta_e N_e^h - \varepsilon_e N_e^e - e N_e^e N_d^n - m_e^e N_e^e \tag{Eqn. 5}$$

$$\frac{dN_e^l}{dt} = \varepsilon_e N_e^e - \lambda_e N_e^l - f N_e^l N_d^n - m_e^l N_e^l \tag{Eqn. 6}$$

$$\frac{dN_e^p}{dt} = \lambda_e N_e^l - \rho_e N_e^p - g N_e^p N_d^n - m_e^p N_e^p \tag{Eqn. 7}$$

$$\frac{dN_e^h}{dt} = \{ \rho_e N_e^p - \tau N_e^h N_d^h - m_e^h N_e^h \} \left[N_e^h \left(\frac{K_e^h - N_e^h}{K_e^h} \right) \right] \tag{Eqn. 8}$$

$$\frac{dN_d^e}{dt} = \beta_d N_d^h - \varepsilon_d N_d^e - m_d^e N_d^e \tag{Eqn. 9}$$

$$\frac{dN_d^n}{dt} = \varepsilon_d N_d^e - \mu_d N_d^n - m_d^n N_d^n - p_d N_d^n \tag{Eqn. 10}$$

$$\frac{dN_d^h}{dt} = \{ \mu_d N_d^p - m_d^h N_d^h \} \left[N_d^h \left(\frac{K_d^h - N_d^h}{K_d^h} \right) \right] \tag{Eqn. 11}$$

where:

$N_m^h, N_m^e, N_m^l, N_m^p$ = Population density of pest mosquito: adult, egg, larvae and pupae.

$N_e^h, N_e^e, N_e^l, N_e^p$ = Population density of *Toxorhynchites*: adult, egg, larvae and pupae.

N_d^h, N_d^e, N_d^n = Population density of *Odonata*: adult, egg, nymph.

K_m^h, K_e^h, K_d^h = Population carrying capacity of the environment for adult: mosquito, *Toxorhynchites*, *Odonata* - respectively.

$m_m^h, m_m^e, m_m^l, m_m^p$ = Pest mosquito mortality rate: adult, egg, larvae and pupae - respectively.

$m_e^h, m_e^e, m_e^l, m_e^p$ = *Toxorhynchites* mortality rate: adult, egg, larvae and pupae - respectively.

m_d^h, m_d^e, m_d^n = *Odonata* mortality rate: adult, egg and nymph - respectively.

ξ = frequency with which an *Toxorhynchites* larva finds and eats a mosquito larva

ζ = frequency with which a dragon fly (*Odonata*) adult finds and eats an adult mosquito.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

τ = frequency with which an *Odonata* adult finds and eats an adult *Toxorhynchites* .

a, b, c = frequency with which an *Odonata* nymph finds and eats a mosquito prey: eggs, larvae and pupae - respectively.

e, f, g = frequency with which a *Odonata* nymph finds and eats an *Toxorhynchites* prey: eggs, larvae and pupae - respectively.

$\beta_m, \beta_e, \beta_d$ = Number of eggs per day from: mosquito, *Toxorhynchites*, *Odonata*

$\epsilon_m, \epsilon_e, \epsilon_d$ = Fraction of eggs hatching into: mosquito larvae, *Toxorhynchites* larvae, *Odonata* nymph

μ_d = Fraction of nymphs changing into *Odonata*

λ_m, λ_e = Fraction of larvae changing to pupae: mosquito, *Toxorhynchites* - respectively

ρ_m, ρ_e = Fraction of pupae turning into: mosquitos and *Toxorhynchites*

The proposed model consists of eleven simultaneous non-linear, ordinary differential equations (1) to (11), which are solved using a 4th order Runge–Kutta method as described by [71], [72], [73]. Using the average life span of all the insect life cycle stages and their mortality rates as described in the previous works of [74]. The following results were obtained from the combination of tables 1, 2 and 3, and the Weibull probability distribution function to determine the various mortality rates of the pests and predators; for the detailed procedure refer to [75].

TABLE I.

Life cycle of the *Odonata* predators as reported by [76],[77]

Odonata (<i>Odonata</i>) Life cycle & expectancy in days	
Maximum no of eggs per day	50
Life expectancy of egg	12-16
Life expectancy of nymph	56-1516
Life expectancy of adult	14-60

TABLE II.

The life span of several classes of Mosquitoes; derived from the works of [8], [10], [78]

Mosquitoes Life cycle & expectancy in days	
Maximum no of eggs per day	100-250
Life expectancy of adult mosquito	4-30
Life expectancy of egg	2-3
Life expectancy of larva or (wigglers)	7-14
Life expectancy of pupa	2-7

TABLE III.

The lifespan & average lifespan of *Toxorhynchites* life cycle as reported by [62], [79], [80], [81] [82].

Toxorhynchites predatory life span in days	
No of eggs laid	8.26
Egg incubation period	1-4
Larval incubation period	4-79
Pupa incubation period	4 - 12
Adult life span	13-107
Daily number of eggs per female	2.52 -14

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

IV. FREQUENCY OF CAPTURE

To determine the frequency with which predators capture prey, we used a negative binomial distribution function (pdf), equation 12 and the negative binomial cumulative distribution function (cdf) equation 13, and the mean equation 14.

$$y = f(x|r, p) = \binom{r+x-1}{x} p^r (1-p)^x I_{(0,1,2,\dots)}(x) \tag{Eqn. 12}$$

$$y = F(x|r, p) = \sum_{i=0}^x \binom{r+i-1}{i} p^r (1-p)^i I_{(0,1,2,\dots)}(i) \tag{Eqn. 13}$$

The mean of the probability distribution is $\eta = \frac{r}{p}$ Eqn. 14

Where eqn. 12 and eqn.13 returns the negative binomial pdf and cdf at each of the values of 'x' using the corresponding number of successes, 'r' and probability of success in a single trial, 'p'. Where X is the number of trials needed to achieve a particular success rate 'r'

This models the scenario for the successive random trials that the predator undertakes, with each predation attempt having a probability of success 'p'. The number of attempts that the predator must perform in order to capture a given number of prey r has a negative binomial distribution where 'I' is the indicator function, which ensures that 'r' only adopts integer values.

Therefore, considering the life span of the predators *Toxorhynchites* larvae, and *Odonata* adult and nymphs and their feeding habits extracted from the research works as shown above. The frequency of capturing their prey can be modelled using the negative binomial function as illustrated in Fig. 4.




Fig. 4 The capturing ability of the predators

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

V. EXPERIMENTAL RESULTS

Scenario I

With no initial infestation, the area is invaded by over two million adult mosquitoes, each laying an average of 175 eggs per day. The results obtained in Fig. 5 illustrate the reproductive capability of a typical pest mosquito, which has had its blood meal. Since there were no control measures in place to manage the mosquito growth rate, within the space of 20 days the mosquito population density reaches its carrying capacity of 4.0×10^7 .




Fig. 5 Population of adult mosquitoes in favourable conditions, starting with zero population of: eggs, larva & pupa . The graph shows the reproduction rate of a typical female mosquito pest after obtaining a blood meal. The adult laid its eggs which transform into larvae then to pupae and finally into adults.

Scenario II.

This illustrates how an already established infestation level develops, as would be the case in the conditions around the residential areas of the Atimbo community, illustrated by Fig. II. We estimate the mosquito (adult, eggs, larvae and pupae) population densities to be at least above 2,000,000 females per km², with each female mosquito laying an average of 175 eggs per day. In the absence of any control measure, the simulated results are shown in Fig. 6.

The result of Fig. 6 shows a great increase in the population of the mosquito: eggs, larvae, pupae - from 2,200,000 to 2.33×10^9 eggs; 1.63×10^9 larvae, 1.07×10^9 pupae and the adult population reaches the carrying capacity of 4.0×10^7 in the environment within 6 days. It is hardly surprising that the health of people living in this dwelling area is very poor.

Scenario III.

The model was set up with different populations of *Odonata* and *Toxorhynchite*. The results reported in Fig.7 are for a starting population of 200 *Odonata* adult and its life cycle stages and 100 adult *Toxorhynchites* and its life cycle stages.

In Fig. 7 with the introduction of 200 *Odonata* adult and all its life cycle forms and 100 *Toxorhynchites* adults and its life cycle stages the mosquito population density was controlled; the population was observed to drop from a peak of 1.35×10^9 to 560 eggs, 8.14×10^8 to 372 larvae, 4.87×10^8 to 207 pupae and 2.90×10^7 to 4 adults in 90 days.




Fig. 6 Mosquito population in the absence of any control measures or predators. The graph of Fig.6 demonstrates the reproduction rate of the female mosquitoes in an established infestation where there is an initial starting population of all the life cycle stages. Because of the initial starting population, the adult pest reaches the carrying capacity earlier when compared to Fig. 5.




Fig. 7 Effect of the introduction of *Toxorhynchites*: adult, eggs, larvae & pupae and *Odonata*: adult, eggs & nymph on mosquitoes infestation. The graph of Fig. 7 shows the semilog plot of the control efficiency of the combined predators when introduced into the mosquito infested habitat. From the legends the black colour lines with the title Mosq., represents the mosquito population of the adults and its life cycle stages in the presence of the predator *Odonata* shown with the red coloured lines and the predator *Toxorhynchites* (Toxo), shown with magenta coloured lines.

There is predation of *Toxorhynchites* adults and its life cycle stages by *Odonata*, which is evident from the results of Fig.7, this is incorporated into equations (5) to (8). 100 *Toxorhynchites* adult and life cycle stages (egg, larvae and pupae) were introduced into the mosquitoes infested habitat, the maximum population observed for the *Toxorhynchites* eggs, larvae and pupae were 905 eggs, 404 larvae and 109 pupae for the first 30 days. Out of the total population of 100 *Toxorhynchites* deployed, the population at 90 days is 148 eggs, 33 larvae, 17 pupae and 5 adults. These findings are consistent with the experimental work of [83], [84], [85], which reports that predation occurs between predators.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

The results of Fig. 5 and Fig. 6 show that mosquito have the ability to multiple uncontrollably especially if a blood meal is obtained as discussed earlier. When any pest mosquito specie is sighted action should be taken.

The result of Fig. 7, illustrate a successful reduction of the population of mosquitoes as the predators established their population in the environment. The result demonstrates the possibility of obtaining a healthier environment with the deployment of *Odonata* and *Toxorhynchites* species into mosquito infested environments.

The combination of predators controls the mosquito population and demonstrates the great potential of this strategy to significantly reduce the mosquito life cycle stages in a reasonable number of days. This duplex approach requires fewer *Odonator* predators to be deployed, reducing the resources required to achieve an impressive result. This proposal considers both long and short term permanent approach to resolve the problems of mosquitoes. Further research in this area will consider an immediate solution along side the long term permanent solutions.

VI. CONCLUSION

It is observed that as long as an environment is left uncared for, it will definitely become a breeding ground providing many mosquito hatcheries. When an environment is occupied with millions of mosquitoes, no preventive measures can cure or manage the mosquito pest. Therefore it is recommended that in addition to managing the environment and preventing it from becoming a breeding zone, a permanent control measure with a combination of predators should be employed.

Breeding, then deploying beneficial insect predators like the *Odonata* and *Toxorhynchites* will transform a hostile environment into a zone habitable by humans, as these predators have a great ability to control mosquitoes at low cost and in a short amount of time. We also advise the restoration of natural habitats that will attract the visits of important naturally beneficial insects.

Finally patient must be exercised to breed beneficial insects as it takes several days before significant results will be noticed and to achieve a mosquito free environment. Any form of insecticides must be avoided in order to preserve the life span of the predators. The goal should be a world free of: disease vector mosquito species and chemical pesticides.

REFERENCES

- [1] Syed,Z. Leal, W. S., "Acute olfactory response of Culex mosquitoes to a human- and bird-derived attractant", Proceedings of the National Academy of Sciences **106** (44): 18803. 2009. doi:10.1073/pnas.0906932106.
- [2] Hill,R. S. Hansson,S.B. and Ignell, R., "Characterization of Antennal TrichoidSensilla from Female Southern House Mosquito, *Culexquinquefasciatus* Say", *Chemical Senses* (Oxford University Press) **34** (3): 231–252. 2009. doi:10.1093/chemse/bjn080.
- [3]SkeeterBite, "The Life Cycle of Mosquitoes",2007. http://www.skeeterbite.info/article_3.html. (Retrieved 21/02/14)
- [4]Goff, G. Le., Bregues, C. and Robert, V., "Stegomyia mosquitoes in Mayotte, taxonomic study and description of *Stegomyiapia* n. sp.". *Parasite***20**: 31, 2013. doi:10.1051/parasite/2013030.
- [5] Scholte, J. E., Schaffner, F., "Waiting for the tiger: establishment and spread of the *Aedesalbopictus* mosquito in Europe", In Takken, W.; Knols, B. G. J. Emerging pests and vector-borne diseases in Europe 1. Wageningen Academic Publishers. ISBN 978-90-8686-053-1. 2007.
- [6] Hochedez, P. S., Debruyne, M., Bossi, P., Hausfater, P., Brucker, G., Bricaire, F., Caumes,E., "Chikungunya Infection in Travelers", *Emerging Infectious Diseases* **12** (10): 1565–1567.2006. doi:10.3201/eid1210.060495. ISSN1080-6040
- [7] Reinert, F. J., Harbach, R. E., and Kitching, I. J., "Phylogeny and classification of Aedini (Diptera: Culicidae), based on morphological characters of all life stages", *Zoological Journal of the Linnean Society***142** (3): 289–368.2004. doi:10.1111/j.1096-3642.2004.00144.x.
- [8] Kim, C. H.,Wilkerson,L.P., and O'guinn, K. L., "New Records and Reference Collection of Mosquitoes on Jeju Island, Republic of Korea", *Entomological Research*. **1** **35**: 55–66. 2005. Retrieved 15 Feb. 2014.
- [9]Bay,D.E. and Harris, R.L., "Introduction to Veterinary Entomology", A Guide to Livestock Insects:43-51. Robert Harris Publisher. 1988.
- [10] Adham,F. K. and Leaf, H. L., "Establishment of a colony of the mosquito *Culisetalongiareolata* under laboratory conditions", *Cellular and Molecular Life Sciences (CMLS)***38** (12): 1498–1499.1982. doi:10.1007/BF01955792.
- [11] Lehrer, S., "Anopheles mosquito transmission of brain tumor", *Medical Hypotheses* **74** (1): 167–168. 2010. doi:10.1016/j.mehy.2009.07.005.
- [12] Wilcox,B. A.,and Ellis, B., "Forests and emerging infectious diseases of humans", *Unasylva***57**. 2006. ISSN0041-6436
- [13]Souza,De, Renato, P.,Foster, G. P.,Sallum, M. A., Coimbra, T.L.M., Maeda, A. Y.,Silveira,V. R., Moreno,E. S., Silva,F. G. da, Rocco, I. M., Ferreira,I. B., Suzuki,A., Oshiro,F. M.,Petrella, S. M.C.N., Pereira, L. E., Katz, G.,Tengan,C. H.,Siciliano, M. M., dos SantosC. L.S. "Detection of a new yellow fever virus lineage within the South American genotype I in Brazil". *J Med Vol.* **82** (1): 175–185. 2010. doi:10.1002/jmv.21606
- [14] Whitehorn, J., and Farrar, J., "Dengue". *Br. Med. Bull.***95**: 161–73. 2010. doi:10.1093/bmb/ldq019.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

- [15] Reiter, P., "Yellow fever and dengue: a threat to Europe?", *Euro Surveill* **15** (10): 19509. (2010-03-11).
- [16] Caglioti, C., Lalle, E., Castilletti, C., Carletti, F., Capobianchi, M. R. and Bordi L., "Chikungunya virus infection: an overview", *The new microbiologica* **36** (3): 211–27.2013.
- [17] Hawman, D. W., Stoermer, K. A., Montgomery, S. A., Pal, P., Oko, L., Diamond, M. S. and Morrison, T. E., "Chronic joint disease caused by persistent chikungunya virus infection is controlled by the adaptive immune response", *Journal of Virology* **87** (24): 13878–88. (Dec 2013). doi:10.1128/JVI.02666-13
- [18] Ghoshal, A., Das, S., Ghosh, S., Mishra, M. K., Sharma, V., Koli, P., Sen, E. and Basu, A., "Proinflammatory mediators released by activated microglia induces neuronal death in Japanese encephalitis", *Glia* **55** (5): 483–96. 2007. doi:10.1002/glia.20474
- [19] Schiøler, K. L., Samuel, M. and Wai, K. L., "Vaccines for preventing Japanese encephalitis", *Cochrane Database Syst Rev* (3): CD004263. 2007. doi:10.1002/14651858.CD004263.
- [20] Ginsberg L., "Difficult and recurrent meningitis", *Journal of Neurology, Neurosurgery, and Psychiatry*. 75 Suppl 1 (90001): 16–21. (March 2004). doi:10.1136/jnnp.2003.034272
- [21] De Van, D. B., de Gans, J., Tunkel, A. R., and Wijdicks, E. F., "Community-acquired bacterial meningitis in adults. The New England Journal of Medicine" **354** (1): 44–53. 2006. doi:10.1056/NEJMr052116
- [22] William, J., Berger, T., and Elston, D., "Andrews' Diseases of the Skin", *Clinical Dermatology* (10th ed.). Saunders. p. 150. 2005. ISBN0-7216-2921-0.
- [23] Champion, R. H., Roberts, S. O. B., Carpenter, R. G., and Roger, J. H., "Urticaria and Angio-Oedema". *British Journal of Dermatology* **81** (8): 588–97. 1969. doi:10.1111/j.1365-2133.1969.tb16041.x
- [24] Nash, D., Mostashari, F., Fine, A., "The outbreak of West Nile virus infection in the New York City area in 1999", *N. Engl. J. Med.* **344** (24): 1807–14. 2001. doi:10.1056/NEJM200106143442401
- [25] Klenk, K., Snow, J., Morgan, K., Bowen, R., Stephens, M., Foster, F., Gordy, P., Beckett, S., Komar, N., Gubler, D., and Bunning, M., "Alligators as West Nile Virus Amplifiers", *Emerging Infectious Diseases* **10** (12): 2150–2155. 2004. doi:10.3201/eid1012.040264
- [26] Cancrini, G., Frangipane, D., Regalbono, A., Riccia, I., Tessarin, C., Gabrielli, S., and Pietrobelli, M., "Aedes albopictus is a natural vector of Dirofilaria immitis in Italy", *Veterinary Parasitology* **118** (3–4): 195–202. 2003. doi:10.1016/j.vetpar.2003.10.011. ISSN0304-4017.
- [27] Facchinelli, L., Valerio, L., Pombi, M., Reiter, P., Costantini, C., and Della A. T., "Development of a novel sticky trap for container-breeding mosquitoes and evaluation of its sampling properties to monitor urban populations of Aedes albopictus", *Medical and veterinary entomology* **21** (2): 183–95. 2007. doi:10.1111/j.1365-2915.2007.00680.x.
- [28] Gama, R. A., Silva, E. M., Silva, I. M., Resende, M. C., and Eiras, Á. E., "Evaluation of the sticky Mosqui TRAP for detecting Aedes (Stegomyia) aegypti (L.) (Diptera: Culicidae) during the dry season in Belo Horizonte, Minas Gerais, Brazil", *Neotropical Entomology* **36** (2): 294–302. 2007. doi:10.1590/S1519-566X2007000200018.
- [29] Meeraus, W. H., Armistead, J. S., and Arias, J. R., "Field comparison of novel and gold standard traps for collecting Aedes albopictus in Northern Virginia", *Journal of the American Mosquito Control Association* **24** (2): 244–248. 2008. doi:10.2987/5676.1
- [30] Karch, S., Monteny, N., Jullien, J. L., Sinègre, G., and Coz J., "Control of Culex pipiens by Bacillus sphaericus and role of nontarget arthropods in its recycling", *J. Am. Mosq. Control Assoc.* 1990 Mar;6(1):47-54. 1990.
- [31] Charles, F. C., Nielson, L., and Deleclus, A., "Bacillus Sphaericus Toxins: Molecular Biology and Mode of Action", *Annual Review of Entomology* Vol. 41: 451-472 (January 1996) DOI: 10.1146/annurev.en.41.010196.002315
- [32] U.S. Environmental Protection Agency. Office of Pesticide Programs. "Bacillus sphaericus serotype H5a5b strain 2362 (128128) Fact Sheet"
- [33] P. E. N. Yufang, S. O. N. Jinzhang, T. I. A. N. Guangzhao, X. U. E. Qunli, G. E. Fengxiang, Y. A. N. Jialun and S. H. I. Qi (1998). Field evaluations of *Romanomermis yunanensis* (Nematoda: Mermithidae) for control of Culicinae mosquitoes in China (1) *Fundam. appl. Nematol.*, 1998, 21 (3), 227-232
- [34] Dong-Kyu L. E. E., "Biological Control of *Culex pipiens pallens* (Diptera, Culicidae) by the Release of Fish Muddy Loach, *Misgurnus mizolepis* in Natural Ponds, Korea", *Entomological Research* Vol. 32 (1), pg 43–47. 2002. DOI: 10.1111/j.1748-5967.2002.tb00008.x
- [35] Machida, H., "The inhibitory effect of a combination of imidacloprid and permethrin on blood feeding by mosquitoes in dogs raised under outdoor conditions", *Vet Parasitol* 154(3-4):318. 2008.
- [36] Scholte, E. J., Ng'habi, K., Kihonda, J., Takken, W., Paaijmans, K., Abdulla, S., Killeen, G. F., and Knols, B. G. J., "An entomopathogenic fungus for control of adult African malaria mosquitoes", *Science* 308: 1641-1642. 2005.
- [37] Faithpraise, F. O., Chatwin, C. R., Obu, J., Olawale, B., Young, R., Birch, P., "Sustainable Control of Anopheles Mosquito Population". *Environment, Ecology & Management*, Vol 3(1). 1-19. (2014a).
- [38] B. J. Sina & K. Aultman. (2001) Resisting resistance. *Trends Parasitol.* 17, 305–306
- [39] Anderson, M., "Mosquito Management: How to Avoid Mosquitoes in your Backyard" (2012). <http://blog.epa.gov/greeningtheapple/2012/05/mosquito-management-how-to-avoid-mosquitoes-in-your/> (Retrieved 20/10/2013)
- [40] Hadley, D., "How to Control Mosquitoes in Your Yard and Home. Limit Habitat for Mosquitoes" 2014. <http://insects.about.com/od/insectpests/a/mosquitocontrol.htm> (Retrieved 16/02/2014)
- [41] U. S. Environmental Protection Agency, "Citronella (Oil of Citronella) (021901) Fact Sheet". 16 February 2011. (Retrieved 1 July, 2011)
- [42] Kazuhiko, N., Alzoreky, S. N., Yoshihashi, T., Nguyen, T. H., and Trakoontivakorn G., "Chemical Composition and Antifungal Activity of Essential Oil from *Cymbopogon nardus* (Citronella Grass)", *Japan International Research Center for Agricultural Sciences* **37** (4): 249–52. (October 2003). INIST:15524982
- [43] Mumcuoglu, K. Y., Magdassi, S., Miller, J., Ben-Ishai, F., Zentner, G., Helbin, V., Friger, M., Kahana, F., and Ingber, A., "Repellency of citronella for head lice: Double-blind randomized trial of efficacy and safety", *The Israel Medical Association Journal* **6** (12): 756–9. 2004.
- [44] Tuetun, B., Choochote, W., Pongpaibul, Y., Junkum, A., Kanjanapothi, D., Chaitong, U., Jitpakdi, A., Riyong, D., Wannasan, A., Pitasawat, B., "Field evaluation of G10, a celery (*Apium graveolens*)-based topical repellent, against mosquitoes (Diptera: Culicidae) in Chiang Mai province, northern Thailand", *Parasitology Research* **104** (3): 515–521. 2008. doi:10.1007/s00436-008-1224-9.
- [45] Enayati, A. A., Hemingway, J., Garner P., "Electronic mosquito repellents for preventing mosquito bites and malaria infection", *Cochrane Database Syst Rev.*, 2007 18; CD005434

- [46]Freudenrich, C.,“BugZappers. How Stuff Works”,<http://home.howstuffworks.com/bug-zapper.htm> Retrieved 2014-02-16
- [47] Science Daily: “Electric Bug Zappers Are Useless For Controlling Mosquitoes, Says UF/IFAS” Pest Expert 30, 1997. <http://www.sciencedaily.com/releases/1997/07/970730060806.htm> (Retrieved 16/02/14)
- [48] Lewis, D.,“Bug Zappers are Harmful, Not Helpful”. Horticulture and Home Pest News (Iowa State University). IC-475 (15).(1996-06-14). Retrieved 2014-02-16.
- [49] Yakob, L.,and Guiyun Y.,“Modeling the Effects of Integrating Larval Habitat Source Reduction and Insecticide Treated Nets for Malaria Control”,PLoS ONE (Public Library of Science) 4 (9).2009. doi:10.1371/journal.
- [50] Killeen,G. F.,and Smith, T. A.,“Exploring the contributions of bed nets, cattle, insecticides and excito-repellency to malaria control: A deterministic model of mosquito host-seeking behaviour and mortality”, American Journal of Tropical Medicine and Hygiene.2007.
- [51] Cohen,J.,and Dupas, P.,“Free Distribution or Cost-Sharing? Evidence from a Randomized Malaria Prevention Experiment”.. Quarterly Journal of Economics 125 (1): 24.2010.
- [52] Williams,C. R., Long, S. A., Webb, C. E.,Bitzhenner,M.,Geier, M., Russell, R. C., and Ritchie, S. A. (2007). *Aedesegypti*Population “Sampling using BG-Sentinel Traps in North Queensland Australia: Statistical Considerations for Trap Deployment and Sampling Strategy”, J. Med. Entomol. 44(2): 345D350 .2007.
- [53] Oluwagbemi, O. O., Fornadel, C. M., Adebiyi, E. F., Norris, D. E., and Rasgon, J. L., “ANOSPEX: A Stochastic, Spatially Explicit Model for Studying Anopheles Metapopulation Dynamics”, PLoS ONE 8(7): e68040. 2013. doi:10.1371/journal.pone.0068040
- [54] Esteva, L., Rivas, G., and Yang, H. M., “Modelling parasitism and predation of mosquitoes by water mites”,J. Math. Biol. (2006) 53:540–555 DOI 10.1007/s00285-006-0020-3
- [55] Faithpraise, F. O., Chatwin, C. R., Young, R. C. D.,and Birch, P.M.,“Timely Control of Aphis craccivora Using an Automatic Robotic Drone management system (ARDMS)”, Technical Report, TR/SU/FF/130617, 17 June 2013, Page 1-19(2014).
- [56] Focks, D. A., Sackett, S. R., and Bailey D. L., “Field experiments on the control of *Aedesegypti* and *Culexquinquefasciatus* by *Toxorhynchitesrutilusrutilus* (Diptera: Culicidae)”,Journal of medical entomology 19 (3): 336–339. 1982.
- [57] Faithpraise, F., Birch,P., Young,R.,Obu,J.,Faithpraise, B., and Chatwin C., “Automatic plant pest detection & recognition using k-means clustering algorithm & correspondence filters”, International Journal of Advanced Biotechnology and Research, Vol. 4, Issue 2, 2013, pp 1052-1062. 2013. ISSN 0976-2612
- [58] Singh, R. K., Dhiman, R. C., and Singh,S. P.,“Laboratory studies on the predatory potential of dragon-fly nymphs on mosquito larvae”,Journal of Communicable Diseases. 35 (2): 96–101.2003.
- [59] Dragonfly site: “What Do Dragonflies Eat?”,<http://www.dragonfly-site.com/what-do-dragonflies-eat.html> (Retrieved 10/03/14)
- [60] Steffan,W. A., and Evenhuis, N. L.,“Biology of *Toxorhynchites*”, Annual Review of Entomology 26: 159.1981. doi:10.1146/annurev.en.26.010181.001111
- [61] Focks, D. A., Sackett, S. R., and Bailey D. L., “Field experimentes on the control of *Aedesegypti* and *Culexquinquefasciatus* by *Toxorhynchitesrutilusrutilus*Diptera: Culicidae”, Journal of Medical Entomology, 193: 336-339. 1982.
- [62] Goettle, B. J.,and Adler, H. P., “Elephant (or Treehole) Predatory mosquito”,1999. www.dnr.sc.gov/cwcs/pdf/Predmosquit (Retrieved 22/02/14)
- [63]Fang,J.,“Ecology: A world without mosquitoes”, Nature (Nature) 466 (7305): 432–4.(July 21, 2010). doi:10.1038/466432a.
- [64] Science Today “Beyond the Headlines Mosquito Eradication”. California Academy of Sciences. 26 July 2010.<http://www.calacademy.org/sciencetoday/mosquito-eradication/> (Retrieved 19 02 2014)
- [65] Villarini, G., Vecchi, G. A., and Smith, J. A., "Modeling of the dependence of tropical storm counts in the North Atlantic Basin on climate indices",Monthly Weather Review 138 (7): 2681–2705. 2010. doi:10.1175/2010MWR3315
- [66]Hilbe, J. M., "Negative Binomial Regression, second edition", 2011 Cambridge, UK: Cambridge University Press
- [67] Bliss, C.I.,“Fitting the negative binomial distribution to biological data”, Biometrics 9:176-200, 1953.
- [68] Greenwood, M.,and Yule, G. U.,“An inquiry into the nature of frequency distributions representative of multiple happenings with particular reference to the occurrence of multiple attacks of disease or of repeated accidents,” Journal of the royal statistical society 83:255-229, 1920
- [69] Hewitt,C.C., Jr.,“The negative binomial applied to the Canadian merit rating plan for individual automobile risks”, Proceedings of the casualty Actuarial Society 47:55-65. 1960.
- [70] Gray, B. C.,“Analysis of frequency count data using negative binomial distribution”, Ecology, 77(8), 1996. Pp.2549-2557
- [71] Klassische, E. F.,“Runge-Kutta-Formeln fünfter und siebenter Ordnung mit Schrittweiten-Kontrolle”, Computing (Arch. Elektron. Rechnen) 4 1969 93-106.
- [72]Dormand,J. R., and Prince, P. J.,“High order embedded Runge-Kutta formulae”, J. Comput. Appl. Math. 7 (1981), no.1, 67-75.
- [73]Schreiber, R.,“MATLAB”,Scholarpedia, 2(7):2929. 2007.
- [74] Faithpraise, F., Idung, J., Chatwin, C., Young, R., Birch, P., “Biological Control of Taro Scarab Beetle (*Papuanauninodis,Coleoptera: Scarabaeidae*) Instars via *scoliid* and *Voriatachinidae*”,International Journal of Applied Biology and Pharmaceutical Technology, Volume 5, Issue 3, in press, 27th April 2014. July –Sept 2014, ISSN: 0976-4550.
- [75] Faithpraise, F., Idung, J., Chatwin, C., Young, R., Birch, P.,Targeting the life cycle stages of the Diamond Back Moth (*PlutellaXylostella*) with three different parasitoid wasps, Technical Report, TR/SU/FF/131202, 02 Nov. 2013, Page 1-18
- [76] Smallshire, D.,and Swash, A., “WILDGuides”, - ISBN 978-1-903657-29-4 .2010.
- [77] The Life-Cycle Of A Odonata. <http://www.british-dragonflies.org.uk/content/biology-ecology>; <http://www.Odonata-site.com/Odonata-facts.html>; Lifespan of Dragonflies. <http://www.desertusa.com/mag98/nov/papr/drangon...> (Retrieved 25/09/13)
- [78] Cross,C.,“The life cycle of *Anopheles* mosquitoes,Female *Anophelesspecies* mosquitoes are the only vector of malaria parasites”,(11/8/04) http://malaria.welcome.ac.uk/doc_WTD023872.html (Retrieved 16/09/13)
- [79] Toma,T.,and Miyagi, I., “Laboratory evaluation of *Toxorhynchitesplendens* (Diptera: Culicidae) for predation of *Aedesalbopictus* mosquito larvae”, Medical and Veterinary Entomology 08/1992; 6(3):281-91

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 3, Issue 5, May 2014

-
- [80] Chohanadisai, L., Benjaphong, N., and Phanthumachinda B., "Laboratory observations on *Toxorhynchites splendens* Wiedemann in Thailand", The Southeast Asian journal of tropical medicine and public health (impact factor: 0.6). 10/1984; 15(3):337-41
- [81] Choochote, W., Jitpakdi, A., Suntaravitun, T., Junkum, A., Rongsriyam, K., and Chaithong U., "A Note on Laboratory Colonization of *Toxorhynchites splendens* by Using an Artificial Mating Technique and Autogenous *Aedes togoi* Larva as Prey", The Journal of Tropical Medicine And Parasitology J. Trop Med Parasitol 2002;25:47-50. Vol 26 (No. 1) June 2003
- [82] Aditya, G., Ash, A., and Saha, G. K., "Predatory activity of *Rhantussikkimensis* and larvae of *Toxorhynchites splendens* on mosquito larvae in Darjeeling", India. J Vector Borne Dis. 2006 Jun;43(2):66-72.
- [83] Polis, G. A., "The evolution and dynamics of intraspecific predation", Annual Review of Ecology and Systematics 12, 225-251 .1981.
- [84] Mehlis, M., Bakker, T. C. M., Engqvist, L., and Frommen, J. G., "To eat or not to eat: egg-based assessment of paternity triggers fine-tuned decisions about filial cannibalism", Proceedings of the Royal Society B-Biological Sciences 277:2627-2635 .2010.
- [85] Volker, H., and Rudolf W., "Impact of Cannibalism on Predator-Prey Dynamics Size-Structured Interactions and Apparent Mutualism", Ecology, 89(6), 2008, pp. 1650-1660 _ 2008 by the Ecological Society of America

BIOGRAPHY

Fina O. Faithpraise was born in Nko- Yakurr Local Government area of Cross River State, Nigeria in 1976. She obtained her first degree at Enugu State University of Science and Technology (ESUT) Nigeria, where she gained Bachelor of Engineering (*computer science and engineering*) in 2001. In 2007, she obtained a Master of Science degree in *engineering physics* from the University of Calabar, Cross River State, Nigeria and started a doctoral research program in the School of Engineering and Informatics (*biomedical engineering*) at the University of Sussex, United Kingdom in 2011.

Fina Faithpraise is also a Lecturer at the University of Calabar, her interest is to find solutions to alleviate pest infestations without resorting to the use of health threatening pesticides.