

Formation of a stable radical by oxidation of a tetraorganoborate

Article (Accepted Version)

Braunschweig, Holger, Krummenacher, Ivo, Mailänder, Lisa, Pentecost, Leanne and Vargas, Alfredo (2016) Formation of a stable radical by oxidation of a tetraorganoborate. *Chemical Communications*, 43. pp. 7005-7008. ISSN 1359-7345

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/60772/>


This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



Formation of a stable radical by oxidation of a tetraorganoborate

Holger Braunschweig,* Ivo Krummenacher,
Lisa Mailänder, Leanne Pentecost and Alfredo Vargas

Oxidation of a tetraorganoborate salt leads to selective formation of the corresponding zwitterionic radical rather than carbon-carbon bond formation of the boron substituents.

Q2

Please check this proof carefully. **Our staff will not read it in detail after you have returned it.**

Translation errors between word-processor files and typesetting systems can occur so the whole proof needs to be read. Please pay particular attention to: tabulated material; equations; numerical data; figures and graphics; and references. If you have not already indicated the corresponding author(s) please mark their name(s) with an asterisk. Please e-mail a list of corrections or the PDF with electronic notes attached – do not change the text within the PDF file or send a revised manuscript. Corrections at this stage should be minor and not involve extensive changes. All corrections must be sent at the same time.

Please bear in mind that minor layout improvements, e.g. in line breaking, table widths and graphic placement, are routinely applied to the final version.

Please note that, in the typefaces we use, an italic vee looks like this: ν , and a Greek nu looks like this: ν .

We will publish articles on the web as soon as possible after receiving your corrections; **no late corrections will be made.**

Please return your **final** corrections, where possible within **48 hours** of receipt, by e-mail to: chemcomm@rsc.org

Queries for the attention of the authors

Journal: ChemComm

Paper: c6cc02916g

Title: Formation of a stable radical by oxidation of a tetraorganoborate

Editor's queries are marked on your proof like this **Q1**, **Q2**, etc. and for your convenience line numbers are indicated like this 5, 10, 15, ...

Please ensure that all queries are answered when returning your proof corrections so that publication of your article is not delayed.

Query reference	Query	Remarks
Q1	For your information: You can cite this article before you receive notification of the page numbers by using the following format: (authors), Chem. Commun., (year), DOI: 10.1039/c6cc02916g.	
Q2	Please carefully check the spelling of all author names. This is important for the correct indexing and future citation of your article. No late corrections can be made.	
Q3	Please check that the inserted CCDC number is correct.	
Q4	Ref. 1d: Please provide the page (or article) number(s).	

Formation of a stable radical by oxidation of a tetraorganoborate†

Cite this: DOI: 10.1039/c6cc02916g

Received 7th April 2016,
Accepted 4th May 2016


DOI: 10.1039/c6cc02916g

www.rsc.org/chemcomm

Herein, we describe the selective formation of a stable neutral spiroborate radical by one-electron oxidation of the corresponding tetraorganoborate salt $\text{Li}[\text{B}(\text{C}_6\text{H}_4)_2]$, formally containing a tetrahedral borate centre and a *s-cis*-butadiene radical cation as the spin-bearing site. Spectroscopic and computational methods have been used to determine the spin distribution and the chromism observed in the solid state.

The oxidation of tetraaryl-, -alkenyl or -alkyl borates with the general formula $[\text{BR}_4]^-$ usually leads to the formation of C–C coupled products by skeletal rearrangements.^{1,2} Oxidation thus represents a versatile method for the synthesis of disubstituted alkynes or biaryl compounds.³ As depicted in Scheme 1, such reactions are also a feature of spiroborates, which in the case of a spirocyclic borafluorene derivative (**1**) afford a bis(biaryl) compound.^{4,5} In general, these reactions can be initiated by a variety of oxidising agents or reaction conditions, such as photo- or electrochemically.⁶ In a few instances, $[\text{BR}_4]^*$ radicals have been suggested as intermediates in the oxidative ligand coupling process, but never directly observed.⁷ Herein, we show with a spiroborate (**2**) that the oxidation can be directed to the exclusive formation of such a boranyl radical species.


Spiroborate **2** was synthesised as previously reported by our group by salt elimination of two equivalents of 1,2-dilithio-1,2,3,4-tetraphenylbutadiene with $\text{BF}_3 \cdot \text{Et}_2\text{O}$.⁸ Whereas the redox chemistry of boroles, five-membered boron-containing, antiaromatic and highly Lewis acidic heterocycles,⁹ has been explored extensively,¹⁰ the redox behaviour of the related spiroborate salt **2** had not been investigated so far. As judged by cyclic voltammetry in tetrahydrofuran solution, compound **2**



Scheme 1 Two examples of oxidation reactions of organoborate salts resulting in C–C coupled products.

undergoes an irreversible oxidation at -0.35 V vs. the $[\text{FeCp}_2]/[\text{FeCp}_2]^+$ couple (see ESI†) in the potential window. We wondered if the oxidation step could be realised on a preparative scale and the formed product be isolated. To this end, a stoichiometric amount of the one-electron oxidation agent ferrocenium hexafluorophosphate was added to a solution of **2** in benzene at rt (Scheme 2). Addition of the Fe(III) salt led to an immediate colour change from yellow to dark green and the precipitation of a colourless solid. Upon completion, the ^{11}B NMR resonance of the starting material (**2**, $\delta_{\text{B}} = -1.4$ ppm) disappeared without the occurrence of a new signal. Similarly, the ^1H NMR spectrum of the crude reaction mixture showed only the formation of ferrocene but no additional signals for the oxidation product. After the separation of $\text{Li}[\text{PF}_6]$ and ferrocene, compound **3** was isolated as green amorphous, or alternatively, red crystalline, solids (80% crystalline yield).

The absence of NMR signals suggested the radical character of **3**, which was supported by EPR spectroscopy (*vide infra*). X-ray diffraction of red single crystals of **3** confirmed the



Scheme 2 Oxidation of spiroborate salt **2** with $[\text{FeCp}_2][\text{PF}_6]$ to radical **3** (only one possible resonance form is drawn).

^a Institut für Anorganische Chemie, Julius-Maximilians-Universität Würzburg, Am Hubland, 97074 Würzburg, Germany. E-mail: h.braunschweig@uni-wuerzburg.de

^b Department of Chemistry, School of Life Sciences, University of Sussex, Brighton BN1 9QJ, Sussex, UK

† Electronic supplementary information (ESI) available: Experimental details, additional crystallographic, computational and electrochemical data. CCDC 1469228. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6cc02916g


1
5
10

Fig. 1 Left: Molecular structure of **3** in the solid state. Hydrogen atoms and the ellipsoids of the phenyl residues are omitted for clarity. The ellipsoids are set to 50% probability. For bond lengths and angles see Table S1 in the ESI.† Right: Side view showing the relative arrangement of the two rings.

15

formation of a charge-neutral species, consistent with a one-electron oxidation of spiroborate **2**.⁸ The D_{2d} symmetry is lifted in the oxidised product as can be seen by the changes of the geometric parameters around the spiro boron atom (*cf.* Table S1 in the ESI†). For instance, the two essentially planar BC_5 rings in **3** deviate significantly from a perpendicular arrangement ($79.0(1)^\circ$) and are displaced relative to each other from the twofold axis (Fig. 1, right). In addition, the one-electron oxidation leads to a shortening of the B–C distances of about 0.1 Å (**3**: $\text{O}_{B-C} = 1.535 \text{ \AA}$ vs. **2**: $\text{O}_{B-C} = 1.631 \text{ \AA}$) and to a reduction in the bond length alternation of the bicyclic rings. The formal double bonds of the borate anion (C1–C2, C3–C4, C5–C6, and C7–C8) are elongated in the oxidised product **3**, consistent with removal of an electron from the HOMO of the anion (**2**) which shows π -binding interactions between these carbon atoms.

20

25

30

The phenyl substituents around the spirocycle are arranged in a propeller-like manner as observed for **2**. However, it is important to note that the phenyl ring at C1 is only tilted by 30.2° with respect to the fused ring plane, whereas the other aryl substituents have torsion angles between 43 and 69° . This might be due to effects of molecular packing or to some degree of spin delocalisation into the aryl π system.

35

40

The continuous-wave ESR spectrum of **3** in benzene solution shows that the unpaired spin density is delocalised into the aryl substituents (Fig. 2). The best-fit parameters for the second-derivative ESR signal ($g_{\text{iso}} = 2.003$) relate to small proton hyperfine couplings (in the range of 0.05 – 0.15 mT) of four equivalent phenyl groups, indicating simultaneous delocalisation of the unpaired spin density over both π planes. The ESR results thus point to a spiroconjugated organic radical with a structure of higher symmetry (D_{2d}) than observed in the solid state.¹¹

45

50

The UV-vis spectrum of **3** differs strongly from that of the borate salt **2**, which shows a lowest absorption maximum at 365 nm ($\epsilon = 34\,490 \text{ L mol}^{-1} \text{ cm}^{-1}$).⁸ In addition to bands in the same wavelength region ($\lambda(\epsilon) = 329 \text{ nm}$ ($24\,894 \text{ L mol}^{-1} \text{ cm}^{-1}$), $\lambda(\epsilon) = 352 \text{ nm}$ ($22\,919 \text{ L mol}^{-1} \text{ cm}^{-1}$)), radical **3** exhibits additional transitions at lower energies ($\lambda(\epsilon) = 463 \text{ nm}$ ($12\,967 \text{ L mol}^{-1} \text{ cm}^{-1}$), $\lambda(\epsilon) = 572 \text{ nm}$ ($2442 \text{ L mol}^{-1} \text{ cm}^{-1}$), $\lambda(\epsilon) = 604 \text{ nm}$ ($2307 \text{ L mol}^{-1} \text{ cm}^{-1}$), $\lambda(\epsilon) = \sim 700.0 \text{ nm}$ ($\sim 750 \text{ L mol}^{-1} \text{ cm}^{-1}$)),

55




Fig. 2 X-band first (grey dotted line) and second (solid black line) derivative ESR spectrum of **3** at room temperature in benzene solution. A simulation of the second derivative spectrum is shown in red. Instrumental parameters: $\nu = 9.379 \text{ GHz}$, modulation frequency = 100 kHz , modulation amplitude = 0.1 G , microwave power = 0.25 mW , conversion time = 40 ms , 4 scans.

1
5
10

15

20

25

30

35

40

45

50

55

which are extended into the near infrared range. The bands between 500 – 900 nm are significantly weaker in intensity than those between 300 – 500 nm and likely can be attributed to intervalence charge transfer transitions. The absorptions from 550 nm to the near IR range are consistent with other organic monoradicals.^{12–14}

Another interesting aspect of this radical species is that its colour is strongly dependent on its physical state. The radical (**3**) is dark green in solution and in its amorphous solid state, whereas it is red in the crystalline state. Presumably, the locked conformation in the crystal structure limits the amount of π conjugation in the molecule, thus leading to a shift of the maximum absorption to shorter wavelengths.

Additional computational studies within the Kohn–Sham density functional theory (DFT) were undertaken. These calculations support the experimental results with respect to the generation of the radical species. Calculated structures of **2** and **3** were in reasonable agreement with those obtained experimentally (see ESI† for Computational details).

Energy minimisation calculations on **3** yielded structures in the gas phase that are distinct from the solvated structures. Furthermore, excited state calculations on the resulting geometries revealed different absorption profiles. These results highly suggest environment-dependent geometries which in turn dictate the observed absorption profile – explaining the experimentally observed differing colours of the species depending on the physical state. Indeed, depending on the structure, *e.g.* solvated structure, lower molecular orbitals are involved in the excitation process. Such observations lead to the idea of environment (*e.g.* solvent)-stabilised charge distribution within the system, which is further supported by computational results as shown by molecular electrostatic potentials (MEP) and natural charges. Further discussions on the theoretical results are given in the ESI.† Moreover, these calculations allowed identification of the experimentally observed transitions:




Fig. 3 MO62X/6-311G* calculated (a) singly-occupied molecular orbital (SOMO), and (b) spin-density distribution for **3** with imposed D_2 symmetry.

HOMO–2 to LUMO (spin-down) at around $15\,500\text{ cm}^{-1}$ (645 nm), and HOMO to LUMO (spin-up) at around $21\,000\text{ cm}^{-1}$ (476 nm).

Another geometry-dependent feature of **3** is that of the spin-density distribution, which unsurprisingly is dictated by the structure. Calculations show that the spin population in the radical is essentially localised on one borole moiety in both the gas and solution phase; the dominant spin density is found on the two carbon atoms adjacent to the boron atom. Imposing D_2 symmetry distributes the spin density evenly on all four carbon atoms adjacent to the boron atom that extends into the phenyl substituents. Furthermore, in this symmetrical configuration the molecular orbitals (MOs) are delocalised across both borole moieties (see Fig. 3). Both the spin density and MO profiles are well in line with experimental results (*vide supra*). The total electronic energy of the symmetrised structure is only slightly (0.6 kcal mol^{-1}) higher than that of the non-symmetrised structure. This could be ascribed in part to interaction of the two perpendicular π planes connected by the boron centre, *via* so-called spiroconjugation.¹⁵

The difference in reactivity of **2** compared to its benzannulated analogue **1**, the latter undergoing coupling of the organic substituents upon oxidation,^{4,5} most likely arises from the stoichiometric use of the outer-sphere oxidant ferrocenium, which can accept precisely one electron from the borate. Isolation of the stable spiro radical (**3**) suggests that a radical mechanism might be operative in the oxidative ligand coupling of other organoborates, as previously proposed.^{4,5,7} The possible resonance forms of the zwitterionic species **3** all consist of a borate anion and a *s-cis*-1,3-butadiene radical cation bearing the unpaired spin density (see Scheme 1). The structure thus draws immediate comparison to simple butadiene radical cations which have attracted considerable theoretical and experimental attention,¹⁶ but have eluded characterisation by X-ray crystallography thus far.¹⁷ Moreover, radical **3** resembles the structures of the neutral radicals reported by Haddon and coworkers,¹⁸ in which two phenalenyl units are attached to a tetrahedral boron atom. In this family of compounds, however, the neutral spiro radical is based on *N,N*- or *N,O*-containing six-membered heterocyclic systems and thus not susceptible to a competing oxidative ligand coupling reaction. Closely related to **3** are also the neutral borocyclic radicals of the general formula

$[(C_6F_5)_2B(O_2C_{14}H_8)]^\bullet$ and $[(C_6F_5)_2B(O_2C_{16}H_8)]^\bullet$ in which the boron atom has two oxygen substituents.¹⁹

In summary, we have shown that by chemical one-electron oxidation of an organoborate, a neutral stable radical of the form $[BR_4]^\bullet$ can be generated. The stability and persistency of the radical species enabled characterisation by X-ray crystallography, besides spectroscopic techniques (ESR and UV-vis). Assisted by DFT and TD-DFT calculations, the striking colour change between the crystalline and the solution/amorphous state could be attributed to structural changes of the spiro compound. Further investigations regarding the intriguing electronic and optical properties of the compound, as well as its reactivity, are underway.

We thank the Deutsche Forschungsgemeinschaft (H. B.), the University of Sussex and the EPSRC (A. V.) for support of this work.

Notes and references

- For references to early work, see: (a) H. W. Spier, *Biochem. Z.*, 1952, **322**, 467–470; (b) D. H. Geske, *J. Phys. Chem.*, 1959, **63**, 1062–1070; (c) D. H. Geske, *J. Phys. Chem.*, 1962, **66**, 1743–1744; (d) A. N. Nesmeyanov, V. A. Sazonova, G. S. Liberman and L. I. Emelyanova, *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 1965, **48**.
- For different outcomes of organoborate oxidations see, for example: (a) S. T. Murphy, C. Zou, J. B. Miers, R. M. Ballew, S. D. Dlott and G. B. Schuster, *J. Phys. Chem.*, 1993, **97**, 13152–13157; (b) A. Y. Polykarpov, S. Hassoon and D. C. Neckers, *Macromolecules*, 1996, **29**, 8274.
- (a) A. Suzuki, N. Miyaura, S. Abiko, M. Itoh, H. C. Brown, J. A. Sinclair and M. M. Midland, *J. Am. Chem. Soc.*, 1973, **95**, 3080–3081; (b) M. Naruse, K. Utimoto and H. Nozaki, *Tetrahedron*, 1974, **30**, 2159–2163; (c) K. Yamada, N. Miyaura, M. Itoh and A. Suzuki, *Tetrahedron Lett.*, 1975, **16**, 1961–1964; (d) A. Suzuki, N. Miyaura, S. Abiko, M. Itoh, M. M. Midland, J. A. Sinclair and H. C. Brown, *J. Org. Chem.*, 1986, **51**, 4507–4511; (e) A. Pelter and R. A. Drake, *Tetrahedron Lett.*, 1988, **29**, 4181–4184.
- J. J. Eisch, J. H. Shah and M. P. Boleslawski, *J. Organomet. Chem.*, 1994, **464**, 11–21.
- J. J. Eisch and R. J. Wilcsek, *J. Organomet. Chem.*, 1974, **71**, C21–C24.
- See, for example: (a) J. L. R. Williams, J. C. Doty, P. J. Gridale, R. Searle, T. H. Regan, G. P. Happ and D. P. Maier, *J. Am. Chem. Soc.*, 1967, **89**, 5153–5157; (b) E. E. Bancroft, H. N. Blount and E. G. Janzen, *J. Am. Chem. Soc.*, 1979, **101**, 3692; (c) J. J. Eisch, M. P. Boleslawski and K. Tamao, *J. Org. Chem.*, 1989, **54**, 1627–1634; (d) S. Boyatzis, J. D. Wilkey and G. B. Schuster, *J. Org. Chem.*, 1990, **55**, 4537–4544; (e) Y. Yasu, T. Koike and M. Akita, *Adv. Synth. Catal.*, 2012, **354**, 3414; (f) G. Duret, R. Quinlan, P. Bisseret and N. Blanchard, *Chem. Sci.*, 2015, **6**, 5366.
- See, for example: (a) P. Abley and J. Halpern, *J. Chem. Soc. D*, 1971, 1238; (b) T. Ishikawa, S. Nonaka, A. Ogawa and T. Hirao, *Chem. Commun.*, 1998, 1209.
- H. Braunschweig, C. Hörl, F. Hupp, K. Radacki and J. Wahler, *Organometallics*, 2012, **31**, 8463–8466.
- For general reviews, see: (a) H. Braunschweig, I. Krummenacher and J. Wahler, *Adv. Organomet. Chem.*, 2013, **61**, 1–53; (b) H. Braunschweig and T. Kupfer, *Chem. Commun.*, 2011, **47**, 10903–10914; (c) A. Steffen, R. M. Ward, W. D. Jones and T. B. Marder, *Coord. Chem. Rev.*, 2010, **254**, 1950–1976.
- For a general review, see: H. Braunschweig and I. Krummenacher, in *Organic Redox Systems: Synthesis, Properties and Applications*, ed. T. Nishinaga, John Wiley & Sons, Inc., 2016, pp. 503–519.
- For an ESR study on a related spirobifluorene compound, see: F. Gerson, B. Kowert and B. M. Peake, *J. Am. Chem. Soc.*, 1974, **96**, 118–120.
- A. Heckmann, C. Lambert, M. Goebel and R. Wortmann, *Angew. Chem., Int. Ed.*, 2004, **43**, 5851–5856.
- L. Fajari, R. Papoular, M. Reig, E. Brillas, J. L. Jorda, O. Vallcorba, J. Rius, D. Velasco and L. Juliá, *J. Org. Chem.*, 2014, **79**, 1771–1777.
- Q. Peng, A. Obolda, M. Zhang and F. Li, *Angew. Chem.*, 2015, **127**, 7197–7201.

- 1 15 For reviews, see: (a) H. Dürr and R. Gleiter, *Angew. Chem., Int. Ed. Engl.*, 1978, **17**, 559–569; (b) R. Gleiter and G. Haberhauer, *Aromaticity and Other Conjugation Effects*, Wiley-VCH, Weinheim, 2012.
- 5 16 For butadiene radical cations see, for example: (a) C. S. Q. Lew, J. R. Brisson and L. J. Johnston, *J. Org. Chem.*, 1997, **62**, 4047; (b) J. Oxgaard and O. Wiest, *J. Phys. Chem. A*, 2001, **105**, 8236.
- 17 A Cambridge Structural Database (CSD) search with the *ConQuest* program resulted in no match (accessed February 2, 2016).
- 18 S. K. Mandal, M. E. Itkis, X. Chi, S. Samanta, D. Lidsky, R. W. Reed, R. T. Oakley, F. S. Tham and R. C. Haddon, *J. Am. Chem. Soc.*, 2005, **127**, 8185–8196.
- 19 L. E. Longobardi, L. Liu, S. Grimme and D. W. Stephan, *J. Am. Chem. Soc.*, 2016, **138**, 2500–2503.
- 10
- 15
- 20
- 25
- 30
- 35
- 40
- 45
- 50
- 55