

Asymptotic safety guaranteed

Article (Published Version)

Litim, Daniel F and Sannino, Francesco (2014) Asymptotic safety guaranteed. *Journal of High Energy Physics*, 2014 (12). p. 178. ISSN 1029-8479

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/63620/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Asymptotic safety guaranteed

Daniel F. Litim^a and Francesco Sannino^b

^a*Department of Physics and Astronomy, University of Sussex,
Falmer Campus, Brighton, BN1 9QH, U.K.*

^b*CP³-Origins & the Danish Institute for Advanced Study Danish IAS, Univ. of Southern Denmark,
Campusvej 55, DK-5230 Odense, Denmark*

E-mail: d.litim@sussex.ac.uk, sannino@cp3-origins.net

ABSTRACT: We study the ultraviolet behaviour of four-dimensional quantum field theories involving non-abelian gauge fields, fermions and scalars in the Veneziano limit. In a regime where asymptotic freedom is lost, we explain how the three types of fields cooperate to develop fully interacting ultraviolet fixed points, strictly controlled by perturbation theory. Extensions towards strong coupling and beyond the large- N limit are discussed.

KEYWORDS: 1/ N Expansion, Renormalization Group

ARXIV EPRINT: [1406.2337](https://arxiv.org/abs/1406.2337)

Contents

1	Introduction	2
2	Origin of interacting UV fixed points	4
2.1	Asymptotic safety	4
2.2	Gravitons	5
2.3	Fermions	5
2.4	Gluons	6
2.5	Scalars	6
3	From asymptotic freedom to asymptotic safety	7
3.1	Gauge-Yukawa theory	7
3.2	Leading order	8
3.3	Next-to-leading order	8
3.4	Next-to-next-to-leading order	12
3.5	UV scaling and Landau pole	15
3.6	UV critical surface	16
3.7	Phase diagram	17
3.8	Mass terms and anomalous dimensions	19
4	Consistency	20
4.1	Stability	20
4.2	Weyl consistency	20
4.3	Universality	21
4.4	Operator ordering	21
4.5	Gap	22
4.6	Unitarity	23
4.7	Triviality	23
5	Towards asymptotic safety at strong coupling	24
5.1	Beyond the Veneziano limit	24
5.2	Infinite order perturbation theory	24
5.3	Finite order perturbation theory	25
6	Conclusion	27

1 Introduction

It is widely acknowledged that ultraviolet (UV) fixed points are central for quantum field theories to be fundamental and predictive up to highest energies [1, 2]. A well-known example is asymptotic freedom of quantum chromodynamics where the UV fixed point is non-interacting [3, 4]. In turn, neither the U(1) nor the scalar sector of the standard model are asymptotically free. This is known as the triviality problem, which limits the predictivity to a scale of maximal UV extension [5]. High-energy fixed points may also be interacting, a scenario referred to as asymptotic safety [6]. It is then tempting to think that theories which are not asymptotically free, or not even renormalisable by power-counting, may well turn out to be fundamental in their own right, provided they develop an interacting UV fixed point [7]. In recent years, asymptotic safety has become a popular scenario to address quantum aspects of gravity [7–13]. In a similar vein, UV conformal extensions of the standard model with and without gravity have received some attention in view of interacting fixed points [14–29] and scale invariance in particle physics and cosmology [30–53].

The most notable difference between asymptotically free and asymptotically safe theories relates to residual interactions at high energies. Canonical power counting becomes modified and the relevant or marginal invariants which dominate high energy physics are no longer known a priori. Couplings may become large and small expansion parameters are often not available. Establishing or refuting asymptotic safety in a reliable manner then becomes a challenging non-perturbative task [54]. A few rigorous results for asymptotically safe UV fixed points have been obtained for certain power-counting non-renormalisable models by taking the space-time dimensionality as a continuous parameter [6, 16, 55–59] in the spirit of the ϵ -expansion [60], or by using large- N techniques [19, 61–68]. Asymptotic safety then arises in the vicinity of the Gaussian fixed point where perturbation theory is applicable. The success of well-controlled model studies provides valuable starting points to search for asymptotic safety at strong coupling.

In this paper, we are interested in the UV behaviour of interacting gauge fields, fermions and scalars in four dimensions. In the regime where asymptotic freedom is lost, we ask the question whether the theory is able, dynamically, to develop an interacting UV fixed point. Our main tool to answer this question is a suitably chosen large- N limit [69] (where N refers to the number of fields), whereby the theory is brought under strict perturbative control. Banks and Zaks [70] have used a similar idea to investigate the presence of interacting infrared (IR) fixed points in gauge theories with fermionic matter. Here, we will discover that all three types of fields are required for an asymptotically safe UV fixed point to emerge. Given that exactly solvable models in four dimensions are hard to come by, our findings are a useful starting point to construct UV safe models of particle physics.

We illustrate the main outcome for gauge-matter theories in the absence of asymptotic freedom in terms of a running non-Abelian coupling (α_g) and a running Yukawa coupling (α_y). Figure 1 shows the phase diagram close to the Gaussian fixed point to leading order in perturbation theory. Renormalization group trajectories are directed towards the IR. We observe that both the Yukawa and the gauge coupling behave QED-like close to the Gaussian. Consequently, not a single trajectory emanates from the Gaussian, meaning




Figure 1. The phase diagram of certain 4D gauge-Yukawa theories in the absence of asymptotic freedom and supersymmetry, in the vicinity of an asymptotically safe fixed point (UV). The renormalization group flow for the gauge coupling α_g and the Yukawa coupling α_y is pointing towards the IR. The two renormalization group trajectories emanating out of the asymptotically safe fixed point define UV finite quantum field theories which at low energies correspond to a weakly (G) and a strongly coupled theory. Other trajectories exist as well, but they do not lead to UV finite theories (within perturbation theory). Parameter choices and further details are given in section 3.7.

that it is an IR fixed point. The main novelty is the occurrence of an interacting fixed point induced by fluctuations of the gauge, fermion, and scalar fields. The fixed point is located close to the Gaussian and controlled by perturbation theory. The UV nature of the fixed point is evidenced by the (two) UV finite renormalization group trajectories running out of it. They lead to sensible theories at all scales: in the UV, they are finite due to the interacting fixed point. In the IR, they correspond to weakly coupled theories with Gaussian scaling, and to strongly coupled theories with confinement and chiral symmetry breaking or conformal behaviour, respectively. We also find UV unstable trajectories which do not emanate from the UV fixed point. They equally approach sensible IR theories, yet their UV predictivity is limited (at least in perturbation theory) by a scale of maximal UV extension. In this sense, asymptotic safety guarantees the existence of UV finite matter-gauge theories even in the absence of asymptotic freedom or supersymmetry.

In the rest of the paper we provide the details of our study. We recall the perturbative origin of interacting UV fixed points and asymptotic safety for sample theories of self-interacting gravitons, fermions, gluons, and scalars (section 2). We then explain why and how asymptotic safety can arise for certain gauge-Yukawa theories in strictly four dimensions. Fully interacting UV fixed points are found and analysed together with their universal scaling exponents, the UV critical surface, and the phase diagram of figure 1 (sec-

tion 3). Aspects such as stability, Weyl consistency, unitarity, and triviality are discussed (section 4), together with further directions for asymptotic safety within perturbation theory and beyond (section 5). We close with some conclusions (section 6).

2 Origin of interacting UV fixed points

In this section, we recall the perturbative origin of asymptotic safety for certain quantum field theories. We discuss key examples and introduce some notation.

2.1 Asymptotic safety

Asymptotic safety is the scenario which generalises the notion of a free, Gaussian, ultraviolet fixed point to an interacting, non-Gaussian one. An asymptotically safe UV fixed point then acts as an anchor for the renormalisation group evolution of couplings, allowing them to approach the high-energy limit along well-defined RG trajectories without encountering divergences such as Landau poles.

Within perturbation theory, the origin for asymptotic safety is best illustrated in terms of a running dimensionless coupling $\alpha = \alpha(t)$ of a hypothetical theory (to be specified below) with its renormalisation group (RG) β -function given by

$$\partial_t \alpha = A \alpha - B \alpha^2. \tag{2.1}$$

Here, $t = \ln(\mu/\Lambda)$ denotes the logarithmic RG ‘time’, μ the RG momentum scale and Λ a characteristic reference scale of the theory. A and B are numbers. We assume that (2.1) arises from a perturbative expansion of the full β -function ($\beta \equiv \partial_t \alpha$), with α reasonably small for perturbation theory to be applicable. The linear term relates to the tree level contribution, reflecting that the underlying coupling is dimensionful with mass dimension $-A$. The quadratic term stands for the one loop contribution. Evidently, the flow displays two types of fixed points, a trivial one at $\alpha_* = 0$, and a non-trivial one at

$$\alpha_* = A/B. \tag{2.2}$$

In the spirit of perturbation theory, the non-trivial fixed point (2.2) is accessible in the domain of validity of the RG flow (2.1) as long as $\alpha_* \ll 1$. This can be achieved in two manners, either by having $A \ll 1$ at fixed B , or by making $1/B \ll 1$ at fixed A . (Below, we discuss examples where both of these options are realised.) Integrating (2.1) in the vicinity of the fixed point, one finds that small deviations from it scale as

$$\delta\alpha = (\alpha - \alpha_*) \propto \left(\frac{\mu}{\Lambda_c}\right)^\vartheta, \tag{2.3}$$

thereby relating the characteristic energy scale Λ_c of the theory to the deviation from the fixed point $\delta\alpha$ at the RG scale μ and a universal number ϑ . The role of Λ_c here is similar to that of Λ_{QCD} in QCD as it describes the cross-over between two different scaling regimes of the theory. The universal scaling index ϑ arises as the eigenvalue of the one-dimensional ‘stability matrix’

$$\vartheta = \left. \frac{\partial\beta}{\partial\alpha} \right|_*. \tag{2.4}$$

It is given by $\vartheta = -A$ at the non-trivial fixed point (2.2), and by $\vartheta = A$ at the Gaussian fixed point.¹ We have that $\vartheta < 0$ at the non-trivial fixed point provided $A > 0$. Consequently, small deviations from the fixed point (2.3) decrease with increasing RG momentum scale meaning that (2.2) is an UV fixed point. If, additionally, $B > 0$, the fixed point obeys $\alpha_* > 0$. Consequently, in this case the Gaussian fixed point of the model (2.1) becomes an infrared (IR) fixed point. Unlike for asymptotically free theories where the RG running close to a trival UV fixed point is logarithmically slow, the RG running close to a non-trivial fixed point is power-like, and thus much faster. We notice that for $A = 0$ the model (2.1) displays a doubly-degenerate Gaussian fixed point. In this light, as soon as the canonical mass dimension of the underlying coupling becomes negative, $A > 0$, such as in theories which are power-counting non-renormalisable, the degeneracy of the perturbative β -function is lifted leading to a pair of non-degenerate fixed points. Provided that α is the sole relevant coupling of the model under consideration, the existence of an interacting UV fixed point can be used to define the theory fundamentally. This are the bare bones of asymptotic safety [1, 2, 6].

2.2 Gravitons

We now recall specific examples for asymptotically safe quantum field theories where the mechanism just described is at work. We start with Einstein gravity with action $(16\pi G_N)^{-1} \int \sqrt{\det \bar{g}} R$ in D dimensions. Newton's coupling G_N has canonical mass dimension $[G_N] = 2 - D$ and the theory is power-counting non-renormalisable above its critical dimension $D_c = 2$. In units of the RG scale, the dimensionless gravitational coupling of the model reads

$$\alpha = G_N(\mu)\mu^{D-2}. \tag{2.5}$$

In $D = D_c + \epsilon$ dimensions, one finds the RG flow (2.1) with $A = \epsilon \ll 1$, $B = 50/3$, and an UV fixed point (2.2) in the perturbative regime by analytical continuation of space-time dimensionality [6, 55, 56, 71]. Results have been extended to two-loop order, also including a cosmological constant [72]. It is the sign of $B > 0$ which enables an interacting UV fixed point for gravity in the perturbative domain. Much of the recent motivation to revisit asymptotic safety for gravity in four dimensions derives from this result close to two dimensions [7–13]. Advanced non-perturbative studies predict a gravitational fixed point (2.5) of order unity [54], see also [73–79].

2.3 Fermions

Next we consider a purely fermionic theory of N_F selfcoupled massless Dirac fermions with, exemplarily, Gross-Neveu-type selfinteraction $\frac{1}{2}g_{\text{GN}}(\bar{\psi}\psi)^2$, [80]. The quartic fermionic self-coupling g_{GN} has canonical mass dimension $[g_{\text{GN}}] = 2 - D$ and the model is perturbatively non-renormalisable above its critical dimension $D_c = 2$. The dimensionless coupling reads

$$\alpha = \frac{g_{\text{GN}}(\mu)}{2\pi N_F} \mu^{2-D}. \tag{2.6}$$

¹The leading exponent ϑ is related to the exponent ν in the statistical physics literature as $\nu = -1/\vartheta$.

Close to two dimensions the β -function (2.1) for (2.6) can be computed within the ϵ -expansion by setting $D = D_c + \epsilon$. The coefficient A , given by minus the canonical mass dimension, becomes $A = \epsilon \ll 1$, while the coefficient $B > 0$, to leading order in ϵ , is of order one and given by the 1-loop coefficient in the two-dimensional theory. Hence, the model has a reliable UV fixed point (2.2) in the perturbative regime. Its renormalisability has been established more rigorously in [58, 59] with the help of the non-perturbative renormalisation group. Similarly, in the large- N_F limit and at fixed dimension $D = 3$, one finds that $A \propto 1/N_F \ll 1$ while the coefficient $B > 0$ remains of order unity, leading to the same conclusion [64]. UV fixed points of four-fermion theories have been studied non-perturbatively in the continuum and on the lattice, e.g. [81–83]. For a simple example, see [84].

2.4 Gluons

Next, we consider pure $SU(N_C)$ Yang-Mills theory with N_C colors in $D = 4 + \epsilon$ dimensions, with action $\sim 1/(g_{\text{YM}}^2) \int \text{Tr} F_{\mu\nu} F^{\mu\nu}$. The canonical mass dimension of the coupling reads $[g_{\text{YM}}^2] = 4 - D$ and the theory is perturbatively non-renormalisable above its critical dimension $D_c = 4$. Introducing the running dimensionless strong coupling as

$$\alpha = \frac{g_{\text{YM}}^2 N_C}{(4\pi)^2} \mu^{D-4}, \quad (2.7)$$

one may derive its β -function (2.1) to leading order in the ϵ -expansion, $\epsilon = D - D_c$. Again, one finds that $A = \epsilon \ll 1$ and $B > 0$ of order unity, thus leading to an UV fixed point (2.2) [57]. The expansion has been extended up to fourth order in ϵ [16] suggesting that this fixed point persists in $D = 5$, in accord with a prediction using functional renormalisation [85].

2.5 Scalars

Finally we turn to self-interacting scalar fields. For scalar field theory with linearly realised $O(N)$ symmetry, the dimensionless version of its quartic self-coupling is given by

$$\alpha = \frac{\lambda N}{(4\pi)^2} \mu^{D-4}. \quad (2.8)$$

The critical dimension of these models is $D_c = 4$. Within the ϵ -expansion away from the critical dimension, using $D = 4 - \epsilon$, the one loop β -function is of the form (2.1) with $A = -\epsilon < 0$ and $B < 0$. Therefore, the fixed point (2.2) is physical below the critical dimension, and it is an infrared one, i.e. the seminal Wilson-Fisher fixed point. For the physically relevant dimension $D = 3$, its existence has been confirmed even beyond perturbation theory [86, 87]. In this light, the Wilson-Fisher fixed point can be viewed as the infrared analogue of asymptotic safety. The search for interacting UV fixed points in $D > 4$ dimensions within perturbation theory has seen renewed interest recently [88]. Functional renormalisation suggests the absence of global fixed points [89], in accord with the qualitative picture obtained here.

In non-linear σ -models, i.e. scalar theories with non-linearly realised internal symmetry, the critical dimension is reduced to $D_c = 2$. The relevant coupling then displays an

UV fixed point within perturbation theory [90, 91]. Fixed points of non-linear σ -models have also been investigated for their similarity with gravity [92]. Lattice results for an interacting UV fixed point in $D = 3$ in accord with functional renormalisation have recently been reported in [93].

3 From asymptotic freedom to asymptotic safety

In this section, we explain the perturbative origin for asymptotic safety in a class of gauge-Yukawa theories in strictly four dimensions.

3.1 Gauge-Yukawa theory

We consider a theory with $SU(N_C)$ gauge fields A_μ^a and field strength $F_{\mu\nu}^a$ ($a = 1, \dots, N_C^2 - 1$), N_F flavors of fermions Q_i ($i = 1, \dots, N_F$) in the fundamental representation, and a $N_F \times N_F$ complex matrix scalar field H uncharged under the gauge group. The fundamental action is taken to be the sum of the Yang-Mills action, the fermion kinetic terms, the Yukawa coupling, and the scalar kinetic and self-interaction Lagrangean $L = L_{YM} + L_F + L_Y + L_H + L_U + L_V$, with

$$L_{YM} = -\frac{1}{2} \text{Tr} F^{\mu\nu} F_{\mu\nu} \tag{3.1}$$

$$L_F = \text{Tr} (\bar{Q} i \not{D} Q) \tag{3.2}$$

$$L_Y = y \text{Tr} (\bar{Q}_L H Q_R + \bar{Q}_R H^\dagger Q_L) \tag{3.3}$$

$$L_H = \text{Tr} (\partial_\mu H^\dagger \partial^\mu H) \tag{3.4}$$

$$L_U = -u \text{Tr} (H^\dagger H)^2 \tag{3.5}$$

$$L_V = -v (\text{Tr} H^\dagger H)^2. \tag{3.6}$$

Tr is the trace over both color and flavor indices, and the decomposition $Q = Q_L + Q_R$ with $Q_{L/R} = \frac{1}{2}(1 \pm \gamma_5)Q$ is understood. This theory has been investigated for its interesting properties in [65–68]. We will motivate it for our purposes while we progress. In four dimensions, the model has four classically marginal coupling constants given by the gauge coupling, the Yukawa coupling y , the quartic scalar couplings u and the ‘double-trace’ scalar coupling v , which we write as

$$\alpha_g = \frac{g^2 N_C}{(4\pi)^2}, \quad \alpha_y = \frac{y^2 N_C}{(4\pi)^2}, \quad \alpha_h = \frac{u N_F}{(4\pi)^2}, \quad \alpha_v = \frac{v N_F^2}{(4\pi)^2}. \tag{3.7}$$

We have normalized the couplings with the appropriate powers of N_C and N_F preparing for the Veneziano limit to be considered below. Notice the additional power of N_F in the definition of the scalar double-trace coupling, meaning that v/u becomes $\alpha_v/(\alpha_h N_F)$. We also use the shorthand notation $\beta_i \equiv \partial_t \alpha_i$ with $i = (g, y, h, v)$ to indicate the β -functions for the couplings (3.7). To obtain explicit expressions for these, we use the results [94–96].

3.2 Leading order

We begin our reasoning with the RG flow for the gauge coupling to one-loop order using the $SU(N_C)$ Yang-Mills Lagrangean (3.1) coupled to N_F fermions (3.2) in the fundamental representation,

$$\beta_g = \partial_t \alpha_g = -B \alpha_g^2. \tag{3.8}$$

Note that a linear term $A\alpha_g$ is absent, unlike in (2.1), as we strictly stick to four dimensions. To this order the gauge β -function (3.8) displays a doubly-degenerated fixed point at

$$\alpha_g^* = 0 \tag{3.9}$$

which is an UV fixed point for positive B . Provided that the coefficient B is numerically very small, $|B| \ll 1$, however, we also have that $\partial_t \alpha_g \ll 1$ close to the Gaussian fixed point, indicating that the theory might develop a non-trivial fixed point with

$$0 < \alpha_g^* \ll 1 \tag{3.10}$$

once higher loop terms are included. For $B > 0$, (3.9) corresponds to asymptotic freedom, in which case (3.10) would then correspond to a conformal infrared fixed point.² In turn, for $B < 0$ asymptotic freedom is lost, the Gaussian fixed point becomes an IR fixed point, and the theory may become asymptotically safe perturbatively at (3.10). In our setup, the one-loop coefficient B depends on both N_C and N_F . Explicitly, $B = -\frac{4}{3}\epsilon$, where

$$\epsilon = \frac{N_F}{N_C} - \frac{11}{2}. \tag{3.11}$$

For $\epsilon > 0$, asymptotic freedom of the gauge sector is lost. The prerequisite for an asymptotically safe fixed point within the perturbative regime thus translates into

$$0 \leq \epsilon \ll 1. \tag{3.12}$$

Consequently, to achieve asymptotic safety in the gauge sector in a controlled perturbative manner, we must perform a Veneziano limit by sending both N_C and N_F to infinity, but keeping their ratio N_F/N_C fixed [69]. The parameter (3.11) thereby becomes continuous and can take any real value including (3.12). In most of the paper, the parameter ϵ will be our primary perturbative control parameter in the regime (3.12), except in section 5 where we also discuss the regime where ϵ becomes large.

3.3 Next-to-leading order

We now must check whether this scenario can be realized upon the inclusion of higher loop corrections. At the next-to-leading (NLO) order in perturbation theory, which is two-loop, the RG flow for the gauge coupling takes the form

$$\partial_t \alpha_g = -B \alpha_g^2 + C \alpha_g^3. \tag{3.13}$$

²There is a vast body of work dealing with the availability of this Banks-Zaks type IR fixed point [70, 97] in the continuum and on the lattice (for recent overviews see [98, 99] and references therein).




Figure 2. The coordinates of the UV fixed point as a function of ϵ at NLO (dashed) and NNLO (full and short-dashed lines). Gauge, Yukawa and scalar couplings are additionally shown in red, magenta and blue, respectively. NNLO corrections lead to a mild enhancement of the gauge and Yukawa couplings over their NLO values. The scalar and Yukawa fixed point couplings are nearly degenerate at NNLO.

As such, the gauge β -function (3.13) may display three fixed points, a doubly-degenerated one at $\alpha_g^* = 0$, and a non-trivial one at

$$\alpha_g^* = B/C. \tag{3.14}$$

The non-trivial fixed point is perturbative as long as $0 \leq \alpha_g^* \ll 1$ along with all the other possible couplings in the theory. In practice, this follows provided that $|B| \ll 1$ and C of order unity, and $B/C > 0$. For this fixed point to be an asymptotically safe one we must have $B < 0$ and $C < 0$. However, in the absence of Yukawa interactions one finds $C = 25$ [97] to leading order in ϵ . Consequently, the would-be fixed point (3.14) resides in the unphysical domain $\alpha_g^* < 0$ where the theory is sick non-perturbatively, see e.g. [100].

This conclusion changes as soon as Yukawa interactions are taken into consideration. The gauge β -function depends on the Yukawa coupling starting from the two-loop order. For this reason, to progress, we must first evaluate the impact of non-trivial Yukawa couplings. At the same time, since the fixed point for the gauge coupling depends on the Yukawa coupling, we must retain its RG flow to its first non-trivial order, which is one-loop. Having introduced Yukawa couplings means that we also have dynamical scalars, (3.3). The simplest choice here is to assume that the scalars are uncharged under the gauge group, whence (3.4). Then neither the gauge nor the Yukawa RG flows depend on the scalar couplings at this order and we can neglect their contribution for now. This ordering of perturbation theory for the different couplings is also favored by considerations related to Weyl consistency conditions (see section 4.2). Hence, following this reasoning and in the

presence of the Yukawa term (3.3), we end up with

$$\begin{aligned}\beta_g &= \alpha_g^2 \left\{ \frac{4}{3}\epsilon + \left(25 + \frac{26}{3}\epsilon\right) \alpha_g - 2 \left(\frac{11}{2} + \epsilon\right)^2 \alpha_y \right\}, \\ \beta_y &= \alpha_y \left\{ (13 + 2\epsilon) \alpha_y - 6 \alpha_g \right\}.\end{aligned}\tag{3.15}$$

The coupled system (3.15) admits three types of fixed points within perturbation theory (3.12). The system still displays a Gaussian fixed point

$$(\alpha_g^*, \alpha_h^*) = (0, 0)\tag{3.16}$$

irrespective of the sign of ϵ . For $\epsilon > 0$, neither the gauge coupling nor the Yukawa coupling are asymptotically free at this fixed point. Ultimately, this is related to the sign of either β -function being positive arbitrarily close to (3.16), making it an IR fixed point. A second, non-trivial fixed point is found as well, $(\alpha_g^*, \alpha_h^*) \neq (0, 0)$ which is an UV fixed point with coordinates

$$\begin{aligned}\alpha_g^* &= \frac{26\epsilon + 4\epsilon^2}{57 - 46\epsilon - 8\epsilon^2} = \frac{26}{57}\epsilon + \frac{1424}{3249}\epsilon^2 + \frac{77360}{185193}\epsilon^3 + \mathcal{O}(\epsilon^4) \\ \alpha_y^* &= \frac{12\epsilon}{57 - 46\epsilon - 8\epsilon^2} = \frac{4}{19}\epsilon + \frac{184}{1083}\epsilon^2 + \frac{10288}{61731}\epsilon^3 + \mathcal{O}(\epsilon^4).\end{aligned}\tag{3.17}$$

Numerically, the series (3.17) reads

$$\begin{aligned}\alpha_g^* &= 0.4561 \epsilon + 0.4383 \epsilon^2 + 0.4177 \epsilon^3 + \mathcal{O}(\epsilon^4) \\ \alpha_y^* &= 0.2105 \epsilon + 0.1699 \epsilon^2 + 0.1667 \epsilon^3 + \mathcal{O}(\epsilon^4)\end{aligned}\tag{3.18}$$

This UV fixed point is physically acceptable in the sense that $(\alpha_g^*, \alpha_y^*) > (0, 0)$ for $\epsilon > 0$. It arises because the gauge and Yukawa couplings contribute with opposite signs to β_g at the two-loop level, allowing for an asymptotically safe fixed point in the physical domain.

As an aside, we also notice the existence of a second interacting fixed point within perturbation theory located at

$$(\alpha_g^*, \alpha_y^*) = \left(-\frac{4\epsilon}{75 + 26\epsilon}, 0 \right).\tag{3.19}$$

For $\epsilon > 0$ this fixed point cannot be reached by any finite RG flow starting from the domain where $\alpha_g > 0$ and is unphysical. For $\epsilon < 0$, (3.19) takes the role of an interacting infrared fixed point *à la* Caswell, Banks and Zaks [70, 97]. Interacting IR fixed points play an important role in extensions of the standard model with a strongly interacting gauge sector and models with a composite Higgs (see [98, 99] and references therein). The IR fixed point arises even in the absence of scalar fields, whereas the UV fixed point necessitates scalar matter with non-vanishing Yukawa interactions. We conclude that the IR fixed point (3.19) is profoundly different from the UV fixed point (3.17).

Returning to our main line of reasoning, we linearize the RG flow in the vicinity of its UV fixed point (3.17),

$$\beta_i = \sum_j M_{ij} (\alpha_j - \alpha_j^*) + \text{subleading}\tag{3.20}$$




Figure 3. Components of the eigenvectors (3.23) and (3.37) (absolute values) corresponding to the relevant eigenvalue ϑ_1 at the UV fixed point at NLO (dashed) and NNLO (full lines), respectively, as functions of ϵ . From top to bottom, the gauge, Yukawa and scalar, and double-trace scalar components are shown. Notice that the relevant eigendirection is largely dominated by the gauge coupling. Each component varies only mildly with ϵ .

where $i = (g, y)$ and $M_{ij} = \partial\beta_i/\partial\alpha_j|_*$ denotes the stability matrix. The eigenvalues of M are universal numbers and characterise the scaling of couplings in the vicinity of the fixed point. They can be found analytically. The first few orders in (3.11) are

$$\begin{aligned} \vartheta_1 &= -\frac{104}{171}\epsilon^2 + \frac{2296}{3249}\epsilon^3 + \frac{1387768}{1666737}\epsilon^4 + \mathcal{O}(\epsilon^4) \\ \vartheta_2 &= \frac{52}{19}\epsilon + \frac{9140}{1083}\epsilon^2 + \frac{2518432}{185193}\epsilon^3 + \mathcal{O}(\epsilon^4). \end{aligned} \tag{3.21}$$

Numerically, the eigenvalues (3.21) read

$$\begin{aligned} \vartheta_1 &= -0.608\epsilon^2 + 0.707\epsilon^3 - 0.833\epsilon^4 + \mathcal{O}(\epsilon^5) \\ \vartheta_2 &= 2.737\epsilon + 8.44\epsilon^2 + 13.599\epsilon^3 + \mathcal{O}(\epsilon^4) \end{aligned} \tag{3.22}$$

A few comments are in order. Firstly, the gauge-Yukawa system at NLO has developed a relevant and an irrelevant eigendirection with eigenvalues $\vartheta_1 < 0$ and $\vartheta_2 > 0$, respectively.³

Secondly, the relevant eigenvalue ϑ_1 is found to be of order ϵ^2 , whereas the irrelevant one is of order ϵ and thus parametrically larger. This is not a coincidence, and its origin can be understood as follows: all couplings settle at values of order ϵ at the fixed point. Hence, $\beta_g \sim \epsilon^3$ and $\beta_y \sim \epsilon^2$ in the vicinity of the fixed point. The relevant eigenvalue originates primarily from the gauge sector, because asymptotic freedom is destabilised due to (3.12). Consequently, at the fixed point, the relevant eigenvalue must scale as $\vartheta_1 \sim \partial\beta_g/\partial\epsilon|_* \sim \epsilon^2$. Conversely, the irrelevant eigenvalue scales as $\vartheta_2 \sim \partial\beta_y/\partial\epsilon|_* \sim \epsilon$. We conclude that the parametric dependence $\sim \epsilon^2$ of the relevant eigenvalue arises because the fixed point in the

³Strictly speaking, there are two further marginal eigenvalues $\vartheta_{3,4} = 0$ related to the scalar selfinteractions which we have taken as classical.

gauge sector stems from a cancellation at two-loop level. Conversely, the behaviour $\sim \epsilon$ of the irrelevant eigenvalue stems from cancellations at the one-loop level. This feature is a direct consequence of the vanishing mass dimension of the couplings. Asymptotic safety then follows as a pure quantum effect rather than through the cancelation of tree-level and one-loop terms.

Finally, introducing a basis in coupling parameter space as $\mathbf{a} = (\alpha_g, \alpha_y)^T$, we denote the relevant eigendirection at the UV fixed point as

$$\mathbf{e}_1 = (e_g, e_y)^T. \tag{3.23}$$

The absolute values of its entries are shown in figure 3 (dashed lines). We find that the relevant eigenvalue is largely dominated by the gauge coupling for all ϵ . Furthermore, the relevant eigendirection is largely independent of ϵ , up to $\epsilon < 0.7$ where couplings become of order one. The domain of validity is further discussed in section 5.1 below.

3.4 Next-to-next-to-leading order

We now move on to the next-to-next-to-leading order (NNLO) in the perturbative expansion where the scalar sector is no longer treated as exactly marginal. Identifying a combined UV fixed point in all couplings becomes a consistency check for asymptotic safety in the full theory. In practice, this amounts to adding the quartic selfinteraction terms (3.5) and (3.6). The one-loop running of the quartic couplings given by

$$\beta_h = -(11 + 2\epsilon)\alpha_y^2 + 4\alpha_h(\alpha_y + 2\alpha_h), \tag{3.24}$$

$$\beta_v = 12\alpha_h^2 + 4\alpha_v(\alpha_v + 4\alpha_h + \alpha_y). \tag{3.25}$$

For consistency, we also include the two-loop corrections to the running of the Yukawa coupling and the three-loop contributions to the running of the gauge coupling,

$$\begin{aligned} \Delta\beta_y^{(2)} &= \alpha_y \left\{ \frac{20\epsilon - 93}{6}\alpha_g^2 + (49 + 8\epsilon)\alpha_g\alpha_y - \left(\frac{385}{8} + \frac{23}{2}\epsilon + \frac{\epsilon^2}{2} \right)\alpha_y^2 - (44 + 8\epsilon)\alpha_y\alpha_h + 4\alpha_h^2 \right\} \\ \Delta\beta_g^{(3)} &= \alpha_g^2 \left\{ \left(\frac{701}{6} + \frac{53}{3}\epsilon - \frac{112}{27}\epsilon^2 \right)\alpha_g^2 - \frac{27}{8}(11 + 2\epsilon)^2\alpha_g\alpha_y + \frac{1}{4}(11 + 2\epsilon)^2(20 + 3\epsilon)\alpha_y^2 \right\}. \end{aligned} \tag{3.26}$$

Several comments are in order. Firstly, both scalar β functions are quadratic polynomials in the couplings and display a Gaussian fixed point. The full system at NNLO then displays a Gaussian fixed point $(\alpha_g^*, \alpha_y^*, \alpha_h^*, \alpha_v^*) = (0, 0, 0, 0)$ as it must.

Secondly, and unlike all other β -functions, the RG flow (3.25) for the double-trace coupling shows no explicit dependence on ϵ . Also, to leading order in $1/N_C$ and $1/N_F$, (3.25) stays quadratic in its coupling to all loop orders [101].

Finally, and most notably, the β -functions of the gauge, Yukawa and single-trace scalar coupling remain independent of the double-trace scalar coupling α_v . In consequence the dynamics of α_v largely decouples from the system and it acts like a spectator without influencing the build-up of the asymptotically safe UV fixed point in the gauge-Yukawa-scalar subsector. Its own RG evolution is primarily fueled by the Yukawa and the single trace coupling, and as such indirectly sensitive to the gauge-Yukawa fixed point. In turn, the scalar coupling α_h couples back into the Yukawa coupling, though not into the gauge coupling.

Since the RG flows at NNLO partly factorize, we can start by first considering the corrections to (3.17) induced by the scalar coupling α_h . This leads to UV fixed points in the gauge-Yukawa-scalar subsystem

$$(\alpha_g^*, \alpha_y^*, \alpha_h^*) \neq (0, 0, 0). \quad (3.27)$$

In the background of the gauge-Yukawa fixed point (3.17), the RG flow (3.24) for α_h admits two fixed points $\alpha_{h2}^* < 0 < \alpha_{h1}^*$, with

$$\alpha_{h1,h2}^* = (\pm\sqrt{23} - 1) \frac{\epsilon}{19} + \mathcal{O}(\epsilon^2), \quad (3.28)$$

irrespective of α_v . Inserting α_{h1}^* together with (3.17) into (3.25) we then also find two solutions for the double trace coupling, with $\alpha_{v2}^* < \alpha_{v1}^*$. These are discussed in more detail below. Conversely, inserting α_{h2}^* together with (3.17) into (3.25) does not offer a fixed point for α_v . We conclude that the fixed point $\alpha_h^* \equiv \alpha_{h1}^* > 0$ is the sole value for the coupling α_h which leads to a UV fixed point in the scalar subsystem. We return to this in section 3.5.

Using (3.15), (3.24), and (3.26), the coordinates of the gauge, Yukawa and scalar coupling are obtained analytically and can be expressed as a power series in ϵ starting as

$$\begin{aligned} \alpha_g^* &= \frac{26}{57} \epsilon + \frac{23(75245 - 13068\sqrt{23})}{370386} \epsilon^2 + \mathcal{O}(\epsilon^3) \\ \alpha_y^* &= \frac{4}{19} \epsilon + \left(\frac{43549}{20577} - \frac{2300\sqrt{23}}{6859} \right) \epsilon^2 + \mathcal{O}(\epsilon^3) \\ \alpha_h^* &= \frac{\sqrt{23} - 1}{19} \epsilon + \frac{1168991 - 202249\sqrt{23}}{82308\sqrt{23}} \epsilon^2 + \mathcal{O}(\epsilon^3). \end{aligned} \quad (3.29)$$

Numerically, the first few orders in the ϵ -expansion read

$$\begin{aligned} \alpha_g^* &= 0.4561 \epsilon + 0.7808 \epsilon^2 + 3.112 \epsilon^3 + \mathcal{O}(\epsilon^4) \\ \alpha_y^* &= 0.2105 \epsilon + 0.5082 \epsilon^2 + 2.100 \epsilon^3 + \mathcal{O}(\epsilon^4) \\ \alpha_h^* &= 0.1998 \epsilon + 0.5042 \epsilon^2 + 2.045 \epsilon^3 + \mathcal{O}(\epsilon^4). \end{aligned} \quad (3.30)$$

The addition of the scalar selfcouplings has led to a physical fixed point $\alpha_h^* > 0$ of order ϵ . NNLO corrections to α_g^* and α_y^* arise only starting at order ϵ^2 without altering the NLO fixed point (3.17). Performing the expansion (3.30) to high orders in ϵ one finds its radius of convergence as

$$\epsilon \leq \epsilon_{\max} = 0.117 \dots \quad (3.31)$$

At ϵ_{\max} in (3.31), the NNLO equations display a bifurcation and the UV fixed point ceases to exist through a merger with a non-perturbative IR fixed point, and the relevant eigenvalue disappears at ϵ_{\max} . The merger at ϵ_{\max} indicates that our working assumption (3.12) should be superseded by $0 < \epsilon \ll \epsilon_{\max}$.

Since α_v does not contribute to the RG flow of the subsystem $(\alpha_g, \alpha_y, \alpha_h)$, the computation of scaling exponents equally factorizes. Linearizing the RG flow in the vicinity




Figure 4. Universal eigenvalues $\vartheta_1 < 0 < \vartheta_2 < \vartheta_3$ at the UV fixed point at NLO (dashed lines) and NNLO (full lines) as function of ϵ (3.11). For all technical purposes, the eigenvalues are dominated by the NLO values except for the close vicinity of $\epsilon \approx \epsilon_{\max}$.

of the fixed point, we have (3.20) where now $i = (g, y, h)$. The eigenvalues ϑ_n are found analytically as a power series in ϵ , the first few orders of which are given by

$$\begin{aligned}
 \vartheta_1 &= -\frac{104}{171} \epsilon^2 + \frac{2296}{3249} \epsilon^3 + \frac{4531558295989 - 922557832416\sqrt{23}}{46931980446} \epsilon^4 + \mathcal{O}(\epsilon^5) \\
 \vartheta_2 &= \frac{52}{19} \epsilon + \frac{136601719 - 22783308\sqrt{23}}{4094823} \epsilon^2 + \mathcal{O}(\epsilon^3) \\
 \vartheta_3 &= \frac{16\sqrt{23}}{19} \epsilon + 4 \frac{217933589\sqrt{23} - 695493228}{94180929} \epsilon^2 + \mathcal{O}(\epsilon^3)
 \end{aligned}
 \tag{3.32}$$

For the relevant eigenvalue, we notice that the first two non-trivial orders have remained unchanged. Numerically, the eigenvalues read

$$\begin{aligned}
 \vartheta_1 &= -0.608 \epsilon^2 + 0.707 \epsilon^3 + 2.283 \epsilon^4 + \dots \\
 \vartheta_2 &= 2.737 \epsilon + 6.676 \epsilon^2 + \dots \\
 \vartheta_3 &= 4.039 \epsilon + 14.851 \epsilon^2 + \dots
 \end{aligned}
 \tag{3.33}$$

The cubic and quartic corrections to the relevant eigenvalue both arise with a sign opposite to the leading term, which is responsible for the smallness of ϑ_1 even for moderate values of ϵ . In turn, the irrelevant eigenvalues receive larger corrections and reach values of the order of $0.1 \div 1$ for moderate ϵ .

We now discuss the role of the double-trace scalar coupling α_v . Its fixed points are entirely induced by the fixed point in the gauge-Yukawa-scalar subsystem (3.29). Two solutions are found,

$$\alpha_{v1,v2}^* = -\frac{1}{19} \left(2\sqrt{23} \mp \sqrt{20 + 6\sqrt{23}} \right) \epsilon + \mathcal{O}(\epsilon^2)
 \tag{3.34}$$

Numerically $\alpha_{v1}^* = -0.1373 \epsilon$ and $\alpha_{v2}^* = -0.8723 \epsilon$, up to quadratic corrections in ϵ . In principle, either of these fixed points can be used in conjunction with (3.29) to define the

combined UV fixed point of the theory. We also note that

$$\alpha_{h1}^* + \alpha_{v2}^* < 0 < \alpha_{h1}^* + \alpha_{v1}^*, \tag{3.35}$$

showing that the scalar field potential is bounded (unbounded) from below for the latter (former) choice of couplings.⁴ At the fixed point α_{v1}^* , the invariant (3.6) becomes perturbatively irrelevant and adds a positive scaling exponent to the spectrum. Conversely, at the fixed point α_{v2}^* , the invariant (3.6) has become perturbatively relevant. Since the RG flow (3.25) is quadratic in the coupling α_v to all orders, the corresponding scaling exponents are equal in magnitude with opposite signs,

$$\vartheta_4 = \frac{8\epsilon}{19} \sqrt{20 + 6\sqrt{23}} + \mathcal{O}(\epsilon^2). \tag{3.36}$$

Numerically, $\vartheta_4 = 2.941\epsilon + \mathcal{O}(\epsilon^2)$. The occurrence of an additional negative eigenvalue is induced by the interacting UV fixed point (3.29).

We may introduce a basis in coupling parameter space as $\mathbf{a} = (\alpha_g, \alpha_y, \alpha_h, \alpha_v)^T$ to denote the eigendirections at the UV fixed point as \mathbf{e}_i . They obey the eigenvalue equation $M\mathbf{e}_i = \vartheta_i \mathbf{e}_i$. The normalised relevant eigendirection \mathbf{e}_1 corresponding to the UV attractive eigenvalue has the components

$$\mathbf{e}_1 = (e_g, e_y, e_h, e_v)^T \tag{3.37}$$

whose values as function of (3.11) are shown in figure 3. Clearly, even at NNLO the gauge coupling dominates the relevant eigendirection. We also find that the eigendirection corresponding to ϑ_4 is independent of the gauge, Yukawa and scalar coupling, $\mathbf{e}_4 = (0, 0, 0, 1)$.

3.5 UV scaling and Landau pole

We summarize the main picture (see also table 1). The asymptotically safe UV fixed point (3.17) in the gauge-Yukawa system bifurcates into several fixed points once the scalar fluctuations are taken into account. In addition to the universal eigenvalues of the gauge-Yukawa system ϑ_1 and ϑ_2 , the scalar sector add the eigenvalues $\pm\vartheta_3$ and $\pm\vartheta_4$. To the leading non-trivial order in ϵ , these are

$$\begin{aligned} \vartheta_1 &= -0.608\epsilon^2 + \mathcal{O}(\epsilon^3) \\ \vartheta_2 &= 2.737\epsilon + \mathcal{O}(\epsilon^2) \\ \vartheta_3 &= 4.039\epsilon + \mathcal{O}(\epsilon^2) \\ \vartheta_4 &= 2.941\epsilon + \mathcal{O}(\epsilon^2). \end{aligned} \tag{3.38}$$

Complete asymptotic safety, e.g. asymptotically safe UV fixed points in all couplings, is achieved at two UV fixed points, FP₁ and FP₂. At FP₁, both scalar invariants are perturbatively irrelevant and the eigenvalue spectrum is

$$\vartheta_1 < 0 < \vartheta_2 < \vartheta_4 < \vartheta_3. \tag{3.39}$$

⁴An inspection of radiative corrections confirms that the effective potential for the scalar fields is stable classically and quantum-mechanically [102].

fixed point	couplings				eigenvalues	
	gauge	Yukawa	scalar	double-trace	relevant	irrelevant
FP ₁	α_g^*	α_y^*	α_{h1}^*	α_{v1}^*	ϑ_1	$\vartheta_2, \vartheta_3, \vartheta_4$
FP ₂	α_g^*	α_y^*	α_{h1}^*	α_{v2}^*	$-\vartheta_4, \vartheta_1$	ϑ_2, ϑ_3
FP ₃	α_g^*	α_y^*	α_{h2}^*	Landau pole	$-\vartheta_3, \vartheta_1$	ϑ_2

Table 1. Summary of UV fixed points of the gauge-Yukawa theory and the number of (ir-)relevant eigenvalues. To leading non-trivial order, the fixed point values are given by (3.17), (3.28), (3.34) and the exponents by (3.38).

At FP₂, the partly decoupled double-trace scalar interaction term with coupling α_v becomes relevant in its own right, and the eigenvalue spectrum, instead, reads

$$-\vartheta_4 < \vartheta_1 < 0 < \vartheta_2 < \vartheta_3. \tag{3.40}$$

At either of these, the UV limit can be taken. We recall that the scalar field potential is stable (unstable) at FP₁ (FP₂), indicating that FP₁ corresponds to a physically acceptable theory at highest energies. Finally, a fixed point FP₃ exists for the gauge-Yukawa-scalar subsystem. Here, the scalar coupling α_h becomes relevant and the eigenvalue spectrum reads

$$-\vartheta_3 < \vartheta_1 < 0 < \vartheta_2. \tag{3.41}$$

Here, however, asymptotic safety is not complete. In fact, the β -function for the double-trace scalar coupling does not show a fixed point in perturbation theory as it remains strictly positive, leading to Landau poles $\alpha_v \rightarrow \pm\infty$ in the IR and in the UV. In the UV, this regime resembles the U(1) or Higgs sector of the standard model. In either case it is no longer under perturbative control. Unless strong-coupling effects resolve this singularity in the UV, this behaviour implies a limit of maximal UV extension of the model close to FP₃.

3.6 UV critical surface

The existence of relevant and irrelevant direction in the UV implies that the short-distance behaviour of the theory is described by a lower-dimensional UV critical surface. We discuss the NLO case and FP₁ at NNLO in detail. This is straightforwardly generalised to the case with two relevant eigendirections. On the critical surface in coupling constant space, we may express the RG running of the irrelevant coupling, say α_i with $(i = y, h, v)$, in terms of the relevant one, say α_g , leading to relations

$$\alpha_i = F_i(\alpha_g). \tag{3.42}$$

To see this more explicitly, we integrate the RG flow in the vicinity of the UV fixed point to find the general solution

$$\alpha_i(\mu) = \alpha_i^* + \sum_n c_n V_i^n \left(\frac{\mu}{\Lambda_c}\right)^{\vartheta_n} + \text{subleading}. \tag{3.43}$$

Here, Λ_c is a reference energy scale, ϑ_n are the eigenvalues of the stability matrix M , V^n the corresponding eigenvectors, and c_n free parameters. The eigenvectors generically mix all couplings. At NLO the eigenvalues obey $\vartheta_1 < 0 < \vartheta_2$. For all coupling α_i to reach the UV fixed point with increasing RG scale $1/\mu \rightarrow 0$ we therefore must set the free parameter $c_2 = 0$. Conversely, the parameter c_1 remains undetermined and should be viewed as a free parameter of the theory. Since this holds true for each coupling α_i , we can eliminate c_1 from (3.43) to express the irrelevant coupling in terms of the relevant one. At NLO, one finds

$$F_y(\alpha_g) = \left(\frac{6}{13} - \frac{88\epsilon}{507} \right) \alpha_g + \frac{8}{171} \epsilon^2 + \mathcal{O}(\epsilon^3) \quad (3.44)$$

to the first few orders in an expansion in ϵ . At NNLO, the hypercritical surface is extended and receives corrections due to the scalar couplings. At FP_1 , for example, one finds

$$\begin{aligned} F_y(\alpha_g) &= (0.4615 + 0.6168 \epsilon) \alpha_g - 0.1335 \epsilon^2 + \mathcal{O}(\epsilon^3) \\ F_h(\alpha_g) &= (0.4380 + 0.5675 \epsilon) \alpha_g - 0.09658 \epsilon^2 + \mathcal{O}(\epsilon^3), \\ F_v(\alpha_g) &= -(0.3009 + 0.3241 \epsilon) \alpha_g + 0.1373 \epsilon + 0.3828 \epsilon^2 + \mathcal{O}(\epsilon^3) \end{aligned} \quad (3.45)$$

to the first few orders in ϵ , and in agreement with (3.44) to order ϵ . The significance of the UV critical surface is that couplings can reach the UV fixed point only along the relevant direction, dictated by the eigenperturbations. This imposes a relation between the relevant and the irrelevant coupling, both of which scale out of the fixed point with the same scaling exponent ϑ_1 . On the critical surface and close to the fixed point, the gauge coupling evolves as

$$\alpha_g(\mu) = \alpha_g^* + \left(\alpha_g(\Lambda_c) - \alpha_g^* \right) \left(\frac{\mu}{\Lambda_c} \right)^{\vartheta_1(\epsilon)} \quad (3.46)$$

and the irrelevant couplings follow suit via (3.42) with (3.44) and (3.45), and with α_g^* and ϑ_1 given by the corresponding expressions at NLO and NNLO, respectively.

3.7 Phase diagram

In figure 1, we show the phase diagram of the asymptotically safe gauge-Yukawa theory for small couplings at NLO, where we have set $\epsilon = 0.05$. The RG trajectories are obtained from integrating (3.15) with arrows pointing towards the IR. For small couplings, one observes the Gaussian and the UV fixed points. At vanishing Yukawa (gauge) coupling, the RG equations (3.15) become infrared free in the gauge (Yukawa) coupling, corresponding to the thick red horizontal (vertical) trajectory in figure 1. This makes the Gaussian fixed point IR attractive in both couplings. The UV fixed point has a relevant and an irrelevant direction, corresponding to the two thick red trajectories one of which is flowing out of and the other into the UV fixed point. These trajectories are distinguished in that they also divide the phase diagram into four regions A, B, C and D .

The trajectory connecting the UV with the Gaussian fixed point is a separatrix, which defines the boundary between the regions A and C , and B and D . Close to the UV fixed point, its coordinates are given by the UV critical surface (3.44). The RG flow along the separatrix is given analytically to very good accuracy by (3.46) and (3.44), and for suitable




Figure 5. The gauge and Yukawa β -functions projected along the separatrix of the phase diagram given in figure 1 (NLO with $\epsilon = 0.05$), also showing the Gaussian and the UV fixed point. For better display, we have rescaled $\beta \rightarrow \beta/(\alpha_g^*)^3$, and $\alpha_g \rightarrow \alpha_g/\alpha_g^*$; see main text.

initial conditions $\alpha_g(\mu = \Lambda_c)$ on the separatrix. Note that due to the smallness of the relevant UV eigenvalue compared to the irrelevant one, $|\vartheta_1/\vartheta_2| = \frac{2}{9}\epsilon + \mathcal{O}(\epsilon^2)$, the RG flow runs very slowly along the separatrix. In turn, trajectories entering into the separatrix (with decreasing RG scale μ) run much faster, reflected by their near-perpendicular angle between these trajectories and the separatrix; see figure 1.

Trajectories in region *A* run towards the Gaussian FP in the IR, and towards strong Yukawa coupling in the UV. Trajectories in *B* run towards a strong coupling regime in the IR. In region *C*, trajectories approach the Gaussian FP in the IR limit. Finally, in region *D* trajectories approach a strongly coupled regime in the IR, outside the domain of applicability of our equations. Notice that the Gaussian fixed point is attractive in all directions. Hence, asymptotically safe trajectories emanating from the UV fixed point either run towards a weakly coupled phase controlled by the Gaussian fixed point in the deep IR, or towards a strongly coupled QCD-like phase characterised by chiral symmetry breaking and confinement. More generally, in figure 1, the boundary between weakly (*A* and *C*) and strongly (*B* and *D*) coupled phases at low energies is given by the UV irrelevant direction, i.e. the two full (red) trajectories running into the UV fixed point.

In figure 5, we show the gauge and Yukawa β -functions projected along the separatrix as functions of the gauge coupling,

$$\begin{aligned} \beta_g^{\text{sep}}(\alpha_g) &\equiv \beta_g(\alpha_g, \alpha_y = F_y(\alpha_g)) \\ \beta_y^{\text{sep}}(\alpha_g) &\equiv \beta_y(\alpha_g, \alpha_y = F_y(\alpha_g)) \end{aligned} \tag{3.47}$$

also using (3.42) with (3.44). Both of them display the Gaussian and the UV fixed point. We also recover the UV relevant eigenvalue

$$\vartheta_1 = \left. \frac{d\beta_i^{\text{sep}}}{d\alpha_g} \right|_* = \left. \frac{\partial \beta_i}{\partial \alpha_g} \right|_* + \left. \frac{\partial F_y}{\partial \alpha_g} \frac{\partial \beta_i}{\partial \alpha_y} \right|_* \tag{3.48}$$

from either of these ($i = g, y$). Close to the UV fixed point the RG running is power-like. Close to the IR fixed point, the running becomes logarithmic. Quantitatively, along the separatrix the crossover from UV scaling to IR scaling takes place once couplings are reduced to about $\sim 65\%$ of their UV fixed point values.

3.8 Mass terms and anomalous dimensions

If mass terms are present, their multiplicative renormalisation is induced through the RG flow of the gauge, Yukawa, and scalar couplings. We now discuss the scaling associated to the scalar wave function renormalization, the scalar mass, and the addition of the fermion mass operator. The former is identified with the anomalous dimension γ_H for the scalar wave function renormalization⁵ Z_H ,

$$\Delta_H = 1 + \gamma_H, \quad \gamma_H \equiv -\frac{1}{2} \frac{d \ln Z_H}{d \ln \mu}. \quad (3.49)$$

Within perturbation theory, the one and two loop contributions to (3.49) read (see e.g. [103])

$$\gamma_H^{(1)} = \alpha_y, \quad (3.50)$$

$$\gamma_H^{(2)} = -\frac{3}{2} \left(\frac{11}{2} + \epsilon \right) \alpha_y^2 + \frac{5}{2} \alpha_y \alpha_g + 2\alpha_h^2. \quad (3.51)$$

Inserting the UV fixed point FP_1 and expanding the anomalous dimension in powers of ϵ , we find

$$\gamma_H = \frac{4\epsilon}{19} + \frac{14567 - 2376\sqrt{23}}{6859} \epsilon^2 + \mathcal{O}(\epsilon^3). \quad (3.52)$$

Notice that the leading and subleading terms are both positive. The anomalous dimension for the scalar mass term m can be derived from the composite operator $\sim m^2 \text{Tr} H^\dagger H$. Introducing $\gamma_m = \frac{1}{2} d \ln m^2 / d \ln \mu$ one finds the mass anomalous dimension

$$\gamma_m^{(1)} = 2\alpha_y + 4\alpha_h + 2\alpha_v \quad (3.53)$$

to one-loop order. We find that (3.53) becomes as large as $\gamma_m^{(1)} \approx 0.09$ for $\epsilon \approx 0.1$ at FP_1 . Evidently, the loop corrections remain small compared to the tree level term leaving $m^2 = 0$ as the sole fixed point within perturbation theory. Analogously, the anomalous dimension for the fermion mass operator is defined as

$$\Delta_F = 3 - \gamma_F, \quad \gamma_F \equiv \frac{d \ln M}{d \ln \mu} \quad (3.54)$$

where M stands for the fermion mass. Within perturbation theory, the one and two loop contributions read

$$\gamma_F^{(1)} = 3\alpha_g - \alpha_y \left(\frac{11}{2} + \epsilon \right), \quad (3.55)$$

$$\gamma_F^{(2)} = (44 + 8\epsilon) \alpha_g \alpha_y + \left(\frac{31}{4} - \frac{5}{3}\epsilon \right) \alpha_g^2 + \frac{1}{4} \left(\frac{11}{2} + \epsilon \right) \left(\frac{23}{2} + \epsilon \right) \alpha_y^2. \quad (3.56)$$

⁵The bare and renormalized fields here are related via $H_B = Z_H^{\frac{1}{2}} H$. Also the wave function definition in [95] is the inverse of the one here.

Inserting the UV fixed point FP_1 and expanding in ϵ we find

$$\gamma_F = \frac{4\epsilon}{19} + \frac{4048\sqrt{23} - 59711}{6859} \epsilon^2 + \mathcal{O}(\epsilon^3). \tag{3.57}$$

The leading and subleading terms are both positive. Interestingly, to one-loop order, the scalar anomalous dimension and the fermion mass anomalous dimension coincide in magnitude. The quantum corrections are bounded, $|\gamma_H^{(1)}| < 1/40$. We stress that the leading order results are entirely fixed by the NLO fixed point (3.17), and insensitive to the details of the scalar sector. The latter only enter starting at order ϵ^2 .

4 Consistency

In this section, we discuss aspects of consistency and the validity of results.

4.1 Stability

In the LO, NLO and NNLO approximations, we have retained the β -functions of the gauge, Yukawa and quartic couplings at different loop levels within perturbation theory. As we have argued, the ordering as shown in table 2 is dictated by the underlying dynamics towards asymptotic safety, centrally controlled by the gauge coupling.

The selfconsistency of our reasoning is confirmed a posteriori by the stability of the result. Firstly, the leading coefficients in ϵ of the NLO fixed point α_g^* and α_y^* remain numerically unchanged at NNLO, see (3.17) and (3.29). We therefore expect that all coefficients up to ϵ^2 of α_g^* and α_y^* and the ϵ coefficient of α_h^* and α_v^* in (3.29), (3.28) and (3.34) remain unchanged beyond NNLO. Secondly, the stability also extends to the universal eigenvalues. Interestingly, here, the first two non-trivial coefficients (up to order ϵ^3) for the relevant eigenvalue ϑ_1 at NLO agree with the NNLO coefficients, see (3.21) and (3.32). For the leading irrelevant eigenvalue ϑ_2 , this agreement holds for the leading (order ϵ) coefficient.

All couplings of the theory have become fully dynamical at NNLO. At N^3LO in the expansion, no new consistency conditions arise. Instead, higher loop corrections will lead to higher order corrections in the results established thus far. Based the observations above, we expect that all coefficients up to ϵ^4 (ϵ^2) [ϵ] of the universal eigenvalues ϑ_1 (ϑ_2) [$\vartheta_{3,4}$] at NNLO in (3.32) are unaffected at N^3LO and beyond.

4.2 Weyl consistency

At a more fundamental level an argument known as Weyl consistency conditions [104, 105] lends a formal derivation of this hierarchical procedure of table 2. Replacing the couplings (3.7) by the set $\{g_i\} \equiv \{g, y, u, v\}$ with β functions $\beta_i = dg_i/d\ln \mu$, the Weyl consistency condition

$$\frac{\partial \beta^j}{\partial g_i} = \frac{\partial \beta^i}{\partial g_j} \tag{4.1}$$

relates partial derivatives of the various β functions to each other, and $\beta^i \equiv \chi^{ij} \beta_j$. The functions χ^{ij} plays the role of a metric in the space of couplings. The relations are expected

coupling	order in perturbation theory		
	α_g	1	2
α_y	0	1	2
α_h	0	0	1
α_v	0	0	1
approximation level	LO	NLO	NNLO

Table 2. Relation between approximation level and the loop order to which couplings are retained in perturbation theory.

to hold in the full theory, and hence it is desirable to obey (4.1) even within finite approximations. The crucial point here is that the metric itself is a function of the couplings. Therefore, a consistent solution to (4.1) will generically relate different orders within a *naïve* fixed-order perturbation theory. In [106] it was shown that these conditions hold for the standard model. For the gauge-Yukawa theory studied here, the metric χ has been given explicitly in [68] showing that the ordering laid out in table 2 is consistent with (4.1).

4.3 Universality

For our explicit computations we have used known RG equations in the MS-bar regularisation scheme. In general, the expansion coefficients of β -functions are non-universal numbers and depend on the adopted scheme. On the other hand, it is well-known that one-loop RG coefficients for couplings with vanishing mass dimension are scheme-independent and universal. Furthermore, the two-loop gauge contribution to the gauge β -function is also known to be universal, provided a mass-scale independent regularisation scheme is adopted. Coefficients at higher loop order are strictly non-universal. The main new effect in our work arises from the two-loop coefficients in the gauge sector, and from the interacting UV fixed point in the Yukawa RG flow at one-loop (3.15). Expressing the Yukawa fixed point in terms of the gauge coupling $\alpha_y^* = \alpha_y^*(\alpha_g)$, one then shows that the fixed point in the gauge sector is invariant to leading order in ϵ under perturbative (non-singular) reparametrisations $\alpha_g \rightarrow \alpha'_g(\alpha_g)$, see (3.15). We therefore conclude that the interacting UV fixed point arises universally, irrespective of the regularisation scheme.

4.4 Operator ordering

Unlike in asymptotically free theories, at an interacting UV fixed point it is not known beforehand which invariants will become relevant since canonical power counting cannot be taken for granted [54]. For asymptotically safe theories with perturbatively small anomalous dimensions and corrections-to-scaling, however, canonical power counting can again be used to conclude that invariants with canonical mass dimension larger than four will remain irrelevant at a perturbative UV fixed point. The reason for this is that corrections to scaling, in the regime (3.12), are too small to change canonical scaling dimensions by an integer, and hence cannot change irrelevant into relevant operators. If masses are switched-on, two such operators are the fermion and scalar mass terms, both of which receive only




Figure 6. The fourfold degeneracy of the classically marginal invariants (3.1), (3.3), (3.5) and (3.6) — schematically indicated by a thick grey line (left panel) — is lifted by residual interactions in the UV, $0 < \epsilon$ (right panel). Also shown are the universal eigenvalues $\vartheta_1 < 0 < \vartheta_2 < \vartheta_4 < \vartheta_3$ (bottom to top, respectively) of the fixed point FP_1 , and the interaction-induced gaps Δ in the eigenvalue spectrum at NNLO as functions of ϵ .

perturbatively small corrections at the fixed point. We conclude that the relevancy of operators continues to be controlled by their canonical mass dimension [54].

On the other hand, residual interactions, even if perturbatively weak, control the scaling of invariants which classically have a vanishing canonical mass dimension and can change these into relevant or irrelevant ones, see figure 6. In our model, we find that the operator ordering of the classically marginal invariants at the fixed point is reflected by our search strategy, see table 2. At LO, the $SU(N_C)$ Yang-Mills Lagrangean (3.1) coupled to N_F fermions (3.2) is assumed to become a relevant operator in the regime (3.12) because asymptotic freedom is lost. This assumption is tested and confirmed at NLO against the inclusion of Yukawa interactions (3.3). The eigenvalue ϑ_1 is dominated by the gauge and ϑ_2 dominated by the Yukawa coupling. This is consistent with the initial assumption inasmuch as the scaling of the Yukawa term provides a subleading correction to the scaling of the Yang-Mills term. At NNLO, two quartic scalar selfinteractions are introduced whose non-trivial fixed points add two eigenvalues to the spectrum. At FP_1 , both of these are irrelevant. At FP_2 , the double trace scalar selfinteraction becomes relevant. The structure of the scalar sector is induced by the fixed point in the gauge-Yukawa subsector. In general, for other values of the gauge and Yukawa couplings, the scalar sector may not offer a fixed point at all.

4.5 Gap

Residual interactions at the UV fixed point have lifted the fourfold degeneracy amongst the classically marginal couplings. In figure 6, we show the eigenvalues to leading order in ϵ at the fixed point FP_1 , except for ϑ_1 which is shown at order ϵ^2 . The difference between the smallest negative and the smallest positive eigenvalue, which we denote as the gap of the eigenvalue spectrum Δ , is then a good quantitative measure for the strength of residual interactions. At the UV fixed point we have $\Delta = \vartheta_2 - \vartheta_1$. Quantitatively, the gap in the

eigenvalue spectrum read

$$\Delta = \frac{52}{19}\epsilon + \mathcal{O}(\epsilon^2), \tag{4.2}$$

where the (sub)leading term in ϵ arises from the (N)NLO approximation. Classically, we have $\Delta = 0$. We notice that the leading and subleading term have the same sign, increasing the gap with increasing ϵ . We stress that the gap in the eigenvalue spectrum is insensitive to the details of the scalar sector and only determined by the gauge-Yukawa subsystem.

4.6 Unitarity

An important constraint on quantum corrections relates to the scaling dimension of primary fields such as scalar fields themselves. For a quantum theory to be compatible with unitarity, it is required that the scaling dimension must be larger than unity, $\Delta_H > 1$. This behaviour can be observed in the result. To leading order in ϵ , γ_H is negative and hence $\Delta_H > 1$. At NNLO, we observe cancellations in (3.51) ensuring that γ_H remains negative. Overall, fluctuation-induced corrections reach values of up to 5% for moderate ϵ .

For the composite scalar operator $\delta^{ij}\bar{Q}_i Q_j$, the leading order corrections in ϵ decrease its scaling dimension Δ_F below its classical value $\Delta_F = 3$, see (3.54). This is further decreased at NNLO where all corrections to Δ_F have the same sign and no cancellations occur. The NNLO corrections are thus stronger than those for Δ_H . Here, corrections push Δ_F down from its classical value by up to 10%, leaving $\Delta_F > 1$. We conclude that the effects of residual interactions are compatible with basic constraints on the scaling of scalar operators.

4.7 Triviality

Triviality bounds often arise when infrared free interactions display a perturbative Landau pole towards high energies, limiting the predictivity of the theory to the scale of maximal UV extension [5]. On a more fundamental level, triviality relates to the difficulty of defining a self-interacting scalar quantum field in four dimensions [60, 107–109], which also puts the existence of elementary scalars into question. In the standard model, the scalar and the U(1) sectors are infrared free. At the UV fixed points detected here, triviality for all three types of fields is evaded through residual interactions. This also indicates that the scalar degrees of freedom may indeed be taken as elementary.

Moreover, we also observe that the avoidance of triviality in the scalar sector is closely linked to the presence of gauge fields, be they asymptotically free or asymptotically safe. In fact, an interacting fixed point in the scalar sector would not arise without an interacting fixed point for the Yukawa coupling, see (3.24), (3.25). Furthermore, without gauge fields, the fermion-boson subsystem does not generate an interacting UV fixed point, and couplings cannot reach the Gaussian fixed point in the UV. With asymptotically free gauge fields (say, for small $\epsilon < 0$), the UV fixed point for the Yukawa coupling remains the trivial one, see (3.15), (3.26). A detailed inspection then shows that complete asymptotic freedom follows, albeit under certain constraints on the parameters [110]. With asymptotically safe gauge fields (for small $\epsilon > 0$), complete asymptotic safety is achieved at two interacting UV fixed points (see table 1). We conclude that triviality is evaded in the large- N limit with and without asymptotic freedom in the gauge sector, although the specific details differ.

Asymptotic freedom in the gauge sector had to be given up for the Yukawa and scalar sectors to develop interacting UV fixed points.

5 Towards asymptotic safety at strong coupling

It would be useful to understand the existence or not of UV fixed points in non-Abelian gauge theories with matter and away from the regime where asymptotic safety is realised perturbatively and ϵ is small. In this section, we indicate some directions towards larger ϵ , with and without scalar matter.

5.1 Beyond the Veneziano limit

Presently, our study is bound to the second nontrivial order within perturbation theory and to the leading order in $1/N_F, 1/N_C \ll 1$, allowing for an accurate determination of the UV fixed point in the regime (3.12). The stability in the result makes it conceivable that the UV fixed point may persist even for finite values of ϵ . With increasing ϵ , the upper bound (3.31) which has arisen at NNLO comes into play. Solutions (N_C, N_F) to the constraint

$$0 \leq \epsilon(N_C, N_F) < \epsilon_{\max}, \tag{5.1}$$

where we take for ϵ_{\max} its value at NNLO given in (3.31), would then be likely candidate theories where the fixed point may exist even for finite but small couplings. The first few such solutions with the smallest numbers of fields are

$$(N_C, N_F) = (5, 28), (7, 39), (9, 50), (10, 56), (11, 61), (12, 67), \dots \tag{5.2}$$

Once $N_C > 12$, more than one solution for N_F may exist. Extending our study to N³LO should improve the estimate for the window (5.1) for large N . For finite values of N_F and N_C , the existence of an asymptotically safe window can in principle be tested using non-perturbative tools such as functional renormalisation [111–114], or the lattice.

5.2 Infinite order perturbation theory

Interestingly, an infinite order result is available for nonabelian gauge theories with a finite number of colors $N_C < \infty$, without scalars, but with $N_F \rightarrow \infty$ many Dirac fermions transforming according to a given representation of the gauge group [115], see also [116, 117] and references therein. In the terminology of this work, this corresponds to the parameter regime

$$1 \ll \epsilon, \tag{5.3}$$

see (3.11). In this limit the parametric deviation from asymptotic freedom is large, and the model becomes partly abelian [118]. Defining $x = 4N_F T_R \alpha$ with $\alpha = g^2/(4\pi)^2$ and T_F the trace normalization, it is possible to sum exactly the infinite perturbative series for the gauge β -function for large numbers of flavors. The all-order result has the form [116, 118, 119]

$$\frac{3}{2x} \frac{\beta(x)}{x} = 1 + \frac{H(x)}{N_F} + \mathcal{O}(N_F^{-2}) \tag{5.4}$$




Figure 7. The fully resummed gauge β function (5.4) is shown to leading order in $1/N_F$ including the Gaussian and the UV fixed point ($N_C = 5$ and $N_F = 28$).

and an integral representation of $H(x)$ can be found in [116] (see figure 7 for an example). Adopting $T_F = \frac{1}{2}$, one can show that the function $H(x)$ is finite for $0 \leq x < 3$ with a negative logarithmic singularity at $x = 3$ where $H(x) = N_C/8 \times \ln|3-x| + \text{const.} + \mathcal{O}(3-x)$. (Similar results are found for other representations as well.) This structure implies the existence of a nontrivial UV fixed point to leading order in $1/N_F$. As a word of caution, however, we remind the reader that an infinite order perturbative result may be upset non-perturbatively, or by higher order terms in N_F^{-1} (see [116] for a discussion of the latter in QED). Expanding about the fixed point, the two leading terms read

$$\alpha^* = \frac{3}{2 N_F} - \frac{1}{2 N_F} \exp\left(-a \cdot \frac{N_F}{N_C} + b(N_C)\right) \tag{5.5}$$

where $a = 8$, and $b(N_C) \simeq 15.857 + 2.632/N_C^2$. The UV fixed point starts dominating the RG running once $(\alpha^* - \alpha)/\alpha^* \lesssim N_C/(16 N_F)$ and its basin of attraction becomes algebraically small for large N_F ; see figure 7. Using the explicit form for $H(x)$ we also find the universal eigenvalue at the fixed point (5.5) of the infinite order β -function (5.4),

$$\vartheta = -\frac{3 N_C}{4 N_F} \exp\left(a \cdot \frac{N_F}{N_C} - b(N_C)\right) \tag{5.6}$$

By construction, the result (5.6) is valid in the limit $N_C/N_F \ll 1$ where the eigenvalue becomes parametrically large. The exponent $\nu = -1/\vartheta$ for the correlation length becomes very small, $\nu \rightarrow 0$. Eigenvalues which grow rapidly with the number of degrees of freedom have been observed previously for quantum gravity in the large dimensional limit in the continuum [17, 18, 73] and from lattice considerations [120].

5.3 Finite order perturbation theory

The origin of asymptotic safety in Yang-Mills theory with (5.5), (5.6) is different from the one observed in section 3, because the vanishing of the gauge β -function (5.4) arises as

an infinite order effect due to gluon and fermion loops for large ϵ , rather than through an order-by-order cancellation of fluctuations from gauge, fermion and scalar fields for small ϵ . It would be useful to understand whether the result (5.5) persists beyond the limit of infinite N_F with fixed and finite N_C . To that end, we test the continuity of the fixed point in (N_F, N_C) by combining two observations. Firstly, we notice that a precursor of the fixed point (5.5) is already visible within perturbation theory at finite orders. To see this explicitly, we come back to our equations at NNLO and switch off the Yukawa and scalar coupling, $\alpha_h = 0, \alpha_y = 0$ and $\alpha_v = 0$. In the parameter regime (5.3), we then find the UV fixed point $\alpha_g^* = 3/(2\sqrt{7}\epsilon) + \text{subleading}$, and the eigenvalue $\vartheta = -4\sqrt{\epsilon}/\sqrt{7} + \text{subleading}$. Adopting the same definition for the coupling as in (5.5), this result translates into

$$\begin{aligned} \alpha^* &= \frac{3}{2\sqrt{7}} \frac{1}{\sqrt{N_C N_F}}, \\ \vartheta &= -\frac{4}{\sqrt{7}} \sqrt{\frac{N_F}{N_C}}, \end{aligned} \tag{5.7}$$

to leading order in $1/\epsilon$ and $1/N_F$. A few comments are in order. Comparing (5.7) with (5.5), (5.6) for fixed N_C , we find that the $1/N_F$ decay of the fixed point is replaced by a softer square-root decay due to the finite order approximation in perturbation theory. The non-analytic dependence on N_F and N_C develops into the result (5.5) with increasing orders in perturbation theory where the power law behaviour becomes $\alpha^* \sim N_F^{(2-p)/(p-1)}$ [115], provided the p -loop coefficient is negative [117]. We also find that the eigenvalue ϑ in (5.7) grows large in the regime (5.3), modulo subleading corrections. While the growth rate $\vartheta \sim -\sqrt{N_F}$ in (5.7) is weaker than the one observed in (5.6), the correlation length exponent ν shows the same qualitative behaviour $\nu \rightarrow 0$ as the infinite order fixed point. We thus may conclude that (5.7) is the low-order precursor to the all-order result (5.5).

Secondly, for the finite order fixed point (5.7) we observe that the limits $1/N_C \rightarrow 0$ with N_F/N_C fixed can be accessed, and hence finite values for ϵ with (5.3), because the underlying NNLO equations remain valid in this parameter regime. Note that this limit is not covered by the rationale which has led to (5.4). The UV fixed point then reads

$$\alpha^* = \frac{33 + 6\epsilon}{4\sqrt{7}\epsilon} \frac{1}{N_F}. \tag{5.8}$$

For fixed ϵ , the fixed point shows the same $1/N_F$ behaviour as the fixed point (5.5). The coefficient in front of $1/N_F$ in (5.8) is larger than the fixed point (5.5) for all finite ϵ . Unlike (5.6), its eigenvalue (5.7) remains bounded since N_F/N_C is finite. It would thus seem that the inclusion of more gluons or less fermions maintains the UV fixed point, albeit with a softened UV scaling behaviour and at stronger coupling. The continuity of results in (N_F, N_C) suggests that the UV fixed point (5.5) is not an artefact of the large- N_F limit, but rather a fingerprint of a fixed point in the physical theory.

In summary, the observations in this section indicate that the matter-gauge systems studied here have a sufficiently rich structure to admit asymptotically safe UV fixed points also for finite (N_C, N_F) , with and without scalar matter, in addition to the weakly coupled UV fixed point for small ϵ . More work is required to identify them reliably within perturbation theory and beyond, and for generic values of ϵ .

6 Conclusion

We have used large- N techniques to understand the ultraviolet behaviour of theories involving fundamental gauge fields, fermions, and scalars. In strictly four space-time dimensions, and in the regime where the gauge sector is no longer asymptotically free, we have identified a perturbative origin for asymptotic safety. We found that all three types of fields are necessary for an interacting UV fixed point to arise. The primary driver towards asymptotic safety are the Yukawa interactions, which source the interacting fixed point for both the gauge fields and the scalars. In return, the gauge fields stabilise an interacting fixed point in the Yukawa sector. Fixed points are established in the perturbative domain, consistent with unitarity. Triviality bounds and Landau poles are evaded. Here the scalar fields can be considered as elementary.

It would be worth extending this picture within perturbation theory and beyond, also taking subleading corrections into consideration, and for fields with more general gauge charges, gauge groups, and Yukawa interactions. Once the number of fields is finite, asymptotic safety can in principle be tested non-perturbatively using the powerful machinery of functional renormalisation [111–114], or the lattice. In a different vein, one might wonder whether the weakly coupled ultraviolet fixed point has a strongly coupled dual. First steps to extend the ideas of Seiberg duality [121] to non-supersymmetric theories have been discussed in [122, 123]. It has also been suggested that UV conformal matter could simplify the quantisation of canonical gravity [39], or help to resolve outstanding puzzles in particle physics and cosmology. Our study offers such candidates.

Acknowledgments

This work is supported by the Science Technology and Facilities Council (STFC) [grant number ST/J000477/1], by the National Science Foundation under Grant No. PHYS-1066293, and by the hospitality of the Aspen Center for Physics. The CP³-Origins centre is partially funded by the Danish National Research Foundation, grant number DNRF90.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] K.G. Wilson, *Renormalization group and critical phenomena. 1. Renormalization group and the Kadanoff scaling picture*, *Phys. Rev.* **B 4** (1971) 3174 [[INSPIRE](#)].
- [2] K.G. Wilson, *Renormalization group and critical phenomena. 2. Phase space cell analysis of critical behavior*, *Phys. Rev.* **B 4** (1971) 3184 [[INSPIRE](#)].
- [3] D.J. Gross and F. Wilczek, *Ultraviolet behavior of non-Abelian gauge theories*, *Phys. Rev. Lett.* **30** (1973) 1343 [[INSPIRE](#)].
- [4] H.D. Politzer, *Reliable perturbative results for strong interactions?*, *Phys. Rev. Lett.* **30** (1973) 1346 [[INSPIRE](#)].

- [5] D.J.E. Callaway, *Triviality pursuit: can elementary scalar particles exist?*, *Phys. Rept.* **167** (1988) 241 [INSPIRE].
- [6] S. Weinberg, *Ultraviolet divergences in quantum theories of gravitation*, in *General relativity: an Einstein centenary survey*, S.W. Hawking and W. Israel eds., (1979), pg. 790 [INSPIRE].
- [7] D.F. Litim, *Renormalisation group and the Planck scale*, *Phil. Trans. Roy. Soc. Lond. A* **369** (2011) 2759 [arXiv:1102.4624] [INSPIRE].
- [8] D.F. Litim, *On fixed points of quantum gravity*, *AIP Conf. Proc.* **841** (2006) 322 [hep-th/0606044] [INSPIRE].
- [9] M. Niedermaier, *The asymptotic safety scenario in quantum gravity: an introduction*, *Class. Quant. Grav.* **24** (2007) R171 [gr-qc/0610018] [INSPIRE].
- [10] M. Niedermaier and M. Reuter, *The asymptotic safety scenario in quantum gravity*, *Living Rev. Rel.* **9** (2006) 5 [INSPIRE].
- [11] R. Percacci, *Asymptotic safety*, arXiv:0709.3851 [INSPIRE].
- [12] D.F. Litim, *Fixed points of quantum gravity and the renormalisation group*, *PoS(QG-Ph)024* [arXiv:0810.3675] [INSPIRE].
- [13] M. Reuter and F. Saueressig, *Quantum Einstein gravity*, *New J. Phys.* **14** (2012) 055022 [arXiv:1202.2274] [INSPIRE].
- [14] D.I. Kazakov, *Ultraviolet fixed points in gauge and SUSY field theories in extra dimensions*, *JHEP* **03** (2003) 020 [hep-th/0209100] [INSPIRE].
- [15] H. Gies, J. Jaeckel and C. Wetterich, *Towards a renormalizable standard model without fundamental Higgs scalar*, *Phys. Rev. D* **69** (2004) 105008 [hep-ph/0312034] [INSPIRE].
- [16] T.R. Morris, *Renormalizable extra-dimensional models*, *JHEP* **01** (2005) 002 [hep-ph/0410142] [INSPIRE].
- [17] P. Fischer and D.F. Litim, *Fixed points of quantum gravity in extra dimensions*, *Phys. Lett. B* **638** (2006) 497 [hep-th/0602203] [INSPIRE].
- [18] P. Fischer and D.F. Litim, *Fixed points of quantum gravity in higher dimensions*, *AIP Conf. Proc.* **861** (2006) 336 [hep-th/0606135] [INSPIRE].
- [19] D.I. Kazakov and G.S. Vartanov, *Renormalizable $1/N_f$ expansion for field theories in extra dimensions*, *JHEP* **06** (2007) 081 [arXiv:0707.2564] [INSPIRE].
- [20] O. Zanusso, L. Zambelli, G.P. Vacca and R. Percacci, *Gravitational corrections to Yukawa systems*, *Phys. Lett. B* **689** (2010) 90 [arXiv:0904.0938] [INSPIRE].
- [21] H. Gies, S. Rechenberger and M.M. Scherer, *Towards an asymptotic-safety scenario for chiral Yukawa systems*, *Eur. Phys. J. C* **66** (2010) 403 [arXiv:0907.0327] [INSPIRE].
- [22] J.-E. Daum, U. Harst and M. Reuter, *Running gauge coupling in asymptotically safe quantum gravity*, *JHEP* **01** (2010) 084 [arXiv:0910.4938] [INSPIRE].
- [23] G.P. Vacca and O. Zanusso, *Asymptotic safety in Einstein gravity and scalar-fermion matter*, *Phys. Rev. Lett.* **105** (2010) 231601 [arXiv:1009.1735] [INSPIRE].
- [24] X. Calmet, *Asymptotically safe weak interactions*, *Mod. Phys. Lett. A* **26** (2011) 1571 [arXiv:1012.5529] [INSPIRE].
- [25] S. Folkerts, D.F. Litim and J.M. Pawłowski, *Asymptotic freedom of Yang-Mills theory with gravity*, *Phys. Lett. B* **709** (2012) 234 [arXiv:1101.5552] [INSPIRE].

- [26] F. Bazzocchi, M. Fabbrichesi, R. Percacci, A. Tonero and L. Vecchi, *Fermions and Goldstone bosons in an asymptotically safe model*, *Phys. Lett. B* **705** (2011) 388 [[arXiv:1105.1968](#)] [[INSPIRE](#)].
- [27] H. Gies, S. Rechenberger, M.M. Scherer and L. Zambelli, *An asymptotic safety scenario for gauged chiral Higgs-Yukawa models*, *Eur. Phys. J. C* **73** (2013) 2652 [[arXiv:1306.6508](#)] [[INSPIRE](#)].
- [28] O. Antipin, M. Mojaza and F. Sannino, *Conformal extensions of the standard model with Veltman conditions*, *Phys. Rev. D* **89** (2014) 085015 [[arXiv:1310.0957](#)] [[INSPIRE](#)].
- [29] P. Doná, A. Eichhorn and R. Percacci, *Matter matters in asymptotically safe quantum gravity*, *Phys. Rev. D* **89** (2014) 084035 [[arXiv:1311.2898](#)] [[INSPIRE](#)].
- [30] A. Bonanno and M. Reuter, *Cosmology of the Planck era from a renormalization group for quantum gravity*, *Phys. Rev. D* **65** (2002) 043508 [[hep-th/0106133](#)] [[INSPIRE](#)].
- [31] K.A. Meissner and H. Nicolai, *Conformal symmetry and the standard model*, *Phys. Lett. B* **648** (2007) 312 [[hep-th/0612165](#)] [[INSPIRE](#)].
- [32] R. Foot, A. Kobakhidze, K.L. McDonald and R.R. Volkas, *A solution to the hierarchy problem from an almost decoupled hidden sector within a classically scale invariant theory*, *Phys. Rev. D* **77** (2008) 035006 [[arXiv:0709.2750](#)] [[INSPIRE](#)].
- [33] J. Hewett and T. Rizzo, *Collider signals of gravitational fixed points*, *JHEP* **12** (2007) 009 [[arXiv:0707.3182](#)] [[INSPIRE](#)].
- [34] D.F. Litim and T. Plehn, *Signatures of gravitational fixed points at the LHC*, *Phys. Rev. Lett.* **100** (2008) 131301 [[arXiv:0707.3983](#)] [[INSPIRE](#)].
- [35] M. Shaposhnikov and D. Zenhausern, *Quantum scale invariance, cosmological constant and hierarchy problem*, *Phys. Lett. B* **671** (2009) 162 [[arXiv:0809.3406](#)] [[INSPIRE](#)].
- [36] M. Shaposhnikov and D. Zenhausern, *Scale invariance, unimodular gravity and dark energy*, *Phys. Lett. B* **671** (2009) 187 [[arXiv:0809.3395](#)] [[INSPIRE](#)].
- [37] M. Shaposhnikov and C. Wetterich, *Asymptotic safety of gravity and the Higgs boson mass*, *Phys. Lett. B* **683** (2010) 196 [[arXiv:0912.0208](#)] [[INSPIRE](#)].
- [38] S. Weinberg, *Asymptotically safe inflation*, *Phys. Rev. D* **81** (2010) 083535 [[arXiv:0911.3165](#)] [[INSPIRE](#)].
- [39] G. 't Hooft, *Probing the small distance structure of canonical quantum gravity using the conformal group*, [arXiv:1009.0669](#) [[INSPIRE](#)].
- [40] E. Gerwick, D. Litim and T. Plehn, *Asymptotic safety and Kaluza-Klein gravitons at the LHC*, *Phys. Rev. D* **83** (2011) 084048 [[arXiv:1101.5548](#)] [[INSPIRE](#)].
- [41] E. Gerwick, *Asymptotically safe gravitons in electroweak precision physics*, *Eur. Phys. J. C* **71** (2011) 1676 [[arXiv:1012.1118](#)] [[INSPIRE](#)].
- [42] M. Hindmarsh, D. Litim and C. Rahmede, *Asymptotically safe cosmology*, *JCAP* **07** (2011) 019 [[arXiv:1101.5401](#)] [[INSPIRE](#)].
- [43] T. Hur and P. Ko, *Scale invariant extension of the standard model with strongly interacting hidden sector*, *Phys. Rev. Lett.* **106** (2011) 141802 [[arXiv:1103.2571](#)] [[INSPIRE](#)].
- [44] B. Dobrich and A. Eichhorn, *Can we see quantum gravity? Photons in the asymptotic-safety scenario*, *JHEP* **06** (2012) 156 [[arXiv:1203.6366](#)] [[INSPIRE](#)].
- [45] G. Marques Tavares, M. Schmaltz and W. Skiba, *Higgs mass naturalness and scale invariance in the UV*, *Phys. Rev. D* **89** (2014) 015009 [[arXiv:1308.0025](#)] [[INSPIRE](#)].

- [46] C. Tamarit, *Running couplings with a vanishing scale anomaly*, *JHEP* **12** (2013) 098 [[arXiv:1309.0913](#)] [[INSPIRE](#)].
- [47] S. Abel and A. Mariotti, *Novel Higgs potentials from gauge mediation of exact scale breaking*, [arXiv:1312.5335](#) [[INSPIRE](#)].
- [48] O. Antipin, J. Krog, M. Mojaza and F. Sannino, *Stable Extensions with(out) gravity*, *Nucl. Phys. B* **886** (2014) 125 [[arXiv:1311.1092](#)] [[INSPIRE](#)].
- [49] M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann and K. Tuominen, *Physical naturalness and dynamical breaking of classical scale invariance*, *Mod. Phys. Lett. A* **29** (2014) 1450077 [[arXiv:1304.7006](#)] [[INSPIRE](#)].
- [50] E. Gabrielli et al., *Towards completing the standard model: vacuum stability, EWSB and dark matter*, *Phys. Rev. D* **89** (2014) 015017 [[arXiv:1309.6632](#)] [[INSPIRE](#)].
- [51] M. Holthausen, J. Kubo, K.S. Lim and M. Lindner, *Electroweak and conformal symmetry breaking by a strongly coupled hidden sector*, *JHEP* **12** (2013) 076 [[arXiv:1310.4423](#)] [[INSPIRE](#)].
- [52] G.C. Dorsch, S.J. Huber and J.M. No, *Cosmological signatures of a UV-conformal standard model*, *Phys. Rev. Lett.* **113** (2014) 121801 [[arXiv:1403.5583](#)] [[INSPIRE](#)].
- [53] A. Eichhorn and M.M. Scherer, *The Planck scale, the Higgs mass and scalar dark matter*, *Phys. Rev. D* **90** (2014) 025023 [[arXiv:1404.5962](#)] [[INSPIRE](#)].
- [54] K. Falls, D.F. Litim, K. Nikolakopoulos and C. Rahmede, *A bootstrap towards asymptotic safety*, [arXiv:1301.4191](#) [[INSPIRE](#)].
- [55] R. Gastmans, R. Kallosh and C. Truffin, *Quantum gravity near two-dimensions*, *Nucl. Phys. B* **133** (1978) 417 [[INSPIRE](#)].
- [56] S.M. Christensen and M.J. Duff, *Quantum gravity in two + ϵ dimensions*, *Phys. Lett. B* **79** (1978) 213 [[INSPIRE](#)].
- [57] M.E. Peskin, *Critical point behavior of the Wilson loop*, *Phys. Lett. B* **94** (1980) 161 [[INSPIRE](#)].
- [58] K. Gawędzki and A. Kupiainen, *Exact renormalization for the Gross-Neveu model of quantum fields*, *Phys. Rev. Lett.* **54** (1985) 2191 [[INSPIRE](#)].
- [59] K. Gawędzki and A. Kupiainen, *Renormalizing the nonrenormalizable*, *Phys. Rev. Lett.* **55** (1985) 363 [[INSPIRE](#)].
- [60] K.G. Wilson and J.B. Kogut, *The renormalization group and the ϵ -expansion*, *Phys. Rept.* **12** (1974) 75 [[INSPIRE](#)].
- [61] E. Tomboulis, *1/N expansion and renormalization in quantum gravity*, *Phys. Lett. B* **70** (1977) 361 [[INSPIRE](#)].
- [62] E. Tomboulis, *Renormalizability and asymptotic freedom in quantum gravity*, *Phys. Lett. B* **97** (1980) 77 [[INSPIRE](#)].
- [63] L. Smolin, *A fixed point for quantum gravity*, *Nucl. Phys. B* **208** (1982) 439 [[INSPIRE](#)].
- [64] C. de Calan, P.A. Faria da Veiga, J. Magnen and R. Seneor, *Constructing the three-dimensional Gross-Neveu model with a large number of flavor components*, *Phys. Rev. Lett.* **66** (1991) 3233 [[INSPIRE](#)].
- [65] O. Antipin, M. Mojaza, C. Pica and F. Sannino, *Magnetic fixed points and emergent supersymmetry*, *JHEP* **06** (2013) 037 [[arXiv:1105.1510](#)] [[INSPIRE](#)].

- [66] O. Antipin, M. Mojaza and F. Sannino, *Light dilaton at fixed points and ultra light scale super Yang-Mills*, *Phys. Lett. B* **712** (2012) 119 [[arXiv:1107.2932](#)] [[INSPIRE](#)].
- [67] O. Antipin, S. Di Chiara, M. Mojaza, E. Mølgaard and F. Sannino, *A perturbative realization of Miransky scaling*, *Phys. Rev. D* **86** (2012) 085009 [[arXiv:1205.6157](#)] [[INSPIRE](#)].
- [68] O. Antipin, M. Gillioz, E. Mølgaard and F. Sannino, *The a theorem for gauge-Yukawa theories beyond Banks-Zaks fixed point*, *Phys. Rev. D* **87** (2013) 125017 [[arXiv:1303.1525](#)] [[INSPIRE](#)].
- [69] G. Veneziano, *U(1) without instantons*, *Nucl. Phys. B* **159** (1979) 213 [[INSPIRE](#)].
- [70] T. Banks and A. Zaks, *On the phase structure of vector-like gauge theories with massless fermions*, *Nucl. Phys. B* **196** (1982) 189 [[INSPIRE](#)].
- [71] H. Kawai, Y. Kitazawa and M. Ninomiya, *Scaling exponents in quantum gravity near two-dimensions*, *Nucl. Phys. B* **393** (1993) 280 [[hep-th/9206081](#)] [[INSPIRE](#)].
- [72] H. Kawai, Y. Kitazawa and M. Ninomiya, *Ultraviolet stable fixed point and scaling relations in $(2 + \epsilon)$ -dimensional quantum gravity*, *Nucl. Phys. B* **404** (1993) 684 [[hep-th/9303123](#)] [[INSPIRE](#)].
- [73] D.F. Litim, *Fixed points of quantum gravity*, *Phys. Rev. Lett.* **92** (2004) 201301 [[hep-th/0312114](#)] [[INSPIRE](#)].
- [74] A. Codello, R. Percacci and C. Rahmede, *Investigating the ultraviolet properties of gravity with a Wilsonian renormalization group equation*, *Annals Phys.* **324** (2009) 414 [[arXiv:0805.2909](#)] [[INSPIRE](#)].
- [75] D. Benedetti, P.F. Machado and F. Saueressig, *Asymptotic safety in higher-derivative gravity*, *Mod. Phys. Lett. A* **24** (2009) 2233 [[arXiv:0901.2984](#)] [[INSPIRE](#)].
- [76] I. Donkin and J.M. Pawłowski, *The phase diagram of quantum gravity from diffeomorphism-invariant RG-flows*, [arXiv:1203.4207](#) [[INSPIRE](#)].
- [77] N. Christiansen, D.F. Litim, J.M. Pawłowski and A. Rodigast, *Fixed points and infrared completion of quantum gravity*, *Phys. Lett. B* **728** (2014) 114 [[arXiv:1209.4038](#)] [[INSPIRE](#)].
- [78] N. Christiansen, B. Knorr, J.M. Pawłowski and A. Rodigast, *Global flows in quantum gravity*, [arXiv:1403.1232](#) [[INSPIRE](#)].
- [79] D. Becker and M. Reuter, *En route to background independence: broken split-symmetry and how to restore it with bi-metric average actions*, *Annals Phys.* **350** (2014) 225 [[arXiv:1404.4537](#)] [[INSPIRE](#)].
- [80] D.J. Gross and A. Neveu, *Dynamical symmetry breaking in asymptotically free field theories*, *Phys. Rev. D* **10** (1974) 3235 [[INSPIRE](#)].
- [81] Y. Kikukawa and K. Yamawaki, *Ultraviolet fixed point structure of renormalizable four fermion theory in less than four-dimensions*, *Phys. Lett. B* **234** (1990) 497 [[INSPIRE](#)].
- [82] H.-J. He, Y.-P. Kuang, Q. Wang and Y.-P. Yi, *Effective potential, renormalization and nontrivial ultraviolet fixed point in D -dimensional four fermion theories $(2 < D < 4)$ to order $1/N$ in $1/N$ expansion*, *Phys. Rev. D* **45** (1992) 4610 [[INSPIRE](#)].
- [83] S. Hands, A. Kocic and J.B. Kogut, *Four Fermi theories in fewer than four-dimensions*, *Annals Phys.* **224** (1993) 29 [[hep-lat/9208022](#)] [[INSPIRE](#)].
- [84] J. Braun, H. Gies and D.D. Scherer, *Asymptotic safety: a simple example*, *Phys. Rev. D* **83** (2011) 085012 [[arXiv:1011.1456](#)] [[INSPIRE](#)].

- [85] H. Gies, *Renormalizability of gauge theories in extra dimensions*, *Phys. Rev. D* **68** (2003) 085015 [[hep-th/0305208](#)] [[INSPIRE](#)].
- [86] A. Pelissetto and E. Vicari, *Critical phenomena and renormalization group theory*, *Phys. Rept.* **368** (2002) 549 [[cond-mat/0012164](#)] [[INSPIRE](#)].
- [87] D.F. Litim and D. Zappala, *Ising exponents from the functional renormalisation group*, *Phys. Rev. D* **83** (2011) 085009 [[arXiv:1009.1948](#)] [[INSPIRE](#)].
- [88] L. Fei, S. Giombi and I.R. Klebanov, *Critical $O(N)$ models in $6 - \epsilon$ dimensions*, *Phys. Rev. D* **90** (2014) 025018 [[arXiv:1404.1094](#)] [[INSPIRE](#)].
- [89] R. Percacci and G.P. Vacca, *Are there scaling solutions in the $O(N)$ -models for large- N in $d > 4$?*, *Phys. Rev. D* **90** (2014) 107702 [[arXiv:1405.6622](#)] [[INSPIRE](#)].
- [90] E. Brézin and J. Zinn-Justin, *Renormalization of the nonlinear σ -model in $2 + \epsilon$ dimensions. Application to the Heisenberg ferromagnets*, *Phys. Rev. Lett.* **36** (1976) 691 [[INSPIRE](#)].
- [91] W.A. Bardeen, B.W. Lee and R.E. Shrock, *Phase transition in the nonlinear σ -model in $2 + \epsilon$ dimensional continuum*, *Phys. Rev. D* **14** (1976) 985 [[INSPIRE](#)].
- [92] R. Percacci, *Asymptotic safety in gravity and σ -models*, *PoS(CLAQG08)002* [[arXiv:0910.4951](#)] [[INSPIRE](#)].
- [93] B.H. Wellegehausen, D. Körner and A. Wipf, *Asymptotic safety on the lattice: the nonlinear $O(N)$ σ -model*, *Annals Phys.* **349** (2014) 374 [[arXiv:1402.1851](#)] [[INSPIRE](#)].
- [94] M.E. Machacek and M.T. Vaughn, *Two loop renormalization group equations in a general quantum field theory. 1. Wave function renormalization*, *Nucl. Phys. B* **222** (1983) 83 [[INSPIRE](#)].
- [95] M.E. Machacek and M.T. Vaughn, *Two loop renormalization group equations in a general quantum field theory. 2. Yukawa couplings*, *Nucl. Phys. B* **236** (1984) 221 [[INSPIRE](#)].
- [96] M.E. Machacek and M.T. Vaughn, *Two loop renormalization group equations in a general quantum field theory. 3. Scalar quartic couplings*, *Nucl. Phys. B* **249** (1985) 70 [[INSPIRE](#)].
- [97] W.E. Caswell, *Asymptotic behavior of non-Abelian gauge theories to two loop order*, *Phys. Rev. Lett.* **33** (1974) 244 [[INSPIRE](#)].
- [98] F. Sannino, *Conformal dynamics for TeV physics and cosmology*, *Acta Phys. Polon. B* **40** (2009) 3533 [[arXiv:0911.0931](#)] [[INSPIRE](#)].
- [99] J. Kuti, *The Higgs particle and the lattice*, *PoS(LATTICE 2013)004* [[INSPIRE](#)].
- [100] F.J. Dyson, *Divergence of perturbation theory in quantum electrodynamics*, *Phys. Rev.* **85** (1952) 631 [[INSPIRE](#)].
- [101] E. Pomoni and L. Rastelli, *Large- N field theory and AdS tachyons*, *JHEP* **04** (2009) 020 [[arXiv:0805.2261](#)] [[INSPIRE](#)].
- [102] D.F. Litim, M. Mojaza and F. Sannino, *Vacuum stability of asymptotically safe gauge-Yukawa theories*, in preparation.
- [103] M.-X. Luo, H.-W. Wang and Y. Xiao, *Two loop renormalization group equations in general gauge field theories*, *Phys. Rev. D* **67** (2003) 065019 [[hep-ph/0211440](#)] [[INSPIRE](#)].
- [104] I. Jack and H. Osborn, *Analogs for the c theorem for four-dimensional renormalizable field theories*, *Nucl. Phys. B* **343** (1990) 647 [[INSPIRE](#)].

- [105] I. Jack and H. Osborn, *Constraints on RG flow for four dimensional quantum field theories*, *Nucl. Phys. B* **883** (2014) 425 [[arXiv:1312.0428](#)] [[INSPIRE](#)].
- [106] O. Antipin, M. Gillioz, J. Krog, E. Mølgaard and F. Sannino, *Standard model vacuum stability and Weyl consistency conditions*, *JHEP* **08** (2013) 034 [[arXiv:1306.3234](#)] [[INSPIRE](#)].
- [107] M. Lüscher and P. Weisz, *Scaling laws and triviality bounds in the lattice ϕ^4 theory. 2. One component model in the phase with spontaneous symmetry breaking*, *Nucl. Phys. B* **295** (1988) 65 [[INSPIRE](#)].
- [108] A. Hasenfratz, K. Jansen, C.B. Lang, T. Neuhaus and H. Yoneyama, *The triviality bound of the four component ϕ^4 model*, *Phys. Lett. B* **199** (1987) 531 [[INSPIRE](#)].
- [109] O.J. Rosten, *Triviality from the exact renormalization group*, *JHEP* **07** (2009) 019 [[arXiv:0808.0082](#)] [[INSPIRE](#)].
- [110] M. Harada, Y. Kikukawa, T. Kugo and H. Nakano, *Nontriviality of gauge Higgs-Yukawa system and renormalizability of gauged NJLS model*, *Prog. Theor. Phys.* **92** (1994) 1161 [[hep-ph/9407398](#)] [[INSPIRE](#)].
- [111] J. Polchinski, *Renormalization and effective Lagrangians*, *Nucl. Phys. B* **231** (1984) 269 [[INSPIRE](#)].
- [112] C. Wetterich, *Exact evolution equation for the effective potential*, *Phys. Lett. B* **301** (1993) 90 [[INSPIRE](#)].
- [113] T.R. Morris, *The exact renormalization group and approximate solutions*, *Int. J. Mod. Phys. A* **9** (1994) 2411 [[hep-ph/9308265](#)] [[INSPIRE](#)].
- [114] D.F. Litim, *Optimized renormalization group flows*, *Phys. Rev. D* **64** (2001) 105007 [[hep-th/0103195](#)] [[INSPIRE](#)].
- [115] C. Pica and F. Sannino, *β -function and anomalous dimensions*, *Phys. Rev. D* **83** (2011) 116001 [[arXiv:1011.3832](#)] [[INSPIRE](#)].
- [116] B. Holdom, *Large- N flavor β -functions: a recap*, *Phys. Lett. B* **694** (2010) 74 [[arXiv:1006.2119](#)] [[INSPIRE](#)].
- [117] R. Shrock, *Study of possible ultraviolet zero of the β -function in gauge theories with many fermions*, *Phys. Rev. D* **89** (2014) 045019 [[arXiv:1311.5268](#)] [[INSPIRE](#)].
- [118] A. Palanques-Mestre and P. Pascual, *The $1/N_f$ expansion of the γ and β -functions in QED*, *Commun. Math. Phys.* **95** (1984) 277 [[INSPIRE](#)].
- [119] J.A. Gracey, *The QCD β -function at $O(1/N_f)$* , *Phys. Lett. B* **373** (1996) 178 [[hep-ph/9602214](#)] [[INSPIRE](#)].
- [120] H.W. Hamber and R.M. Williams, *Quantum gravity in large dimensions*, *Phys. Rev. D* **73** (2006) 044031 [[hep-th/0512003](#)] [[INSPIRE](#)].
- [121] N. Seiberg, *Electric-magnetic duality in supersymmetric non-Abelian gauge theories*, *Nucl. Phys. B* **435** (1995) 129 [[hep-th/9411149](#)] [[INSPIRE](#)].
- [122] F. Sannino, *QCD dual*, *Phys. Rev. D* **80** (2009) 065011 [[arXiv:0907.1364](#)] [[INSPIRE](#)].
- [123] F. Sannino, *Magnetic S -parameter*, *Phys. Rev. Lett.* **105** (2010) 232002 [[arXiv:1007.0254](#)] [[INSPIRE](#)].