

Quantum metrology with mixed states: when recovering lost information is better than never losing it

Article (Published Version)

Haine, Simon A and Szigeti, Stuart S (2015) Quantum metrology with mixed states: when recovering lost information is better than never losing it. *Physical Review A*, 92 (3). p. 2317. ISSN 1050-2947

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/66133/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Quantum metrology with mixed states: When recovering lost information is better than never losing it

Simon A. Haine

School of Mathematics and Physics, University of Queensland, Brisbane, Queensland, 4072, Australia

Stuart S. Szigeti

*School of Mathematics and Physics, University of Queensland, Brisbane, Queensland, 4072, Australia
and ARC Centre for Engineered Quantum Systems, University of Queensland, Brisbane, Queensland 4072, Australia*

(Received 10 July 2015; published 17 September 2015)

Quantum-enhanced metrology can be achieved by entangling a probe with an auxiliary system, passing the probe through an interferometer, and subsequently making measurements on both the probe and auxiliary system. Conceptually, this corresponds to performing metrology with the purification of a (mixed) probe state. We demonstrate via the quantum Fisher information how to design mixed states whose purifications are an excellent metrological resource. In particular, we give examples of mixed states with purifications that allow (near) Heisenberg-limited metrology and provide examples of entangling Hamiltonians that can generate these states. Finally, we present the optimal measurement and parameter-estimation procedure required to realize these sensitivities (i.e., that saturate the quantum Cramér-Rao bound). Since pure states of comparable metrological usefulness are typically challenging to generate, it may prove easier to use this approach of entanglement and measurement of an auxiliary system. An example where this may be the case is atom interferometry, where entanglement with optical systems is potentially easier to engineer than the atomic interactions required to produce nonclassical atomic states.

DOI: [10.1103/PhysRevA.92.032317](https://doi.org/10.1103/PhysRevA.92.032317)

PACS number(s): 03.67.-a, 42.50.Dv, 42.50.St, 03.75.Dg

I. INTRODUCTION

There is currently great interest in quantum metrology, the science of estimating a classical parameter ϕ with a quantum probe at a higher precision than is possible with a classical probe of identical particle flux. Given a fixed number of particles, N , the ultimate limit to the sensitivity is the Heisenberg limit $\Delta\phi = 1/N$ [1,2]. Naïvely, the choice of probe state is a solved problem; for instance, symmetric Dicke states [1,3] and spin-cat states [4,5] input into a Mach-Zehnder (MZ) interferometer yield sensitivities of $\sqrt{2}/N$ and $1/N$, respectively. However, in practice achieving quantum-enhanced sensitivities is a significant challenge. This is due to both the deleterious effect of losses [6] and the challenges associated with preparing nonclassical states with an appreciable number of particles [7–11]. For example, protocols for generating a spin-cat state commonly require a large Kerr nonlinearity, which either is unavailable (e.g., in optical systems [12]), is difficult to engineer (e.g., in microwave cavities [13,14]), or detrimentally affects the efficient operation of the metrological device (e.g., atom interferometers where the interfering modes have large space-time separation [15–17]).

In this paper, we present an alternative route to quantum-enhanced metrology based on purifications of mixed states. Physically, this involves entangling the probe with an auxiliary system before the probe is affected by ϕ , making measurements on *both* the probe and auxiliary system, and subsequently using correlations between the two measurement outcomes in order to reduce the uncertainty in the estimated parameter (see Fig. 1). This approach is advantageous in cases where it is easier to entangle the probe system with another system, rather than directly create highly entangled states of the probe system itself. An example of this is atom interferometry; although quantum squeezing can be

produced in atomic systems via atomic interactions [18–31], the technical requirements of high-sensitivity, path-separated atom interferometers are better suited to enhancement via entanglement with an optical system [32–37] and information recycling [38–41].

The structure of this paper is as follows. In Sec. II we introduce in detail the central idea of this paper: that purifications of mixed states can possess a large quantum Fisher information (QFI) and therefore represent an excellent resource for quantum metrology. In Sec. III we specialize to an N -boson probe state and Mach-Zehnder (MZ) interferometer, and show how to engineer purifications that yield sensitivities at and near the Heisenberg limit. Finally, in Sec. IV we present optimal measurement schemes that allow these quantum-enhanced sensitivities to be achieved in practice.

II. QUANTUM FISHER INFORMATION FOR A PURIFICATION

We can determine the best sensitivity possible for any given metrology scheme via the QFI, \mathcal{F} , which places an absolute lower bound on the sensitivity, $\Delta\phi \geq 1/\sqrt{\mathcal{F}}$, called the quantum Cramér-Rao bound (QCRB) [42–45]. This bound is independent of the choice of measurement and parameter estimation procedure, and depends only on the input state. Explicitly, if a state $\hat{\rho}_A$ is input into a metrological device described by the unitary operator $\hat{U}_\phi = \exp(-i\phi\hat{G}_A)$, then the QFI is

$$\mathcal{F}_A \equiv \mathcal{F}[\hat{G}_A, \hat{\rho}_A] = 2 \sum_{i,j} \frac{(\lambda_i - \lambda_j)^2}{\lambda_i + \lambda_j} |\langle e_i | \hat{G}_A | e_j \rangle|^2, \quad (1)$$




FIG. 1. The unitary $\hat{U}_{AB} = \exp(-i\hat{\mathcal{H}}_{AB}t/\hbar)$ entangles system A (probe) with system B (auxiliary) before system A passes through a measurement device described by $\hat{U}_\phi = \exp(-i\phi\hat{G}_A)$. If measurements are restricted to system A, then the QFI for an estimate of ϕ is $\mathcal{F}_A = \mathcal{F}[\hat{G}_A, \hat{\rho}_A]$, where $\hat{\rho}_A = \text{Tr}_B\{|\Psi_{AB}\rangle\langle\Psi_{AB}|\}$. If measurements on both systems are permitted, then the QFI is $\mathcal{F}_{AB} = \mathcal{F}[\hat{G}_A, |\Psi_{AB}\rangle] = 4\text{Var}(\hat{G}_A)_{\hat{\rho}_A} \geq \mathcal{F}_A$.

where λ_i and $|e_i\rangle$ are the eigenvalues and eigenvectors of $\hat{\rho}_A$, respectively. If $\hat{\rho}_A$ is pure, then Eq. (1) reduces to $\mathcal{F}_A = 4\text{Var}(\hat{G}_A)$.

A naïve consideration of the pure state QFI suggests that engineering input states with a large variance in \hat{G}_A is an excellent strategy for achieving a high-precision estimate of ϕ . However, there are many operations on $\hat{\rho}_A$ that increase $\text{Var}(\hat{G}_A)$ at the expense of a *decrease* to the purity $\gamma = \text{Tr}\{\hat{\rho}_A^2\}$. Since the QFI is convex in the state, any process that mixes the state typically decreases the QFI. Consequently, any improvement due to a larger $\text{Var}(\hat{G}_A)$ is usually overwhelmed by reductions in the QFI due to mixing.

In order to concretely demonstrate this point, we focus on an N -boson state input into a MZ interferometer. As discussed in Ref. [46], this system is conveniently described by the SU(2) Lie algebra $[\hat{J}_i, \hat{J}_j] = i\epsilon_{ijk}\hat{J}_k$, where ϵ_{ijk} is the Levi-Civita symbol, for $i = x, y, z$. A MZ interferometer is characterized by $\hat{G}_A = \hat{J}_y$; therefore, for pure states, a large QFI requires a large $\text{Var}(\hat{J}_y)$.

Without loss of generality, we restrict ourselves to the class of input states

$$\hat{\rho}_A = \int_0^{2\pi} d\varphi \mathcal{P}(\varphi) \left| \alpha\left(\frac{\pi}{2}, \varphi\right) \right\rangle \left\langle \alpha\left(\frac{\pi}{2}, \varphi\right) \right|. \quad (2)$$

Here $|\alpha(\theta, \varphi)\rangle = \exp(-i\varphi\hat{J}_z)\exp(-i\theta\hat{J}_y)|j, j\rangle$ are spin coherent states, where $|j, m\rangle$ are Dicke states with total angular momentum $j = N/2$ and \hat{J}_z projection m . We focus on the following three states in class (2), which are in order of increasing $\text{Var}(\hat{J}_y)$:

$$\text{Case I: } \mathcal{P}(\varphi) = \delta(\varphi), \quad (3a)$$

$$\text{Case II: } \mathcal{P}(\varphi) = \frac{1}{2\pi}, \quad (3b)$$

$$\text{Case III: } \mathcal{P}(\varphi) = \frac{1}{2} \left[\delta\left(\varphi - \frac{\pi}{2}\right) + \delta\left(\varphi + \frac{\pi}{2}\right) \right]. \quad (3c)$$

These states can be conveniently visualized by plotting the Husimi- \mathcal{Q} function [47,48]

$$\mathcal{Q}(\theta, \varphi) = \frac{2j+1}{4\pi} \langle \alpha(\theta, \varphi) | \hat{\rho}_A | \alpha(\theta, \varphi) \rangle, \quad (4)$$

and the \hat{J}_y projection of the state, $P(J_y) = \langle J_y | \hat{\rho}_A | J_y \rangle$, where $\hat{J}_y |J_y\rangle = J_y |J_y\rangle$ (see Fig. 2).

None of these states yield sensitivities that surpass the standard quantum limit (SQL), $\Delta\phi = 1/\sqrt{N}$. In case I, $\hat{\rho}_A$ is a pure spin coherent state, $|\alpha(\pi/2, 0)\rangle$, with $\mathcal{F}_A = 4\text{Var}(\hat{J}_y) =$




FIG. 2. (Color online) Husimi- \mathcal{Q} function for case I (a), case II (b), and case III (c). The projection in the \hat{J}_y basis, $P(J_y)$ is shown for case I (d), case II (e), and case III (f). $N = 20$ for all frames.

N . In case II, $\hat{\rho}_A$ is an incoherent mixture of Dicke states (i.e., it contains no off-diagonal terms in the $|j, m\rangle$ basis). Although $4\text{Var}(\hat{J}_y) = N(N+1)/2$ is much larger than for case I, the QFI is only $\mathcal{F}_A = N/2$. Finally, case III is an incoherent mixture of maximal and minimal \hat{J}_y eigenstates with $4\text{Var}(\hat{J}_y) = N^2$, which is the maximum possible value in SU(2). However, since the state is mixed the QFI is significantly less than this, with $\mathcal{F}_A = N/2$.

However, suppose the mixing in $\hat{\rho}_A$ arises from entanglement with an auxiliary system B before system A passes through the metrological device (see Fig. 1). Specifically, for an input pure state $|\Psi_{AB}\rangle$ of a composite system $A \otimes B$, where $\hat{\rho}_A = \text{Tr}_B\{|\Psi_{AB}\rangle\langle\Psi_{AB}|\}$, the QFI is

$$\begin{aligned} \mathcal{F}_{AB} &\equiv \mathcal{F}[\hat{G}_A, |\Psi_{AB}\rangle] \\ &= 4(\langle\Psi_{AB}|\hat{G}_A^2|\Psi_{AB}\rangle - \langle\Psi_{AB}|\hat{G}_A|\Psi_{AB}\rangle^2) \\ &= 4(\text{Tr}_A[\hat{G}_A^2\hat{\rho}_A] - \text{Tr}_A[\hat{G}_A\hat{\rho}_A]^2) \\ &\equiv 4\text{Var}(\hat{G}_A)_{\hat{\rho}_A}. \end{aligned} \quad (5)$$

Consequently, for a purification of $\hat{\rho}_A$ the QFI only depends on the variance in \hat{G}_A of $\hat{\rho}_A$ [41,49]. Our naïve strategy of preparing a state with large $\text{Var}(\hat{G}_A)$ *irrespective of its purity* is now an excellent approach. Indeed, in this situation the states in cases I–III are now also arranged in order of increasing QFI, with cases II and III attaining a QFI of $N(N+1)/2$ and N^2 , respectively. It is interesting to note that the QFI for case III is the maximum allowable for N particles in SU(2) [50] and is usually obtained via the difficult to generate spin-cat state, which is a macroscopic superposition, rather than a classical mixture, of spin coherent states. Note also that \mathcal{F}_{AB} is independent of any particular purification, and convexity implies that $\mathcal{F}_{AB} \geq \mathcal{F}_A$. That is, in principle any purification of $\hat{\rho}_A$ is capable of achieving sensitivities at least as good as, and usually much better than, $\hat{\rho}_A$ itself.

Quantum metrology with purifications is not simply a mathematical “trick”; physically, a purification corresponds to entangling the probe system A with some auxiliary system B, and permitting measurements on both systems [51]. Therefore, the practical utility of our proposal depends crucially on the existence of an entangling Hamiltonian that can prepare $\hat{\rho}_A$ in a state with large $\text{Var}(\hat{G}_A)_{\hat{\rho}_A}$.

For the three cases described by Eqs. (2) and (3), a purification of $\hat{\rho}_A$ can be written as

$$|\Psi_{AB}\rangle = \sum_{m=-j}^j c_m |j, m\rangle \otimes |B_m\rangle, \quad (6)$$

with case I corresponding to $\langle B_m|B_n\rangle = 1$, case II corresponding to $\langle B_m|B_n\rangle = \delta_{n,m}$, and case III corresponding to $\langle B_m|B_n\rangle = 1(0)$ for $|n-m|$ even (odd). In the following section, we present a simple scheme that converts a shot-noise limited spin coherent state [such as case I] to the enhanced QFI purifications of cases II and III.

III. EXAMPLE ENTANGLING DYNAMICS LEADING TO INCREASED QFI

Consider again the N -boson probe state (system A) input into a MZ interferometer (i.e., $\hat{G}_A = \hat{J}_y$). The QFI for a purification of $\hat{\rho}_A$ can be written as

$$\mathcal{F}_{AB} = 4(\langle \hat{J}_y^2 \rangle - \langle \hat{J}_y \rangle^2) = \mathcal{F}_0 + \mathcal{F}_1 + \mathcal{F}_2, \quad (7)$$

with

$$\mathcal{F}_0 = \frac{N}{2}(N+2) - 2\langle \hat{J}_z^2 \rangle, \quad (8a)$$

$$\mathcal{F}_1 = -\langle i(\hat{J}_+ - \hat{J}_-) \rangle^2, \quad (8b)$$

$$\mathcal{F}_2 = -\langle \hat{J}_+^2 + \hat{J}_-^2 \rangle, \quad (8c)$$

where $\hat{J}_\pm = \hat{J}_x \pm i\hat{J}_y$. Note that $\mathcal{F}_0, \mathcal{F}_1$, and \mathcal{F}_2 depend only on the matrix elements of $\hat{\rho}_A$ with $|n-m|$ equal to 0, 1, and 2, in the \hat{J}_z basis; writing \mathcal{F}_{AB} in this form is very convenient for what follows.

Before the interferometer, we assume the probe is coupled to some auxiliary system B via the Hamiltonian

$$\hat{\mathcal{H}}_{AB} = \hbar g \hat{J}_z \hat{\mathcal{H}}_B. \quad (9)$$

When system B is a photon field and $\hat{\mathcal{H}}_B$ is proportional to the number of photons in the field, then $\hat{\mathcal{H}}_{AB}$ describes the weak probing of the population difference of an ensemble of two-level atoms with far-detuned light [37,52–61], or dispersive coupling between a microwave cavity and a superconducting qubit [62–64]. We will explore this specific case shortly; however, for now we keep $\hat{\mathcal{H}}_B$ completely general. If the initial system state is a product state $|\Psi_{AB}(0)\rangle = |\Psi_A\rangle \otimes |\Psi_B\rangle$, after some evolution time the state of the system will be given by Eq. (6) with $c_m = \langle m|\Psi_A\rangle$ and $|B_m\rangle = \exp(-imgt\hat{\mathcal{H}}_B)|\Psi_B\rangle$. The reduced density operator of system A is then

$$\hat{\rho}_A = \sum_{n,m=-j}^j c_n c_m^* \mathcal{C}_{n-m} |j, n\rangle \langle j, m|, \quad (10)$$

where the coherence of system A is determined via

$$\mathcal{C}_{n-m} = \langle B_m|B_n\rangle = \langle \Psi_B|e^{-i(n-m)gt\hat{\mathcal{H}}_B}|\Psi_B\rangle. \quad (11)$$

When $\mathcal{C}_{n-m} = 1$, the system remains separable and system A is a pure state, whereas if $\mathcal{C}_{n-m} = \delta_{n,m}$ then $\hat{\rho}_A$ is an incoherent mixture of Dicke states.

Using Eq. (10), $\mathcal{F}_0, \mathcal{F}_1$, and \mathcal{F}_2 can be written as

$$\mathcal{F}_0 = \frac{N}{2}(N+2) - 2\langle \hat{J}_z^2 \rangle, \quad (12a)$$

$$\mathcal{F}_1 = -\langle i(\mathcal{C}_1 \hat{J}_+ - \mathcal{C}_1^* \hat{J}_-) \rangle^2, \quad (12b)$$

$$\mathcal{F}_2 = -(\mathcal{C}_2 \langle \hat{J}_+^2 \rangle + \mathcal{C}_2^* \langle \hat{J}_-^2 \rangle), \quad (12c)$$

where the above expectation values are calculated with respect to $|\Psi_A\rangle$. The effect of the entanglement between systems A and B is entirely encoded in the coherences \mathcal{C}_1 and \mathcal{C}_2 ; coherences greater than second order do not affect the QFI.

Let us consider the effect on the QFI of each term in Eq. (7). \mathcal{F}_0 is independent of the entanglement between systems A and B , and will be of order $N^2/2$ if $|\Psi_A\rangle$ has $\langle \hat{J}_z^2 \rangle \sim N$ (e.g., the spin coherent state $|\alpha(\pi/2, \varphi)\rangle$ has $\langle \hat{J}_z^2 \rangle = N/4$). This suggests that a sufficient condition for Heisenberg scaling is $\mathcal{F}_1 \sim \mathcal{F}_2 \sim \langle \hat{J}_z^2 \rangle \sim N$. In fact, since $\mathcal{F}_1 \leq 0$, the maximum QFI state must necessarily have $\mathcal{C}_1 = 0$. In contrast, \mathcal{F}_2 can be positive or negative, in which case a state with $\mathcal{C}_2 = 0$ and another state with $\mathcal{C}_2 = 1$ and $\mathcal{F}_2 \sim +N^2/2$ might both be capable of (near) Heisenberg-limited metrology. We consider examples of both states below.

A. Case II: Example dynamics yielding $\mathcal{F}_{AB} \simeq N^2/2$

To concretely illustrate the increased QFI a purification of $\hat{\rho}_A$ can provide, we assume system B is a single bosonic mode, described by annihilation operator \hat{b} , and take $\hat{\mathcal{H}}_B = \hat{b}^\dagger \hat{b}$ such that

$$\hat{\mathcal{H}}_{AB} = \hbar g \hat{J}_z \hat{b}^\dagger \hat{b}. \quad (13)$$

If the initial state of system B is a Glauber coherent state $|\beta\rangle$ [65], then the coherences described by Eq. (11) simplify to

$$\mathcal{C}_{n-m} = \exp[-|\beta|^2(1 - e^{-i(n-m)gt})]. \quad (14)$$

$|\mathcal{C}_{n-m}|^2$ decays on a time scale $gt \sim [(n-m)|\beta|^2]^{-1}$. Although the nonorthogonality of $|\beta|\beta e^{i\theta}\rangle$ ensures that \mathcal{C}_{n-m} never actually reaches zero, it becomes very small for even modest values of $|\beta|^2$.

If the initial condition of system A is $|\Psi_A\rangle = |\alpha(\theta, \phi)\rangle$, then \mathcal{F}_{AB} has the simple analytic form given by (see Appendix)

$$\mathcal{F}_0 = N \left(1 + \frac{(N-1)}{2} \sin^2 \theta \right), \quad (15a)$$

$$\mathcal{F}_1 = -N^2 \sin^2 \theta \sin^2(|\beta|^2 \sin(gt) + \phi) e^{-4|\beta|^2 \sin^2(gt/2)}, \quad (15b)$$

$$\mathcal{F}_2 = \frac{N(N-N)}{2} \sin^2 \theta \cos(|\beta|^2 \sin(2gt) + 2\phi) e^{-2|\beta|^2 \sin^2(gt)}. \quad (15c)$$

In contrast, calculating \mathcal{F}_A via Eq. (1) requires the diagonalization of $\hat{\rho}_A$, which must be performed numerically.

We first demonstrate the effect of vanishing first- and second-order coherence on \mathcal{F}_{AB} by preparing system A in the maximal \hat{J}_x eigenstate, $|\alpha(\pi/2, 0)\rangle$, with $N = 100$, and a Glauber coherent state for system B with average particle number $|\beta|^2 = 500$. The initial state for system A is precisely case I [see Eq. (3a)] and has a QFI of N . As shown in Fig. 3,




FIG. 3. (Color online) Time snapshots of the Husimi- \mathcal{Q} function and \hat{J}_y projection illustrating the evolution of a maximal \hat{J}_x eigenstate under entangling Hamiltonian Eq. (13). The snapshots were chosen to correspond to times when the rotation around the J_z axis is such that $\langle \hat{J}_y \rangle = 0$, which roughly corresponds to the local maxima of \mathcal{F}_{AB} in Fig. 4(c). (Parameters: $N = 100, |\beta|^2 = 500$.)

under the evolution of Eq. (13), $\hat{\rho}_A$ tends towards an incoherent mixture of Dicke states [case II], with the corresponding broadening of the $P(J_y)$ distribution.

Figure 4(a) shows that both coherences \mathcal{C}_1 and \mathcal{C}_2 rapidly approach zero, which causes \mathcal{F}_1 and \mathcal{F}_2 to vanish [see Fig. 4(b)]. Consequently, \mathcal{F}_{AB} approaches $\mathcal{F}_0 = N(N+1)/2$, which allows a phase sensitivity of approximately $\sqrt{2} \times$ Heisenberg limit [see Fig. 4(c)]. In contrast, the effect of the mixing causes the QFI of $\hat{\rho}_A$ itself to decrease from N to $\mathcal{F}_A = N/2$, with $\mathcal{F}_A \leq N$ for all t . This remains true even if \hat{G}_A is rotated to lie in an arbitrary direction on the Bloch sphere.

The oscillations in \mathcal{F}_1 and \mathcal{F}_2 (and consequently \mathcal{F}_A and \mathcal{F}_{AB}) before the plateau are due to the complex rotation of \mathcal{C}_1 and \mathcal{C}_2 , which causes rotations of $\hat{\rho}_A$ around the J_z axis before being overwhelmed by the overall decay in magnitude. Furthermore, although the purity of the state also




FIG. 4. (Color online) Evolution of a maximal \hat{J}_x eigenstate under entangling Hamiltonian Eq. (13). (a) Coherences $|\mathcal{C}_1|^2$ (solid line), $|\mathcal{C}_2|^2$ (dashed line), and purity γ (dot-dashed line). (b) Three components of \mathcal{F}_{AB} [see Eq. (7)]: \mathcal{F}_0 (dot-dashed line), \mathcal{F}_1 (solid line), and \mathcal{F}_2 (dashed line). (c) QFI for $\hat{\rho}_A, \mathcal{F}_A$ (dashed line) and a purification of $\hat{\rho}_A, \mathcal{F}_{AB}$ (solid line). For reference, we have included N (dotted line) and $\mathcal{F}_0 = N(N+1)/2 \approx N^2/2$ (dot-dashed line), which correspond to phase sensitivities at the SQL and $\sqrt{2} \times$ Heisenberg limit, respectively. (Parameters: $N = 100, |\beta|^2 = 500$.)




FIG. 5. (Color online) Husimi- \mathcal{Q} function and \hat{J}_y projection for an initial state $|\Psi_{AB}(0)\rangle = |\alpha(\pi/2, \pi/2)\rangle \otimes |\beta\rangle$ under the evolution of Eq. (13) for different values of gt . The \mathcal{Q} function is symmetric about reflection of the J_y axis, resulting in part of the function being hidden from view on the reverse side of the sphere. (Parameters: $N = 100, |\beta|^2 = 500$.)

decays, it never vanishes, thereby illustrating that it is not the entanglement *per se* that is causing the QFI enhancement for a purification of $\hat{\rho}_A$.

B. Case III: Example dynamics yielding $\mathcal{F}_{AB} = N^2$

At $gt = \pi$, there is a revival in $|\mathcal{C}_n|^2$ for n even, but not for n odd. Figure 5 shows the Husimi- \mathcal{Q} function under the evolution of Eq. (13) for times close to $gt = \pi$, when the initial state of system A is the maximal \hat{J}_y eigenstate $|\alpha(\pi/2, \pi/2)\rangle$.

The QFI is initially zero, but the decay of \mathcal{C}_1 and \mathcal{C}_2 rapidly increases to $\mathcal{F}_{AB} = \mathcal{F}_0 = N(N+1)/2$ as in the previous example. As $gt \rightarrow \pi$, the revival of $|\mathcal{C}_2|^2$ causes \mathcal{F}_{AB} to briefly increase to N^2 (see Fig. 6). This is the Heisenberg limit, which is the QFI of a (pure) spin-cat state and the maximum QFI for $SU(2)$ [50]. At $gt = \pi, \hat{\rho}_A$ is identical to a classical mixture of $|\alpha(\pi/2, \pi/2)\rangle$ and $|\alpha(\pi/2, -\pi/2)\rangle$; however, its \mathcal{Q} function is similar to that of a spin-cat state, and purifications of it behave




FIG. 6. (Color online) Evolution of a maximal \hat{J}_y eigenstate near $gt = \pi$ under entangling Hamiltonian Eq. (13). (a) $|\mathcal{C}_1|^2$ (solid line), $|\mathcal{C}_2|^2$ (dashed line), and γ (dot-dashed line). (b) \mathcal{F}_0 (dot-dashed line), \mathcal{F}_1 (solid line), and \mathcal{F}_2 (dashed line). (c) \mathcal{F}_A (dashed line) and \mathcal{F}_{AB} (solid line). For comparison, we have included $\mathcal{F}_0 = N(N+1)/2 \approx N^2/2$ and the Heisenberg limit N^2 (dot-dashed lines); $\mathcal{F}_A \leq N$ for all t . (Parameters: $N = 100, |\beta|^2 = 500$.)

as a spin-cat state for metrological purposes. For these reasons, we call this state a *pseudo-spin-cat state*.

C. Example dynamics for particle-exchange Hamiltonian

In the previous two examples the \hat{J}_z projection was a conserved quantity, so any entanglement between systems A and B can only degrade the coherence in the \hat{J}_z basis of system A (ultimately resulting in an enhanced QFI). The situation is more complicated when considering a Hamiltonian that does not conserve the \hat{J}_z projection, such as when a spin flip in system A is correlated with the creation or annihilation of a quantum in system B . Here, we encounter scenarios where the interaction can either create or destroy coherences in the \hat{J}_z basis of system A , and although a significant QFI enhancement is still possible, it depends upon the initial state of system B .

As a concrete illustration, consider the particle-exchange Hamiltonian

$$\hat{\mathcal{H}}_{\pm} = \hbar g (\hat{J}_{\pm} \hat{b}^{\dagger} + \hat{J}_{\mp} \hat{b}), \quad (16)$$

and assume that system A is initially prepared in the maximal \hat{J}_z eigenstate $|\Psi_A\rangle = |\alpha(0,0)\rangle = |j,j\rangle$ (note this has $\text{Var}(\hat{J}_y) = N/4$, and therefore a QFI of N). Then $\hat{\mathcal{H}}_-$ and $\hat{\mathcal{H}}_+$ physically correspond to Raman superradiance [38,66] and quantum state transfer [34–37,39,40] processes, respectively. After some period of evolution, the combined state of systems $A \otimes B$ takes the form of Eq. (6).

First, consider the case when the initial state for system B is a large amplitude coherent state (i.e., $|\Psi_B\rangle = |\beta\rangle$). Here the addition or removal of a quantum to or from system B has a minimal effect on the state and the system remains approximately separable, since $|\langle B_n | B_m \rangle|^2 \approx 1$. It is therefore reasonable to make the undepleted pump approximation $\hat{b} \rightarrow \beta$, such that $\hat{\mathcal{H}}_{\pm} \rightarrow \hbar g \beta \hat{J}_x$ (assuming β is real). Hence, the effect of the interaction is simply a rotation around the J_x axis, which can create coherence in the \hat{J}_z basis, and so $\mathcal{F}_A = \mathcal{F}_{AB} \leq N$ for all time.

In the opposite limit where the initial state of system B is a Fock state with N_B particles, $|\Psi_B\rangle = |N_B\rangle$, then

$$|B_m\rangle = |N_B \pm (m - j)\rangle, \quad (17)$$

and $\langle B_m | B_n \rangle = \delta_{n,m}$. This ensures that the first- and second-order coherences vanish, and $\mathcal{F}_1 = \mathcal{F}_2 = 0$ for all time. That is, as illustrated in Fig. 7, the state moves towards the equator




FIG. 7. (Color online) Husimi- \mathcal{Q} function and \hat{J}_y projection for an initial state $|\Psi_{AB}(0)\rangle = |\alpha(0,0)\rangle \otimes |N_B\rangle$ under the evolution of $\hat{\mathcal{H}}_-$ for different values of gt . (Parameters: $N = 100, N_B = 20$).




FIG. 8. (Color online) \mathcal{F}_A (dashed line) and \mathcal{F}_{AB} (solid line) for an initial state $|\Psi_{AB}(0)\rangle = |\alpha(0,0)\rangle \otimes |N_B\rangle$ under the evolution of $\hat{\mathcal{H}}_-$. We have indicated N and $N^2/2$ with black dotted lines for comparison. (Parameters: $N = 100, N_B = 20$).

and ultimately evolves to an incoherent Dicke mixture [i.e., case II]. As described in Sec. III A, and shown in Fig. 8, the QFI increases to a maximum of approximately $\mathcal{F}_{AB} \approx N^2/2$. Although setting $N_B = 0$ (i.e., a vacuum state) leads to a larger variance in \hat{J}_z , \mathcal{F}_{AB} still reaches approximately 70% of $N^2/2$.

We therefore see that for the Hamiltonian (16), a large QFI enhancement is achieved provided the initial state $|\Psi_B\rangle$ has small number fluctuations. Compare this to the Hamiltonian (13), where the choice $|\Psi_B\rangle = |N_B\rangle$ leads to no entanglement between systems A and B , while in contrast an initial state with small phase fluctuations (and therefore large number fluctuations), such as a coherent state, causes rapid decoherence in $\hat{\rho}_A$.

IV. OPTIMAL MEASUREMENT SCHEMES

Although the QFI determines the optimum sensitivity for a given initial state, it is silent on the question of how to achieve this optimum. It is therefore important to identify (a) which measurements to make on each system and (b) a method of combining the outcomes of these measurements—which we refer to as a *measurement signal* (\hat{S})—that saturates the QCRB. We do this below for purifications of the incoherent Dicke mixture (case II) and the pseudo-spin-cat state (case III).

A. Optimal measurements for incoherent Dicke mixture (case II)

It is worthwhile briefly recounting the optimal estimation procedure for a symmetric Dicke state $|j,0\rangle$ input into a MZ interferometer. A MZ interferometer rotates \hat{J}_z according to $\hat{J}_z(\phi) = \hat{U}_\phi^\dagger \hat{J}_z \hat{U}_\phi = \cos \phi \hat{J}_z - \sin \phi \hat{J}_x$. Since symmetric Dicke states satisfy $\langle \hat{J}_x \rangle = \langle \hat{J}_z \rangle = \langle \hat{J}_z^2 \rangle = 0$ and $\langle \hat{J}_x^2 \rangle = N(N+2)/8$, it is clear that the *fluctuations* in $\hat{J}_z(\phi)$ contain the phase information, and therefore the quantity $\hat{S} = [\hat{J}_z(\phi)]^2$ oscillates between 0 and $N(N+2)/8$. It can be shown that at the operating point $\phi \rightarrow 0$, $\text{Var}(\hat{S}) \rightarrow 0$, and the quantity $(\Delta\phi)^2 \rightarrow \text{Var}(\hat{S})/(\partial_\phi \langle \hat{S} \rangle)^2 = 1/\mathcal{F}_A$, and therefore the signal \hat{S} saturates the QCRB [3,67,68].

For an incoherent Dicke mixture, we have $\langle \hat{J}_x \rangle = \langle \hat{J}_y \rangle = \langle \hat{J}_z \rangle = 0$, and $\langle \hat{J}_x^2 \rangle = N(N+1)/8$. Unfortunately, the nonzero variance in \hat{J}_z (i.e., $\langle \hat{J}_z^2 \rangle = N/4$) implies that $\text{Var}(\hat{S}) \gg 0$ for all ϕ , and the signal no longer saturates the QCRB. However, since the states $|B_m\rangle$ in the purification Eq. (6) are orthonormal,

a projective measurement of some system B operator diagonal in the $|B_m\rangle$ basis projects system A into a \hat{J}_z eigenstate (i.e., a Dicke state). That is, these measurement outcomes on system B are correlated with \hat{J}_z measurement outcomes on system A . Therefore, subtracting both measurements yields a quantity with very little quantum noise.

More precisely, if we can construct an operator \hat{S}_B on system B that is correlated with \hat{J}_z measurements on system A (i.e., $\hat{S}_B|\Psi_{AB}\rangle = \hat{J}_z|\Psi_{AB}\rangle$), then we can construct the quantity $\hat{S}_0 = \hat{J}_z - \hat{S}_B$ which has the property $\langle \hat{S}_0 \rangle = \langle \hat{S}_0^2 \rangle = 0$. This motivates the signal choice

$$\hat{S} = (\hat{U}_\phi^\dagger \hat{S}_0 \hat{U}_\phi)^2 = (\cos \phi \hat{J}_z - \sin \phi \hat{J}_x - \hat{S}_B)^2. \quad (18)$$

Using $\hat{S}_B|\Psi_{AB}\rangle = \hat{J}_z|\Psi_{AB}\rangle$ and the fact that non- \hat{J}_z -conserving terms vanish due to the absence of off-diagonal terms in the \hat{J}_z representation of $\hat{\rho}_A$ (e.g., expectation values with an odd power of \hat{J}_x vanish), we can show that

$$\langle \hat{S} \rangle = \langle \hat{J}_z^2 \rangle (\cos \phi - 1)^2 + \langle \hat{J}_x^2 \rangle \sin^2 \phi, \quad (19a)$$

$$\begin{aligned} \langle \hat{S}^2 \rangle &= \langle \hat{J}_z^4 \rangle (\cos \phi - 1)^4 + \langle \hat{J}_x^4 \rangle \sin^4 \phi \\ &+ \langle \hat{J}_z^2 \hat{J}_x^2 + \hat{J}_x^2 \hat{J}_z^2 + 4\hat{J}_z \hat{J}_x \hat{J}_x \hat{J}_z \rangle \sin^2 \phi (\cos \phi - 1)^2 \\ &+ 2i \langle (\hat{J}_z \hat{J}_x \hat{J}_y - \hat{J}_y \hat{J}_x \hat{J}_z) \rangle \sin^2 \phi \cos \phi (\cos \phi - 1) \\ &+ \langle \hat{J}_y^2 \rangle \cos^2 \phi \sin^2 \phi. \end{aligned} \quad (19b)$$

Note that the above expectation values can be taken with respect to $|\Psi_{AB}\rangle$ or $\hat{\rho}_A$. The best sensitivity occurs at small displacements around $\phi = 0$. Taking the limit as $\phi \rightarrow 0$ and noting that $\langle \hat{J}_x^2 \rangle = \langle \hat{J}_y^2 \rangle$ gives

$$(\Delta \phi)^2 = \left. \frac{\text{Var}(\hat{S})}{(\partial_\phi \langle \hat{S} \rangle)^2} \right|_{\phi=0} = \frac{1}{4\langle \hat{J}_x^2 \rangle} = \frac{1}{4\langle \hat{J}_y^2 \rangle} = \frac{1}{\mathcal{F}_{AB}}. \quad (20)$$

This demonstrates that the signal Eq. (18) is optimal since it saturates the QCRB.

B. Optimal measurements for pseudo-spin-cat state (case III)

Pure spin-cat states have the maximum QFI possible for N particles in $SU(2)$, are eigenstates of the parity operator, and indeed parity measurements saturate the QCRB [69]. Pseudo-spin-cat states (case III) also have maximal QFI, and since $\langle B_n | B_m \rangle = 1(0)$ for $|n - m|$ even (odd), a projective measurement of system B yields no information other than the parity of the \hat{J}_z projection. This suggests that a measurement of parity could be optimal.

In analogy with case II, our aim is to construct an operator \hat{S}_0 where the correlations between systems A and B lead to a reduction in $\text{Var}(\hat{S}_0)$ and the system mimics a pure spin-cat state. Introducing the quantity

$$\hat{S}_0 = \hat{\Pi}_A \hat{S}_B \equiv \hat{\Pi}_A \hat{\Pi}_B, \quad (21)$$

where $\hat{\Pi}_{A(B)}$ is the parity operator for system $A(B)$, defined by $\hat{\Pi}_A |j, m\rangle = (-1)^m |j, m\rangle$ and $\hat{\Pi}_B |B_m\rangle = (-1)^m |B_m\rangle$, we see that pseudo-spin-cat states satisfy $\hat{S}_0 |\Psi_{AB}\rangle = |\Psi_{AB}\rangle$, and therefore $\text{Var}(\hat{S}_0) = 0$. This motivates the signal choice $\hat{S} = \hat{U}_\phi^\dagger \hat{S}_0 \hat{U}_\phi$.

To calculate the sensitivity, we need to compute $\langle \hat{S} \rangle$ and $\langle \hat{S}^2 \rangle$. Trivially, $\langle \hat{S}^2 \rangle = 1$ for all states. For $\phi \ll 1$, expanding

\hat{U}_ϕ to second order in ϕ gives

$$\begin{aligned} \langle \hat{S} \rangle &\approx \langle (1 + i\phi \hat{J}_y - \frac{1}{2}\phi^2 \hat{J}_y^2) \hat{S}_0 (1 - i\phi \hat{J}_y - \frac{1}{2}\phi^2 \hat{J}_y^2) \rangle \\ &= 1 + i\phi (\langle \hat{J}_y \hat{S}_0 \rangle - \langle \hat{S}_0 \hat{J}_y \rangle) \\ &+ \phi^2 [\langle \hat{J}_y \hat{S}_0 \hat{J}_y \rangle - \frac{1}{2} (\langle \hat{J}_y^2 \hat{S}_0 \rangle + \langle \hat{S}_0 \hat{J}_y^2 \rangle)] + \mathcal{O}(\phi^3), \end{aligned} \quad (22)$$

The relation $\langle B_n | B_{n\pm 1} \rangle = 0$ ensures that terms linear in \hat{J}_y go to zero:

$$\begin{aligned} \langle \hat{J}_+ \rangle &= \sum_{m,n} c_m c_n^* \langle j, n | \hat{J}_+ | j, m \rangle \langle B_n | B_m \rangle \\ &\propto \sum_{n,m} c_m c_n^* \delta_{n,m+1} \langle B_n | B_m \rangle \\ &= \sum_m c_m c_{m+1}^* \langle B_{m+1} | B_m \rangle = 0. \end{aligned} \quad (23)$$

However, unlike case II, the condition $\langle B_n | B_{n\pm 2} \rangle = 1$ preserves terms such as $\langle \hat{J}_+^2 \rangle$. Noting that \hat{J}_y flips the parity of any state in subsystem A but not subsystem B :

$$\hat{\Pi}_A \hat{J}_y |\Psi_{AB}\rangle = -\hat{J}_y \hat{\Pi}_A |\Psi_{AB}\rangle, \quad (24a)$$

$$\hat{\Pi}_B \hat{J}_y |\Psi_{AB}\rangle = \hat{J}_y \hat{\Pi}_B |\Psi_{AB}\rangle, \quad (24b)$$

and using $\hat{S}_0 |\Psi_{AB}\rangle = |\Psi_{AB}\rangle$ gives

$$\langle \hat{J}_y \hat{S}_0 \hat{J}_y \rangle = -\langle \hat{S}_0 \hat{J}_y^2 \rangle = -\langle \hat{J}_y^2 \hat{S}_0 \rangle = -\langle \hat{J}_y^2 \rangle. \quad (25)$$

Therefore

$$\langle \hat{S} \rangle = 1 - 2\phi^2 \langle \hat{J}_y^2 \rangle + \mathcal{O}(\phi^3). \quad (26)$$

Since $\hat{S}_0^2 = 1$ implies that $\hat{S}^2 = 1$, we obtain

$$\text{Var}(\hat{S}) = 4\phi^2 \langle \hat{J}_y^2 \rangle + \mathcal{O}(\phi^4), \quad (27)$$

and consequently

$$(\Delta \phi)^2 = \frac{\text{Var}(\hat{S})}{(\partial_\phi \langle \hat{S} \rangle)^2} = \frac{4\phi^2 \langle \hat{J}_y^2 \rangle}{16\phi^2 \langle \hat{J}_y^2 \rangle^2} = \frac{1}{4\langle \hat{J}_y^2 \rangle} = \frac{1}{\mathcal{F}_{AB}}. \quad (28)$$

This demonstrates that the signal saturates the QCRB and is therefore optimal.

The optimal estimation schemes presented in Secs. IV A and IV B illustrate a somewhat counterintuitive fact: Although the optimal measurement of system B for a pseudo-spin-cat state provides *less* information about system A than for an incoherent Dicke mixture, the pseudo-spin-cat state yields the better (in fact *best*) sensitivity.

C. System B observables that approximate optimal measurements

We now turn to the explicit construction of physical observables that approximate \hat{S}_B . In general, the choice of \hat{S}_B depends upon the specific purification of $\hat{\rho}_A$. Physically, the initial state of system B and the entangling Hamiltonian matter. However, there is no guarantee that \hat{S}_B exists, and if it does there is no guarantee that a measurement of this observable can be made in practice. Nevertheless, as we show below, it may be possible to make a measurement of an observable that *approximates* \hat{S}_B and can therefore give near-optimal sensitivities.

I. Case II

To begin, consider the situation in Sec. III A: the evolution of the state $|\alpha(\pi/2, 0)\rangle \otimes |\beta\rangle$ under the Hamiltonian (13). We require $\hat{S}_B |\Psi_{AB}\rangle = \hat{J}_z |\Psi_{AB}\rangle$. After some evolution time t ,

$$|\Psi_{AB}\rangle = \sum_{m=-j}^j c_m |j, m\rangle |\beta e^{-imgt}\rangle. \quad (29)$$

Clearly, the *phase* of the coherent state is correlated with the \hat{J}_z projection of system B . This can be extracted via a homodyne measurement of the phase quadrature $\hat{Y}_B = i(\hat{b} - \hat{b}^\dagger)$ [70]. In fact, provided $mg t \ll 1$, phase quadrature measurements of $|\beta \exp(-imgt)\rangle$ are linearly proportional to the \hat{J}_z projection:

$$\langle \beta e^{-imgt} | \hat{Y}_B | \beta e^{-imgt} \rangle = 2\beta \sin(mgt) \approx 2\beta mgt, \quad (30)$$

where without loss of generality we have taken β to be real and positive. Consequently, the scaled phase quadrature

$$\hat{S}_B = \frac{\hat{Y}_B}{2\beta gt} \quad (31)$$

satisfies

$$\langle \Psi_{AB} | (\hat{J}_z - \hat{S}_B) | \Psi_{AB} \rangle \approx 0, \quad (32a)$$

$$\langle \Psi_{AB} | (\hat{J}_z - \hat{S}_B)^2 | \Psi_{AB} \rangle \approx \frac{1}{(2\beta gt)^2}, \quad (32b)$$

and so the fluctuations in $(\hat{J}_z - \hat{S}_B)$ become arbitrarily small (and \hat{S}_B becomes perfectly correlated with \hat{J}_z) as $(\beta gt)^2$ becomes large. This suggests that Eqs. (18) and (31) should be a good approximation to an optimal measurement signal.

More precisely, assume that

$$\frac{1}{\beta} \ll \frac{Ngt}{2} \ll 1. \quad (33)$$

The first inequality ensures that $\langle \beta e^{-imgt} | \beta e^{-imgt} \rangle \approx \delta_{n,m}$ and so $\hat{\rho}_A$ is approximately an incoherent Dicke mixture, while the second inequality implies that we are in the linearized regime where Eqs. (30) and (32) hold. Then the signal

$$\hat{S}_{\text{approx}} = \left(\hat{J}_z \cos \phi - \hat{J}_x \sin \phi - \frac{\hat{Y}_B}{2gt} \right)^2 \quad (34)$$

yields the sensitivity

$$(\Delta\phi)^2 \approx \frac{1}{(\partial_\phi \langle \hat{S} \rangle)^2} \left\{ \text{Var}(\hat{S}) - \frac{\sin^2(\phi/2)}{\beta^2} \langle \hat{J}_z^4 \rangle + \frac{2}{(2\beta gt)^4} + \frac{4}{(2\beta gt)^2} [(\cos \phi - 1)^2 \langle \hat{J}_z^2 \rangle + \sin^2 \phi \langle \hat{J}_x^2 \rangle] \right\}, \quad (35)$$

where $\text{Var}(\hat{S})$ and $\partial_\phi \langle \hat{S} \rangle$ are given by the expectations (19) of the optimal signal Eq. (18). Figure 9 shows Eq. (35) compared to an exact numeric calculation.

Condition (33) typically ensures that the term proportional to $\langle \hat{J}_z^4 \rangle$ is small in comparison to the term proportional to $1/(2\beta gt)^2$. We therefore see that our approximate signal \hat{S}_{approx} gives a sensitivity worse than the QCRB, and furthermore at an operating point $\phi \neq 0$. Nevertheless, $\Delta\phi$ approaches the QCRB at $\phi = 0$ as β^2 and $(2\beta gt)^2$ approach infinity. Therefore, for a sufficiently large βgt , we can achieve near-optimal




FIG. 9. (Color online) $\Delta\phi$ vs ϕ using Eq. (34) for a state of the form Eq. (29), with $N = 100$, and $gt = 10^{-2}$. The dot-dashed line is with $|\beta|^2 = 10^6$ ($\beta gt = 10$), and the solid line is for $|\beta|^2 = 10^4$ ($\beta gt = 1$). The dashed line shows the approximate expression for the sensitivity [Eq. (35)] for $|\beta|^2 = 10^4$. For $|\beta|^2 = 10^6$, the numerical calculation and Eq. (35) are identical. The upper and lower black dotted lines represent the standard quantum limit ($1/\sqrt{N}$) and $\sqrt{2}/N$, respectively. The divergence in $\Delta\phi$ close to $\phi = 0$ in both cases is due to the imperfect correlations between \hat{S}_B and \hat{J}_z leading to nonzero variance in \hat{S} . If the correlations were perfect and $\text{Var}(\hat{S})|_{\phi=0} = 0$, $\Delta\phi$ would reach exactly $1/\sqrt{\mathcal{F}_{AB}}$ at $\phi = 0$.

sensitivities close to $\phi = 0$. This is illustrated in Fig. 9. When $\beta gt = 10$, we find that $\Delta\phi$ is very close to the QCRB. In contrast, for $\beta gt = 1$, the imperfect correlations between \hat{J}_z and \hat{S}_B prevent the sensitivity from reaching the QCRB; nevertheless, the sensitivity is still below the SQL. Note that there is a slight deviation between Eq. (35) and the numerical calculation of the sensitivity using the state (29). This is due to terms neglected by our approximations, in particular, the nonlinear terms ignored by linearizations such as Eq. (30) and those neglected terms that arise due to the small (but strictly nonzero) off-diagonal elements of $\hat{\rho}_A$.

2. Case III

Now, consider the situation in Sec. III B: the evolution of the state $|\alpha(0, 0)\rangle \otimes |\beta\rangle$ under the Hamiltonian (13) that at $gt = \pi$ approximately results in a pseudo-spin-cat state.

In order to find an operator that approximates $\hat{S}_B = \hat{\Pi}_B$, we introduce the amplitude quadrature operator $\hat{X}_B = (\hat{b} + \hat{b}^\dagger)$, and notice that

$$\begin{aligned} \langle \beta e^{-im\pi} | \hat{X}_B | \beta e^{-im\pi} \rangle &= 2\beta(-1)^m \\ &= 2\beta \langle j, m | \hat{\Pi}_A | j, m \rangle. \end{aligned} \quad (36)$$

That is, amplitude quadrature measurements of $|\beta e^{-im\pi}\rangle$ are proportional to parity measurements on system B , which are directly correlated with parity measurements on \hat{J}_z eigenstates. Indeed, the quantity

$$\hat{S}_0 = \hat{\Pi}_A \frac{\hat{X}_B}{2\beta}, \quad (37)$$

has a variance $\text{Var}(\hat{S}_0) = 1/(2\beta)^2$ that becomes vanishingly small as the amplitude of the coherent state is increased. We therefore expect the signal

$$\hat{S}_{\text{approx}} = \hat{U}_\phi^\dagger \left(\hat{\Pi}_A \frac{\hat{X}_B}{2\beta} \right) \hat{U}_\phi \quad (38)$$




FIG. 10. (Color online) Phase sensitivity of the approximate signal Eq. (38) for a state of the form Eq. (29) at $gt = \pi$ (i.e., a pseudo-spin-cat state) with $N = 20$. The solid line and dashed line are for $|\beta|^2 = 30$ and $|\beta|^2 = 5$, respectively. The black dotted line indicates the Heisenberg limit $\Delta\phi = 1/N$ (which is the QCRB). Note that the vertical axis is a linear scale.

will be a good approximation to the optimal measurement $\hat{\mathcal{S}}$.

Figure 10 shows the sensitivity for a state of the form Eq. (29) at $gt = \pi$ with $N = 20$. When $|\beta|^2 = 30$, the sensitivity is very close to the Heisenberg limit, while for $|\beta|^2 = 5$ there is a slight degradation in the sensitivity due to imperfect correlations. In contrast to the approximate optimal measurement scheme for case II, which requires a large amplitude coherent state, here the signal (38) is almost optimal even for small amplitude coherent states. This is because $(\beta e^{-i\pi}|\beta\rangle) = \exp(-2|\beta|^2)$ is approximately zero even for modest values of β .

In situations where system A is an ensemble of atoms, and system B is an optical mode, it would be challenging to achieve the strong atom-light coupling regime required for $gt = \pi$. On the other hand, the choice of an initial coherent state for system B ensures that the sensitivity is reasonably insensitive to losses in system B . In particular, since particle loss from a coherent state acts only to reduce the state's amplitude, provided the coherent state remains sufficiently large *after* losses in order to satisfy the requirements for near-optimal measurements, near-Heisenberg-limited sensitivities should be obtainable.

3. Particle-exchange Hamiltonian

Finally, for completeness we include the optimal measurement scheme for the state attained after evolving the product state $|\alpha(0,0)\rangle \otimes |N_B\rangle$ under the Hamiltonian (16) (see Sec. III C). The optimal measurement signal is simply Eq. (18) with

$$\hat{\mathcal{S}}_B = \frac{N}{2} \pm (N_b - \hat{b}^\dagger \hat{b}). \quad (39)$$

This choice of $\hat{\mathcal{S}}_B$ can be constructed by counting the number of particles in system B , and it satisfies $\hat{\mathcal{S}}_B|\Psi_{AB}\rangle = \hat{J}_z|\Psi_{AB}\rangle$ as required. As *any* entangling Hamiltonian of the form

$$\hat{\mathcal{H}}_{\pm} = \sum_k \mathcal{A}_k (\hat{J}_{\pm} \hat{b}^\dagger + \hat{J}_{\mp} \hat{b})^k + \sum_{j,k} \mathcal{B}_{j,k} \hat{J}_z^k (\hat{b}^\dagger \hat{b})^j \quad (40)$$

will lead to a state of the form Eq. (6), with $|B_m\rangle$ given by Eq. (17) (assuming an initial state $|\Psi_A\rangle = |\alpha(0,0)\rangle, |\Psi_B\rangle = |N_B\rangle$), Eq. (39) also transfers to these systems.

V. DISCUSSION AND CONCLUSIONS

We have shown that purifications of mixed states represent an excellent resource for quantum metrology. In particular, we showed that if probe system A and auxiliary system B are entangled such that the first- and second-order coherences of system A vanish, then near-Heisenberg-limited sensitivities can be achieved provided measurements on both systems A and B are allowed. Although we focused on the situation where this entanglement is generated via a few specific Hamiltonians, our conclusions hold irrespective of the specific entanglement generation scheme.

It is important to note that when the QFI approaches the Heisenberg limit $\mathcal{F}_{AB} = N^2$ (in case III, for example), this is not the *true* Heisenberg limit, since N is only the number of particles in system A (which pass through the interferometer), rather than the *total* number of particles N_t in system A and system B . However, we have chosen to report the QFI in terms of N instead of N_t , since we envisage our protocol to be useful in situations where the number of particles in system A is by far the more valuable resource. For example, consider the case of inertial sensing with atom interferometry, where system A is atoms and system B is photons. Here the atoms are sensitive to an inertial phase shift, but it is difficult to arbitrarily increase the atomic flux. Using the techniques discussed above, an improved inertial measurement can be achieved at the cost of introducing some number of auxiliary photons, which is cheap compared to increasing the total number of atoms.

While preparing this paper, we also numerically examined the effect of decoherence on the sensitivity of our metrological schemes. In particular, we found that the effect of particle loss, spin flips, and phase diffusion on purifications of the pseudo-spin-cat state from Fig. 2 was identical to that of a pure spin-cat state.

Although these purified states are no more or less robust to decoherence than other nonclassical pure states, there are situations where they are easier to generate. The example we are most familiar with is atom interferometry, where atom-light entanglement and information recycling is more compatible with the requirements of high-precision atom interferometry than the preparation of nonclassical atomic states via interatomic interactions [39]. However, controlled interactions are routinely engineered between atoms and light [37,71–73], superconducting circuits and microwaves [74,75], light and mechanical systems [76], and ions and light [77–79]. Given that high-efficiency detection is available in all these systems [80–85], the application of our proposal to a range of metrological platforms is plausible in the near term.

Finally, we note that not all quantum systems are created equal; certain quantum information protocols, such as quantum error correction [86] and no-knowledge feedback [87], are better suited to some platforms than others. Our proposal allows an experimenter to both perform quantum-enhanced metrology and take advantage of any additional benefits a hybrid quantum system provides.

ACKNOWLEDGMENTS

We acknowledge useful discussions with Carlton Caves, Joel Corney, Jamie Fiess, Samuel Nolan, and Murray Olsen. This work was supported by Australian Research Council (ARC) Discovery Project No. DE130100575 and the ARC Centre of Excellence for Engineered Quantum Systems (Project No. CE110001013).

APPENDIX: DERIVATION OF EQS. (15)

Here we derive the QFI \mathcal{F}_{AB} for a MZ interferometer with the following entangled input:

$$|\Psi_{AB}(t)\rangle = e^{-igt\hat{J}_z\hat{N}_b}|\theta, \varphi\rangle \otimes |\beta\rangle, \quad (\text{A1})$$

where $\hat{N}_b = \hat{b}^\dagger\hat{b}$, system B is initially in a coherent state $|\beta\rangle$, and system A is initially in a spin coherent state $|\theta, \varphi\rangle$. Any spin coherent state can be defined by rotating the maximal Dicke state on the top pole of the Bloch sphere an angle θ about the J_y axis and an angle φ about the J_z axis:

$$|\theta, \varphi\rangle \equiv \hat{R}(\theta, \varphi)|j, j\rangle = e^{-i\varphi\hat{J}_z}e^{-i\theta\hat{J}_y}|j, j\rangle. \quad (\text{A2})$$

Recall that $j = N/2$, where N is the total number of system A particles.

The QFI is

$$\mathcal{F}_{AB} = 4\text{Var}(e^{igt\hat{J}_z\hat{N}_b}\hat{J}_ye^{-igt\hat{J}_z\hat{N}_b}), \quad (\text{A3})$$

where the expectations in the variance are taken with respect to the initial separable state, $|\theta, \varphi\rangle \otimes |\beta\rangle$.

By application of the Baker-Campbell-Hausdorff formula

$$\begin{aligned} e^{\hat{A}}\hat{B}e^{-\hat{A}} &= \hat{B} + [\hat{A}, \hat{B}] + \frac{1}{2!}[\hat{A}, [\hat{A}, \hat{B}]] \\ &+ \frac{1}{3!}[\hat{A}, [\hat{A}, [\hat{A}, \hat{B}]]] + \dots \end{aligned} \quad (\text{A4})$$

it can be shown that

$$e^{igt\hat{J}_z\hat{N}_b}\hat{J}_ye^{-igt\hat{J}_z\hat{N}_b} = \sin(gt\hat{N}_b)\hat{J}_x + \cos(gt\hat{N}_b)\hat{J}_y. \quad (\text{A5})$$

Therefore, since the initial state is separable, we obtain

$$\begin{aligned} \mathcal{F}_{AB} &= \langle \sin^2(gt\hat{N}_b) \rangle \langle \hat{J}_x^2 \rangle + \langle \cos^2(gt\hat{N}_b) \rangle \langle \hat{J}_y^2 \rangle \\ &+ \langle \sin(gt\hat{N}_b) \cos(gt\hat{N}_b) \rangle \langle \hat{J}_x\hat{J}_y + \hat{J}_y\hat{J}_x \rangle \\ &- (\langle \sin(gt\hat{N}_b) \rangle \langle \hat{J}_x \rangle + \langle \cos(gt\hat{N}_b) \rangle \langle \hat{J}_y \rangle)^2. \end{aligned} \quad (\text{A6})$$

The system A expectations are more easily computed by rotating the operators by $\hat{R}(\theta, \varphi)$ and then taking expectations with respect to the Dicke state $|j, j\rangle$. Specifically,

by virtue of

$$\hat{R}^\dagger(\theta, \varphi)\hat{J}_x\hat{R}(\theta, \varphi) = \cos\theta\cos\varphi\hat{J}_x + \cos\theta\sin\varphi\hat{J}_y - \sin\theta\hat{J}_z, \quad (\text{A7a})$$

$$\hat{R}^\dagger(\theta, \varphi)\hat{J}_y\hat{R}(\theta, \varphi) = -\sin\varphi\hat{J}_x + \cos\varphi\hat{J}_y, \quad (\text{A7b})$$

and the application of $\hat{J}_\pm = \hat{J}_x \pm i\hat{J}_y$ with

$$\hat{J}_\pm|j, m\rangle = \sqrt{j(j+1) - (m \pm 1)^2}|j, m \pm 1\rangle, \quad (\text{A8})$$

we obtain

$$\langle \hat{J}_x \rangle = j \sin\theta \cos\varphi, \quad (\text{A9a})$$

$$\langle \hat{J}_y \rangle = j \sin\theta \sin\varphi, \quad (\text{A9b})$$

$$\langle \hat{J}_x^2 \rangle = \frac{j}{2}(1 + (2j-1)\sin^2\theta\cos^2\varphi), \quad (\text{A9c})$$

$$\langle \hat{J}_y^2 \rangle = \frac{j}{2}(1 + (2j-1)\sin^2\theta\sin^2\varphi), \quad (\text{A9d})$$

$$\langle \hat{J}_x\hat{J}_y + \hat{J}_y\hat{J}_x \rangle = j(2j-1)\sin^2\theta\sin\varphi\cos\varphi. \quad (\text{A9e})$$

With some simplification this gives

$$\begin{aligned} \mathcal{F}_{AB} &= 2j(1 + \sin^2\theta[(2j-1)(\sin^2(gt\hat{N}_b + \varphi)) \\ &- 2j\langle \sin(gt\hat{N}_b + \varphi) \rangle^2]). \end{aligned} \quad (\text{A10})$$

Incidentally, by setting $t = 0$ we can see that the QFI for a spin coherent state input never exceeds the standard quantum limit:

$$\mathcal{F}[\hat{J}_y, |\theta, \varphi\rangle] = 2j(1 - \sin^2\theta\sin^2\varphi) \leq N. \quad (\text{A11})$$

In order to compute the system B expectations, note that

$$\langle \sin(gt\hat{N}_b + \varphi) \rangle = -\frac{i}{2}(\langle e^{i(gt\hat{N}_b + \varphi)} \rangle - \langle e^{-i(gt\hat{N}_b + \varphi)} \rangle) \quad (\text{A12a})$$

$$\langle \sin^2(gt\hat{N}_b + \varphi) \rangle = \frac{2 - \langle e^{2i(gt\hat{N}_b + \varphi)} \rangle - \langle e^{-2i(gt\hat{N}_b + \varphi)} \rangle}{4}. \quad (\text{A12b})$$

Furthermore, for any m ,

$$\begin{aligned} \langle e^{im(gt\hat{N}_b + \varphi)} \rangle &= e^{im\varphi} \langle \beta | \beta e^{imgt} \rangle \\ &= \exp[-|\beta|^2(1 - e^{imgt}) + 2m\varphi]. \end{aligned} \quad (\text{A13})$$

Substituting Eqs. (A12) and (A13) into Eq. (A10) gives

$$\mathcal{F}_{AB} = \mathcal{F}_0 + \mathcal{F}_1 + \mathcal{F}_2, \quad (\text{A14})$$

with the expressions for $\mathcal{F}_0, \mathcal{F}_1$, and \mathcal{F}_2 listed in Eqs. (15).

- [1] M. J. Holland and K. Burnett, Interferometric Detection of Optical Phase Shifts at the Heisenberg Limit, *Phys. Rev. Lett.* **71**, 1355 (1993).
 [2] V. Giovannetti, S. Lloyd, and L. Maccone, Quantum Metrology, *Phys. Rev. Lett.* **96**, 010401 (2006).

- [3] B. Lücke, M. Scherer, J. Kruse, L. Pezzé, F. Deuretzbacher, P. Hyllus, O. Topic, J. Peise, W. Ertmer, J. Arlt, L. Santos, A. Smerzi, and C. Klempt, Twin matter waves for interferometry beyond the classical limit, *Science* **334**, 773 (2011).

- [4] B. C. Sanders and G. J. Milburn, Optimal Quantum Measurements for Phase Estimation, *Phys. Rev. Lett.* **75**, 2944 (1995).
- [5] L. Pezzé and A. Smerzi, Ultrasensitive Two-Mode Interferometry with Single-Mode Number Squeezing, *Phys. Rev. Lett.* **110**, 163604 (2013).
- [6] R. Demkowicz-Dobrzanski, J. Kolodynski, and M. Guta, The elusive Heisenberg limit in quantum-enhanced metrology, *Nat. Commun.* **3**, 1063 (2012).
- [7] D. Leibfried, E. Knill, S. Seidelin, J. Britton, R. B. Blakestad, J. Chiaverini, D. B. Hume, W. M. Itano, J. D. Jost, C. Langer, R. Ozeri, R. Reichle, and D. J. Wineland, Creation of a six-atom “Schrodinger cat” state, *Nature (London)* **438**, 639 (2005).
- [8] W. Wieczorek, R. Krischek, N. Kiesel, P. Michelberger, G. Tóth, and H. Weinfurter, Experimental Entanglement of a Six-Photon Symmetric Dicke State, *Phys. Rev. Lett.* **103**, 020504 (2009).
- [9] W.-B. Gao, C.-Y. Lu, X.-C. Yao, P. Xu, O. Guhne, A. Goebel, Y.-A. Chen, C.-Z. Peng, Z.-B. Chen, and J.-W. Pan, Experimental demonstration of a hyper-entangled ten-qubit Schrodinger cat state, *Nat. Phys.* **6**, 331 (2010).
- [10] B. Vlastakis, G. Kirchmair, Z. Leghtas, S. E. Nigg, L. Frunzio, S. M. Girvin, M. Mirrahimi, M. H. Devoret, and R. J. Schoelkopf, Deterministically encoding quantum information using 100-photon Schrödinger cat states, *Science* **342**, 607 (2013).
- [11] A. Signoles, A. Facon, D. Grosso, I. Dotsenko, S. Haroche, J.-M. Raimond, M. Brune, and S. Gleyzes, Confined quantum Zeno dynamics of a watched atomic arrow, *Nat. Phys.* **10**, 715 (2014).
- [12] H. Jeong, M. S. Kim, T. C. Ralph, and B. S. Ham, Generation of macroscopic superposition states with small nonlinearity, *Phys. Rev. A* **70**, 061801 (2004).
- [13] S. Rebić, J. Twamley, and G. J. Milburn, Giant Kerr Nonlinearities in Circuit Quantum Electrodynamics, *Phys. Rev. Lett.* **103**, 150503 (2009).
- [14] X.-Y. Lü, Xin-You, W.-M. Zhang, S. Ashhab, Y. Wu, and F. Nori, Quantum-criticality-induced strong Kerr nonlinearities in optomechanical systems, *Sci. Rep.* **3**, 2943 (2013).
- [15] J. E. Debs, P. A. Altin, T. H. Barter, D. Döring, G. R. Dennis, G. McDonald, R. P. Anderson, J. D. Close, and N. P. Robins, Cold-atom gravimetry with a Bose-Einstein condensate, *Phys. Rev. A* **84**, 033610 (2011).
- [16] S. S. Szigeti, J. E. Debs, J. J. Hope, N. P. Robins, and J. D. Close, Why momentum width matters for atom interferometry with Bragg pulses, *New J. Phys.* **14**, 023009 (2012).
- [17] N. P. Robins, P. A. Altin, J. E. Debs, and J. D. Close, Atom lasers: Production, properties and prospects for precision inertial measurement, *Phys. Rep.* **529**, 265 (2013).
- [18] K. V. Kheruntsyan, M. K. Olsen, and P. D. Drummond, Einstein-Podolsky-Rosen Correlations via Dissociation of a Molecular Bose-Einstein Condensate, *Phys. Rev. Lett.* **95**, 150405 (2005).
- [19] H. Pu and P. Meystre, Creating macroscopic atomic Einstein-Podolsky-Rosen states from Bose-Einstein condensates, *Phys. Rev. Lett.* **85**, 3987 (2000).
- [20] C. Gross, H. Strobel, E. Nicklas, T. Zibold, N. Bar-Gill, G. Kurizki, and M. K. Oberthaler, Atomic homodyne detection of continuous-variable entangled twin-atom states, *Nature (London)* **480**, 219 (2011).
- [21] J.-C. Jaskula, M. Bonneau, G. B. Partridge, V. Krachmalnicoff, P. Deuar, K. V. Kheruntsyan, A. Aspect, D. Boiron, and C. I. Westbrook, Sub-Poissonian Number Differences in Four-Wave Mixing of Matter Waves, *Phys. Rev. Lett.* **105**, 190402 (2010).
- [22] R. Bucker, J. Grond, S. Manz, T. Berrada, T. Betz, C. Koller, U. Hohenester, T. Schumm, A. Perrin, and J. Schmiedmayer, Twin-atom beams, *Nat. Phys.* **7**, 608 (2011).
- [23] S. A. Haine and A. J. Ferris, Surpassing the standard quantum limit in an atom interferometer with four-mode entanglement produced from four-wave mixing, *Phys. Rev. A* **84**, 043624 (2011).
- [24] R. J. Lewis-Swan and K. V. Kheruntsyan, Proposal for demonstrating the Hong-Ou-Mandel effect with matter waves, *Nat. Commun.* **5**, 3752 (2014).
- [25] M. Kitagawa and M. Ueda, Squeezed spin states, *Phys. Rev. A* **47**, 5138 (1993).
- [26] A. Søndberg Sørensen, Bogoliubov theory of entanglement in a Bose-Einstein condensate, *Phys. Rev. A* **65**, 043610 (2002).
- [27] C. Gross, T. Zibold, E. Nicklas, J. Estève, and M. K. Oberthaler, Nonlinear atom interferometer surpasses classical precision limit, *Nature (London)* **464**, 1165 (2010).
- [28] M. F. Riedel, P. Böhi, Y. Li, T. W. Hänsch, A. Sinatra, and P. Treutlein, Atom-chip-based generation of entanglement for quantum metrology, *Nature (London)* **464**, 1170 (2010).
- [29] M. T. Johnsson and S. A. Haine, Generating Quadrature Squeezing in an Atom Laser Through Self-Interaction, *Phys. Rev. Lett.* **99**, 010401 (2007).
- [30] S. A. Haine and M. T. Johnsson, Dynamic scheme for generating number squeezing in Bose-Einstein condensates through nonlinear interactions, *Phys. Rev. A* **80**, 023611 (2009).
- [31] S. A. Haine, J. Lau, R. P. Anderson, and M. T. Johnsson, Self-induced spatial dynamics to enhance spin squeezing via one-axis twisting in a two-component Bose-Einstein condensate, *Phys. Rev. A* **90**, 023613 (2014).
- [32] H. Jing, J.-L. Chen, and M.-L. Ge, Quantum-dynamical theory for squeezing the output of a Bose-Einstein condensate, *Phys. Rev. A* **63**, 015601 (2000).
- [33] M. Fleischhauer and S. Gong, Stationary Source of Nonclassical or Entangled Atoms, *Phys. Rev. Lett.* **88**, 070404 (2002).
- [34] S. A. Haine and J. J. Hope, Outcoupling from a Bose-Einstein condensate with squeezed light to produce entangled-atom laser beams, *Phys. Rev. A* **72**, 033601 (2005).
- [35] S. A. Haine and J. J. Hope, A multi-mode model of a non-classical atom laser produced by outcoupling from a Bose-Einstein condensate with squeezed light, *Laser Phys. Lett.* **2**, 597 (2005).
- [36] S. A. Haine, M. K. Olsen, and J. J. Hope, Generating Controllable Atom-Light Entanglement with a Raman Atom Laser System, *Phys. Rev. Lett.* **96**, 133601 (2006).
- [37] K. Hammerer, A. S. Sørensen, and E. S. Polzik, Quantum interface between light and atomic ensembles, *Rev. Mod. Phys.* **82**, 1041 (2010).
- [38] S. A. Haine, Information-Recycling Beam Splitters for Quantum Enhanced Atom Interferometry, *Phys. Rev. Lett.* **110**, 053002 (2013).
- [39] S. S. Szigeti, B. Tonekaboni, W. Y. S. Lau, S. N. Hood, and S. A. Haine, Squeezed-light-enhanced atom interferometry below the standard quantum limit, *Phys. Rev. A* **90**, 063630 (2014).
- [40] B. Tonekaboni, S. A. Haine, and S. S. Szigeti, Heisenberg-limited metrology with a squeezed vacuum state, three-mode mixing, and information recycling, *Phys. Rev. A* **91**, 033616 (2015).

- [41] S. A. Haine, S. S. Szigeti, M. D. Lang, and C. M. Caves, Heisenberg-limited metrology with information recycling, *Phys. Rev. A* **91**, 041802 (2015).
- [42] S. L. Braunstein and C. M. Caves, Statistical Distance and the Geometry of Quantum States, *Phys. Rev. Lett.* **72**, 3439 (1994).
- [43] M. G. A. Paris, Quantum estimation for quantum technology, *Int. J. Quantum. Inform.* **7**, 125 (2009).
- [44] G. Tóth and I. Apellaniz, Quantum metrology from a quantum information science perspective, *J. Phys. A* **47**, 424006 (2014).
- [45] R. Demkowicz-Dobrzański, M. Jarzyna, J. Kołodyński, and E. Wolf, Quantum limits in optical interferometry, in *Progress in Optics*, Vol. 60 (Elsevier, Amsterdam, 2015), p. 345.
- [46] B. Yurke, S. L. McCall, and J. R. Klauder, SU(2) and SU(1,1) interferometers, *Phys. Rev. A* **33**, 4033 (1986).
- [47] F. T. Arecchi, E. Courtens, R. Gilmore, and H. Thomas, Atomic coherent states in quantum optics, *Phys. Rev. A* **6**, 2211 (1972).
- [48] G. S. Agarwal, State reconstruction for a collection of two-level systems, *Phys. Rev. A* **57**, 671 (1998).
- [49] D. Mondal, Generalized Fubini-study metric and Fisher information metric, [arXiv:1503.04146](https://arxiv.org/abs/1503.04146).
- [50] L. Pezzé and A. Smerzi, Entanglement, Nonlinear Dynamics, and the Heisenberg Limit, *Phys. Rev. Lett.* **102**, 100401 (2009).
- [51] If we restricted measurements to system A , then this is equivalent to tracing out system B , in which case the QFI is \mathcal{F}_A .
- [52] W. Wasilewski, K. Jensen, H. Krauter, J. J. Renema, M. V. Balabas, and E. S. Polzik, Quantum Noise Limited and Entanglement-Assisted Magnetometry, *Phys. Rev. Lett.* **104**, 133601 (2010).
- [53] M. H. Schleier-Smith, I. D. Leroux, and V. Vuletić, Squeezing the collective spin of a dilute atomic ensemble by cavity feedback, *Phys. Rev. A* **81**, 021804 (2010).
- [54] Z. Chen, J. G. Bohnet, S. R. Sankar, J. Dai, and J. K. Thompson, Conditional Spin Squeezing of a Large Ensemble via the Vacuum Rabi Splitting, *Phys. Rev. Lett.* **106**, 133601 (2011).
- [55] S. S. Szigeti, M. R. Hush, A. R. R. Carvalho, and J. J. Hope, Continuous measurement feedback control of a Bose-Einstein condensate using phase-contrast imaging, *Phys. Rev. A* **80**, 013614 (2009).
- [56] S. S. Szigeti, M. R. Hush, A. R. R. Carvalho, and J. J. Hope, Feedback control of an interacting Bose-Einstein condensate using phase-contrast imaging, *Phys. Rev. A* **82**, 043632 (2010).
- [57] T. Vanderbruggen, S. Bernon, A. Bertoldi, A. Landragin, and P. Bouyer, Spin-squeezing and Dicke-state preparation by heterodyne measurement, *Phys. Rev. A* **83**, 013821 (2011).
- [58] S. Bernon, T. Vanderbruggen, R. Kohlhaas, A. Bertoldi, A. Landragin, and P. Bouyer, Heterodyne nondemolition measurements on cold atomic samples: Towards the preparation of nonclassical states for atom interferometry, *New J. Phys.* **13**, 065021 (2011).
- [59] N. Brahm, T. Botter, S. Schreppler, D. W. C. Brooks, and D. M. Stamper-Kurn, Optical Detection of the Quantization of Collective Atomic Motion, *Phys. Rev. Lett.* **108**, 133601 (2012).
- [60] J. G. Bohnet, K. C. Cox, M. A. Norcia, J. M. Weiner, Z. Chen, and J. K. Thompson, Reduced spin measurement back-action for a phase sensitivity ten times beyond the standard quantum limit, *Nat. Photon.* **8**, 731 (2014).
- [61] K. C. Cox, J. M. Weiner, G. P. Greve, and J. K. Thompson, Generating entanglement between atomic spins with low-noise probing of an optical cavity, [arXiv:1504.05160v1](https://arxiv.org/abs/1504.05160v1).
- [62] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R. S. Huang, J. Majer, S. Kumar, S. M. Girvin, and R. J. Schoelkopf, Strong coupling of a single photon to a superconducting qubit using circuit quantum electrodynamics, *Nature (London)* **431**, 162 (2004).
- [63] D. I. Schuster, A. Wallraff, A. Blais, L. Frunzio, R.-S. Huang, J. Majer, S. M. Girvin, and R. J. Schoelkopf, ac Stark Shift and Dephasing of a Superconducting Qubit Strongly Coupled to a Cavity Field, *Phys. Rev. Lett.* **94**, 123602 (2005).
- [64] J. A. Haigh, N. J. Lambert, A. C. Doherty, and A. J. Ferguson, Dispersive readout of ferromagnetic resonance for strongly coupled magnons and microwave photons, *Phys. Rev. B* **91**, 104410 (2015).
- [65] D. F. Walls and G. J. Milburn, *Quantum Optics*, 2nd ed. (Springer-Verlag, Berlin, 2008).
- [66] M. G. Moore and P. Meystre, Generating Entangled Atom-Photon Pairs from Bose-Einstein Condensates, *Phys. Rev. Lett.* **85**, 5026 (2000).
- [67] P. Bouyer and M. A. Kasevich, Heisenberg-limited spectroscopy with degenerate Bose-Einstein gases, *Phys. Rev. A* **56**, R1083 (1997).
- [68] T. Kim, O. Pfister, M. J. Holland, J. Noh, and J. L. Hall, Influence of decorrelation on Heisenberg-limited interferometry with quantum correlated photons, *Phys. Rev. A* **57**, 4004 (1998).
- [69] J. J. Bollinger, Wayne M. Itano, D. J. Wineland, and D. J. Heinzen, Optimal frequency measurements with maximally correlated states, *Phys. Rev. A* **54**, R4649 (1996).
- [70] H. A. Bachor and T. C. Ralph, *A Guide to Experiments in Quantum Optics*, 2nd ed. (Wiley, New York, 2004).
- [71] I. Bloch, J. Dalibard, and W. Zwerger, Many-body physics with ultracold gases, *Rev. Mod. Phys.* **80**, 885 (2008).
- [72] A. D. Cronin, J. Schmiedmayer, and D. E. Pritchard, Optics and interferometry with atoms and molecules, *Rev. Mod. Phys.* **81**, 1051 (2009).
- [73] L. Li, Y. O. Dudin, and A. Kuzmich, Entanglement between light and an optical atomic excitation, *Nature (London)* **498**, 466 (2013).
- [74] H. Paik, D. I. Schuster, Lev S. Bishop, G. Kirchmair, G. Catelani, A. P. Sears, B. R. Johnson, M. J. Reagor, L. Frunzio, L. I. Glazman, S. M. Girvin, M. H. Devoret, and R. J. Schoelkopf, Observation of High Coherence in Josephson Junction Qubits Measured in a Three-Dimensional Circuit QED Architecture, *Phys. Rev. Lett.* **107**, 240501 (2011).
- [75] Z.-L. Xiang, S. Ashhab, J. Q. You, and F. Nori, Hybrid quantum circuits: Superconducting circuits interacting with other quantum systems, *Rev. Mod. Phys.* **85**, 623 (2013).
- [76] M. Aspelmeyer, T. J. Kippenberg, and F. Marquardt, Cavity optomechanics, *Rev. Mod. Phys.* **86**, 1391 (2014).
- [77] D. Leibfried, R. Blatt, C. Monroe, and D. Wineland, Quantum dynamics of single trapped ions, *Rev. Mod. Phys.* **75**, 281 (2003).
- [78] D. J. Wineland, Nobel lecture: Superposition, entanglement, and raising Schrödinger's cat, *Rev. Mod. Phys.* **85**, 1103 (2013).
- [79] A. Stute, B. Casabone, B. Brandstatter, K. Friebe, T. E. Northup, and R. Blatt, Quantum-state transfer from an ion to a photon, *Nat. Photon* **7**, 219 (2013).
- [80] M. A. Rowe, D. Kielpinski, V. Meyer, C. A. Sackett, W. M. Itano, C. Monroe, and D. J. Wineland, Experimental violation of a Bell's inequality with efficient detection, *Nature (London)* **409**, 791 (2001).

- [81] D. H. Smith, G. Gillett, M. P. de Almeida, C. Branciard, A. Fedrizzi, T. J. Weinhold, A. Lita, B. Calkins, T. Gerrits, H. M. Wiseman, S. W. Nam, and A. G. White, Conclusive quantum steering with superconducting transition-edge sensors, *Nat. Commun.* **3**, 625 (2012).
- [82] C. Guerlin, J. Bernu, S. Deleglise, C. Sayrin, S. Gleyzes, S. Kuhr, M. Brune, J.-M. Raimond, and S. Haroche, Progressive field-state collapse and quantum non-demolition photon counting, *Nature (London)* **448**, 889 (2007).
- [83] J. P. Groen, D. Ristè, L. Tornberg, J. Cramer, P. C. de Groot, T. Picot, G. Johansson, and L. DiCarlo, Partial-Measurement Backaction and Nonclassical Weak Values in a Superconducting Circuit, *Phys. Rev. Lett.* **111**, 090506 (2013).
- [84] K. W. Murch, S. J. Weber, C. Macklin, and I. Siddiqi, Observing single quantum trajectories of a superconducting quantum bit, *Nature (London)* **502**, 211 (2013).
- [85] J. D. Cohen, S. M. Meenehan, and O. Painter, Optical coupling to nanoscale optomechanical cavities for near quantum-limited motion transduction, *Opt. Express* **21**, 11227 (2013).
- [86] E. M. Kessler, I. Lovchinsky, A. O. Sushkov, and M. D. Lukin, Quantum Error Correction for Metrology, *Phys. Rev. Lett.* **112**, 150802 (2014).
- [87] S. S. Szigeti, A. R. R. Carvalho, J. G. Morley, and M. R. Hush, Ignorance is Bliss: General and Robust Cancellation of Decoherence via No-Knowledge Quantum Feedback, *Phys. Rev. Lett.* **113**, 020407 (2014).