

The global atlas of podoconiosis

Article (Published Version)

Deribe, Kebede, Cano, Jorge, Newport, Melanie J, Pullan, Rachel L, Noor, Abdisalan M, Enquesselassie, Fikre, Murray, Christopher J L, Hay, Simon I, Brooker, Simon J and Davey, Gail (2017) The global atlas of podoconiosis. *Lancet Global Health*, 5 (5). e477-e479. ISSN 2214-109X

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/67394/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



The global atlas of podoconiosis

The world stands on the edge of an historic public health success with the imminent eradication of dracunculiasis (guinea-worm disease) and polio. Since the World Health Assembly called for the eradication of dracunculiasis in 1986¹ and poliomyelitis in 1988,² astonishing progress has been made. In 2016, only 25 human cases of dracunculiasis were reported from three countries,³ transmission of wild poliovirus was found in only three countries, and 37 cases of polio were reported worldwide.⁴ In addition to these achievements, there has been progress in the elimination of the little-known disease podoconiosis (endemic non-filarial elephantiasis).⁵ Essential to this elimination effort is the need to understand the geography of the disease to identify potentially at-risk areas and set a benchmark for evaluating elimination efforts.⁶

The global burden of podoconiosis is not at present clearly defined, although available estimates suggest that globally there are 4 million people with podoconiosis,⁷ mainly in tropical countries of Africa, Central and South America, and southeast Asia. Tropical African countries bear the highest disease burden. A literature search identifies 32 countries as being either known or suspected to be endemic for podoconiosis. In Africa, the disease has been reported in Angola, Burundi, Cameroon, Cape Verde, Chad, Democratic Republic of Congo, Equatorial Guinea, Ethiopia, Kenya, Madagascar, Mozambique, Niger, Nigeria, Rwanda, São Tomé and Príncipe, Sudan, Tanzania, and Uganda. Podoconiosis has been reported in the Latin American highlands in Brazil, Colombia, Costa Rica, Ecuador, El Salvador, French Guiana, Guatemala, Honduras, Mexico, Peru, and Suriname. In Asia, although filarial elephantiasis predominates in India, podoconiosis has been reported in the northwestern part of the country, as well as in Sri Lanka and Indonesia.⁸ However, under-reporting of podoconiosis is possible because of diagnostic challenges and a low index of suspicion. Podoconiosis is an environment-related disease, caused by long-term barefoot exposure to red clay soil.^{6,9,10} Therefore, environmental factors, such as soil, and other climatic factors that affect the generation of soil, can help predict the occurrence of podoconiosis. An individual's vulnerability to the disease is also aggravated by poverty and insufficient access to water for foot hygiene.¹¹

We have received funding from the Wellcome Trust¹² to develop a global atlas of podoconiosis. We aim to advance new knowledge on the geographical distribution and spatial epidemiology of the disease. A first step in this work will be to establish the geographical absence of disease by applying an evidence consensus approach (thorough literature searches and contacting ministries of health).^{13,14} The project will also use environmental predictors to determine environmental suitability for the occurrence of podoconiosis. In our previous work in Ethiopia we have identified important covariates that drive the spatial distribution of podoconiosis. Using boosted regression tree modelling, we mapped the environmental limits of podoconiosis, with high accuracy for determining the presence or absence of podoconiosis.⁶ Once validated in other countries this model will have the potential to define the environmental limits of podoconiosis globally, thus targeting survey efforts to high-priority countries.

Population-based surveys are important sources of data for mapping of podoconiosis.¹⁵ Surveys informed by podoconiosis risk stratification will also be important. Such surveys will provide statistically powered and spatially representative sampling, which will capture environmental risk drivers of podoconiosis within a

Panel: Standard podoconiosis survey

Podoconiosis surveys are done in individuals aged 15 years or older who have lived in the implementation unit (administrative units used as the basis for making decisions about morbidity management) for more than 10 years.¹⁵ Podoconiosis is a clinical diagnosis based on exclusion of other potential causes of lymphoedema and disease-specific tests, such as the Alere Filariasis Test Strip and Wb123 test, to exclude lymphoedema due to lymphatic filariasis. For the surveyed cluster or districts, the following data are required:

- The name of the site (district or village names)
- Region and district where the survey was done
- Coordinates (longitude and latitude) of community or cluster (if available)
- Survey methods and population sampling
- Year of survey
- Age range
- Number of individuals examined
- Number of people with lymphoedema
- Number of people diagnosed with podoconiosis
- Disease stage for podoconiosis¹⁶

For the integrated mapping of NTDs see <http://aiminitiative.org/>

country. A recommended survey of podoconiosis is done among individuals aged 15 years or older (podoconiosis commonly occurs after the age of 15 years), with a clear diagnostic algorithm using history, physical diagnosis, and disease-specific tests (panel).¹⁵ The usefulness of the data can be enhanced by Bayesian modelling to identify risk factors at different scales and to predict the prevalence and burden of podoconiosis in unsampled locations. Such a Bayesian framework has been used to effectively map the global distribution of several infectious diseases, including malaria, dengue virus infection, and soil-transmitted helminth infection.^{14,17}

Data collected through dedicated epidemiological surveys can be expensive and inefficient, however, especially for countries with low disease prevalence. To reduce reporting costs, suspected endemic countries can also collect information about podoconiosis through routine surveillance systems. For example, since Ethiopia included podoconiosis into the national health management information system in 2013, an increased number of cases have been reported. The use of community health workers to identify cases of podoconiosis, which are subsequently validated by experienced health workers, has been piloted with good success.^{18,19} There is also opportunity for the integrated reporting of morbidity cases due to other neglected tropical diseases, such as leprosy, Buruli ulcer, and trichiasis, and subsequent management through the primary health-care system. Clinical diagnosis of podoconiosis based on pathognomonic signs and symptoms has been found to be workable in the regular health system. Case definitions for podoconiosis have been developed and can be used for case identification through routine surveillance.^{5,20}

Over the span of 5 years, the global atlas of podoconiosis will define the epidemiology and distribution of podoconiosis globally. Through collection and collation of the available evidence, generation of new epidemiological data, and the currently available state of the art geostatistical and machine learning approaches, the global limits of podoconiosis will be defined and the population at risk and burden of the disease estimated for the first time. Further collaborations from endemic countries, partners, and WHO country offices are crucial to advance the project goals. Additionally, we are exploring strategies to integrate podoconiosis mapping with

ongoing mapping of other neglected tropical diseases. The atlas will provide vital evidence of the geographical distribution and burden of podoconiosis globally, and provide an important basis for expanding prevention and treatment services on the path towards a world without podoconiosis.

**Kebede Deribe, Jorge Cano, Melanie J Newport, Rachel L Pullan, Abdisalan M Noor, Fikre Enquselassie, Christopher J L Murray, Simon I Hay, Simon J Brooker, Gail Davey*

Wellcome Trust Brighton and Sussex Centre for Global Health Research, Brighton and Sussex Medical School, Brighton, BN1 9PX, UK (KD, MJN, GD); School of Public Health, Addis Ababa University, Addis Ababa, Ethiopia (KD, FE); Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, UK (JC, RLP); Kenya Medical Research Institute-Wellcome Trust Research Programme, Nairobi, Kenya (AMN); Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, USA (CJLM, SIH); Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK (SIH); and Bill and Melinda Gates Foundation, Seattle, WA, USA (SJB) kebededeka@yahoo.com

KD is funded by a Wellcome Trust Intermediate Fellowship in Public Health and Tropical Medicine (201900). SIH is funded by a Senior Research Fellowship from the Wellcome Trust (095066), and grants from the Bill and Melinda Gates Foundation (OPP1119467, OPP1093011, OPP1106023, and OPP1132415). We declare no competing interests.

Copyright © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY license.

- 1 World Health Assembly. Resolution WHA 39.21: elimination of dracunculiasis. Geneva: World Health Organization, 1986. http://www.who.int/neglected_diseases/mediacentre/WHA_39_21_Eng.pdf (accessed Feb 14, 2017).
- 2 World Health Assembly. Resolution WHA 41.28: global eradication of poliomyelitis by the year 2000. Geneva: World Health Organization, 1988. <http://www.who.int/ihr/polioresolution4128en.pdf> (accessed Feb 14, 2017).
- 3 WHO. Dracunculiasis eradication: Mali reports zero cases in 2016. Geneva: World Health Organization, 2017. http://www.who.int/neglected_diseases/news/Mali_reports_zero_cases_in_2016/en/ (accessed Feb 14, 2017).
- 4 WHO. Executive board EB140/13. Poliomyelitis. Geneva: World Health Organization, 2016. http://polioeradication.org/wp-content/uploads/2016/07/EB140_13-en.pdf (accessed Feb 14, 2017).
- 5 Deribe K, Wanji S, Shafi O, et al. The feasibility of eliminating podoconiosis. *Bull World Health Organ* 2015; **93**: 712–18.
- 6 Deribe K, Cano J, Newport MJ, et al. Mapping and modelling the geographical distribution and environmental limits of podoconiosis in Ethiopia. *PLoS Negl Trop Dis* 2015; **9**: e0003946.
- 7 Tekola Ayele F, Adeyemo A, Finan C, et al. HLA class II locus and susceptibility to podoconiosis. *N Engl J Med* 2012; **366**: 1200–08.
- 8 Price EW. Podoconiosis: non-filarial elephantiasis. Oxford: Oxford Medical Publications, 1990.
- 9 Molla YB, Wardrop NA, Le Blond JS, et al. Modelling environmental factors correlated with podoconiosis. *Int J Health Geogr* 2014; **13**: 24.
- 10 Price EW, Bailey D. Environmental factors in the etiology of endemic elephantiasis of the lower legs in tropical Africa. *Trop Geogr Med* 1984; **36**: 1–5.
- 11 Deribe K, Brooker SJ, Pullan RL, et al. Epidemiology and individual, household and geographical risk factors of podoconiosis in Ethiopia: results from the first nationwide mapping. *Am J Trop Med Hyg* 2015; **92**: 148–58.

- 12 Wellcome Trust. Intermediate fellowships in public health and tropical medicine. 2016. <https://wellcome.ac.uk/what-we-do/directories/intermediate-fellowships-public-health-tropical-medicine-people-funded> (accessed Feb 24, 2017).
- 13 Brady OJ, Gething PW, Bhatt S, et al. Refining the global spatial limits of dengue virus transmission by evidence-based consensus. *PLoS Negl Trop Dis* 2012; **6**: e1760.
- 14 Bhatt S, Gething PW, Brady OJ, et al. The global distribution and burden of dengue. *Nature* 2013; **496**: 504–07.
- 15 Sime H, Deribe K, Assefa A, et al. Integrated mapping of lymphatic filariasis and podoconiosis: lessons learnt from Ethiopia. *Parasit Vectors* 2014; **7**: 397.
- 16 Tekola F, Ayele Z, Mariam DH, Fuller C, Davey G. Development and testing of a de novo clinical staging system for podoconiosis (endemic non-filarial elephantiasis). *Trop Med Int Health* 2008; **13**: 1277–83.
- 17 Pullan RL, Smith JL, Jasrasaria R, Brooker SJ. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasit Vectors* 2014; **7**: 37.
- 18 Alemu G, Tekola Ayele F, Daniel T, Ahrens C, Davey G. Burden of podoconiosis in poor rural communities in Gulliso woreda, West Ethiopia. *PLoS Negl Trop Dis* 2011; **5**: e1184.
- 19 Desta K, Ashine M, Davey G. Predictive value of clinical assessment of patients with podoconiosis in an endemic community setting. *Trans R Soc Trop Med Hyg* 2007; **101**: 621–23.
- 20 Deribe K, Wanji S, Shafi O, Muheki E, Umulisa I, Davey G. Measuring elimination of podoconiosis, endemicity classifications, case definition and targets: an international Delphi exercise. *Int Health* 2015; **7**: 306–16.